

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 July 2004 (15.07.2004)

PCT

(10) International Publication Number
WO 2004/058309 A1

(51) International Patent Classification⁷: **A61K 49/00**,
C07K 14/525

MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2003/040979

(22) International Filing Date:
22 December 2003 (22.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/435,262 23 December 2002 (23.12.2002) US
60/467,198 2 May 2003 (02.05.2003) US

(71) Applicant (for all designated States except US): **HUMAN GENOME SCIENCES, INC.** [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PARMELEE, David** [US/US]; 11512 Parkedge Drive, Rockville, MD 20852 (US). **YEH, Ren-Hwa** [CN/US]; 13834 Lullaby Road, Germantown, MD 20874 (US). **GALPERINA, Olga** [US/US]; 14029 Saddlevue Drive, North Potomac, MD 20878 (US). **HILBERT, David** [US/US]; 8501 Meadowlark Lane, Bethesda, MD 20817 (US). **ROSEN, Craig, A.** [US/US]; 22400 Rolling Hill Lane, Laytonsville, MD 20882 (US).

(74) Agents: **STEFFE, ERIC, K.** et al.; Sterne, Kessler, Goldstein & Fox, P.L.L.C., 1100 New York Avenue, N.W., Washington, DC 20005 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,

(84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEUTROKINE-ALPHA CONJUGATE, NEUTROKINE-ALPHA COMPLEX, AND USES THEREOF

(57) Abstract: The present invention is directed to a neutrokin- α conjugate, wherein said neutrokin- α conjugate comprises a neutrokin- α protein and a chelator. The present invention is also directed to a neutrokin- α complex, wherein said neutrokin- α complex comprises a neutrokin- α conjugate and a metal ion wherein said metal ion is noncovalently associated with the chelator of said conjugate. The present invention is further directed to compositions, specifically pharmaceutical and diagnostic compositions, comprising a neutrokin- α conjugate or complex as described herein. Therapeutic and diagnostic uses of said conjugate, complex, and compositions are described.



WO 2004/058309 A1

NEUTROKINE-ALPHA CONJUGATE, NEUTROKINE-ALPHA COMPLEX, AND USES THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention is directed to a modified Neutrokin- α protein that can be labeled with a metal ion, such as a radionuclide, and is useful for the treatment, diagnosis, and imaging of certain conditions such as cancer and autoimmune disease.

Background Art

[0002] Neutrokin- α (also known as BLySTM protein (B-Lymphocyte Stimulator); also known as, *inter alia*, TALL-1, THANK, and BAFF) is a member of the tumor necrosis factor (TNF) superfamily that induces B cell proliferation and immunoglobulin secretion (Moore *et al.*, *Science* 285:260-263 (1999)). Neutrokin- α also appears to be a key regulator of peripheral B cell populations *in vivo* due to the role Neutrokin- α plays in regulating B cell survival (Mackay *et al.*, *J. Exp. Med.* 190:1697-1710 (1999); Do *et al.*, *J. Exp. Med.* 192:953-964 (2000); and Hsu *et al.*, *J. Immunol.* 168:5993-6 (2002)). Like other members of the TNF family, Neutrokin- α is a type-II membrane protein that can be cleaved at the cell surface to form a soluble protein (Mariani *et al.*, *J. Cell Biol.* 137:221-229 (1997)). Neutrokin- α , like other members of the TNF ligand family described to date, forms biologically active trimers.

[0003] Like other members of the TNF family, Neutrokin- α is a ligand that interacts with several receptors. Neutrokin- α was initially shown to interact with TACI (trans-membrane activator and CAML interactor) and BCMA (B cell maturation antigen) (Gross *et al.*, *Nature* 404:995-999 (2000)). Both receptors were found to bind APRIL as well (Marsters *et al.*, *Curr. Biol.* 10:785-788 (2000); Wu *et al.*, *J. Biol. Chem.* 275:35478-35485 (2000)),

- 2 -

APRIL being the TNF ligand that has the highest degree of sequence homology with Neutrokin- α . Most recently, a third receptor, termed BAFF-R, has been identified. This receptor apparently does not interact with APRIL or any known TNF ligand other than Neutrokin- α (Thompson *et al.*, *Science* 293:2108-2111 (2001)).

[0004] Neutrokin- α receptor expression, defined functionally by the binding of biotinylated Neutrokin- α to cells, is found predominantly on B cells (see, *e.g.*, Moore *et al.*, *Science* 285:260-263 (1999), although TACI has also been reported to be expressed on activated T cells (Wang *et al.*, *Nature Immunol.* 2:577-8 (2001). The expression of Neutrokin- α is observed on normal B cells as well as on tumorous B cells, including Non-Hodgkin's lymphoma cells and Chronic Lymphocytic Leukemia cells. Furthermore, Neutrokin- α receptor expression is not observed in pre-B cells; rather, Neutrokin- α receptor expression becomes observable at the same stage in B cell development when surface Ig expression becomes apparent (see, *e.g.*, Hsu *et al.*, *J. Immunol.* 168:5993-5996 (2002).

[0005] The restricted expression profile of Neutrokin- α receptors on lymphoid cells, and predominantly on B cells, makes Neutrokin- α an attractive vehicle for targeting therapies to lymphocytes, and to B lineage cells in particular. Neutrokin- α thus may be used to treat and diagnose diseases and disorders of the immune system, particularly those associated with aberrant B cells numbers or function, including for example, autoimmune diseases, B cell cancers and inflammation.

[0006] Autoimmune diseases are characterized, in part, by a failure of the immune system to distinguish the body's own cells and tissues from those of pathogens. B cells that produce antibodies that recognize parts of the normal body (autoantibodies) are characteristic of many autoimmune diseases. Systemic lupus erythematosus, rheumatoid arthritis, Sjögren's syndrome, Wegener's granulomatous, myasthenia gravis, multiple sclerosis, diabetes, and some forms of asthma are all examples of autoimmune diseases that are associated with autoantibodies. Thus, an agent that inhibits the proliferation,

- 3 -

differentiation or survival of B cells, *e.g.*, a drug comprising Neutrokin- α linked to a source of radiation, could be used to treat or prevent diseases such as systemic lupus erythematosus, rheumatoid arthritis, Sjögren's syndrome, Wegener's granulomatous, myasthenia gravis, multiple sclerosis, diabetes and some forms of asthma. Such an agent would also be useful in the treatment of cancers, particularly B cell cancers or cancer cells in close proximity to B cells.

[0007] Moreover, the development of a Neutrokin- α protein associated with a metal ion would be useful in diagnostic processes. Such a protein conjugate could be linked to a metal ion which could be visualized using any number of known techniques including single photon emission computed tomography (SPECT), positron emission tomography (PET), and magnetic resonance imaging (MRI). A Neutrokin- α protein associated with a metal ion could further be used, *inter alia*, as an agent to inhibit the proliferation of or kill B-cells. Thus, a Neutrokin- α protein associated with a metal ion would be useful in the treatment of diseases caused by or involving aberrant B cell proliferation, activation or survival including, but not limited to, B cell cancers, autoimmune diseases and inflammation.

[0008] The use of a chelator conjugated to a protein for labeling said protein with radiometals allows one to label the same protein-chelator conjugate with different radiometals providing a number of choices of half-life and types of emissions for various medical applications in both diagnosis and therapy. Radiometals also offer significant advantages over iodine when used to label proteins. Radiometal labeling, for example, avoids the deleterious effects of oxidation experienced in direct iodination reactions. Labeling with metals can also overcome problems of *in vivo* deiodination by tumor and normal tissues, particularly when using rapidly internalized proteins.

BRIEF SUMMARY OF THE INVENTION

[0009] The present invention provides compositions useful in the treatment and diagnosis of diseases and disorders associated with cells bearing

- 4 -

Neutrokin- α receptors on their surface. As Neutrokin- α receptors are found predominantly on cells of lymphoid origin, and particularly on B cells, in one embodiment the present invention provides compositions useful in the treatment and diagnosis of diseases and disorders associated with aberrant lymphocyte, and particularly B lymphocyte, number or function. In specific embodiments, the present invention provides compositions useful in the treatment and diagnosis of autoimmune diseases with a humoral autoimmune component including, but not limited to, Systemic lupus Erythematosus, rheumatoid arthritis, Sjögren's syndrome, Wegener's granulomatous, myasthenia gravis, multiple sclerosis, diabetes, and autoimmune forms of asthma. In other specific embodiments, the present invention provides compositions useful in the treatment and diagnosis of cancers of cells bearing neutrokin- α receptors including B cell cancers such as Non-Hodgkins' lymphoma, chronic lymphocytic leukemia, and multiple myeloma, or cancers of other cell types that express Neutrokin- α receptors, (e.g., epithelial cells such as epithelial cells in the lung or colon.). In other specific embodiments, the present invention provides compositions useful in the treatment and diagnosis of cancers of cells that are in close proximity to cells expressing Neutrokin- α receptors, for example, cancers that have metastasized through the lymphatic system.

[0010] In one embodiment, the present invention provides a Neutrokin- α conjugate comprising a Neutrokin- α protein and a chelator. In another embodiment, the present invention provides a Neutrokin- α complex comprising a Neutrokin- α conjugate and a metal ion. In specific embodiments, a composition of the invention comprises a Neutrokin- α conjugate or a Neutrokin- α complex in an acceptable carrier.

[0011] The present invention also provides methods of preparing Neutrokin- α conjugates and Neutrokin- α complexes. In one embodiment, a Neutrokin- α conjugate is prepared by reacting a Neutrokin- α protein with a chelator. In another embodiment, a Neutrokin- α complex is prepared by reacting a Neutrokin- α conjugate with a metal ion. In

- 5 -

another embodiment, a Neutrokin- α complex is prepared by reacting a Neutrokin- α protein with a chelator-metal ion complex.

[0012] In preferred embodiments, the Neutrokin- α protein component of a Neutrokin- α conjugate or Neutrokin- α complex is a protein consisting of a trimer of three identical subunits, each subunit comprising, or alternatively consisting of the protein of SEQ ID NO:3, namely amino acids 134-285 of the full-length Neutrokin- α protein which is shown in SEQ ID NO:2. In other embodiments, the Neutrokin- α protein component of a Neutrokin- α conjugate or Neutrokin- α complex, comprises or alternatively consists of the Neutrokin- α protein of SEQ ID NO:2, or fragments or variants thereof, *e.g.*, amino acids 134-285 of SEQ ID NO:2.

[0013] In preferred embodiments, the chelator component of a Neutrokin- α conjugate or Neutrokin- α complex is α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7,10-tetraacetic acid. In other embodiments, the chelator of a Neutrokin- α conjugate or Neutrokin- α complex has a chemical formula according to any one of the formulas selected from the group consisting of Formula I, II, III, or IV described below.

[0014] In preferred embodiments, the metal ion component is a radionuclide that emits gamma rays, positrons, x-rays, fluorescence, beta particles, alpha particles, or auger electrons. In other preferred embodiments, the metal ion component is a radionuclide which is an electron or neutron-capturing agent.

[0015] In preferred embodiments, the metal ion component of Neutrokin- α complex or chelator-metal-ion complex is yttrium-90 (^{90}Y). In other embodiments, the metal ion component of a Neutrokin- α complex or chelator-metal-ion complex is indium-111 (^{111}In), leuticium-177 (^{177}Lu), holmium-166 (^{166}Ho), bismuth-215 (^{215}Bi), and actinium-225 (^{225}Ac).

[0016] The present invention further encompasses methods and compositions for killing cells bearing Neutrokin- α receptors and/or cells in close proximity to cells bearing Neutrokin- α receptors, comprising, or alternatively consisting of, contacting a Neutrokin- α conjugate and/or a

- 6 -

Neutrokin-alpha complex of the invention with cells bearing Neutrokin-alpha receptors. In preferred embodiments, the cells bearing Neutrokin-alpha receptors are B cells.

[0017] The present invention further encompasses methods and compositions for killing cells bearing Neutrokin-alpha receptors or cells in close proximity to cells bearing a Neutrokin-alpha receptor, comprising, or alternatively consisting of, administering to an animal in which such killing is desired, a Neutrokin-alpha conjugate and/or a Neutrokin-alpha complex in an amount effective to kill cells bearing Neutrokin-alpha receptors and/or cells in close proximity to cells bearing a Neutrokin-alpha receptor. In preferred embodiments, the cells bearing Neutrokin-alpha receptors are B cells.

[0018] The present invention also provides methods of using the compositions of the invention for the diagnosis and/or treatment of B-cell cancers. Specifically contemplated is the use of Neutrokin-alpha for the diagnosis and/or treatment of non-Hodgkin's lymphoma, multiple myeloma, chronic lymphocytic leukemia (CLL), acute lymphocytic leukemia (ALL), Burkitt's lymphoma, and/or Epstein Barr-Virus (EBV) transformed B cell cancers.

[0019] The present invention also provides methods of using the compositions of the invention for the diagnosis and/or treatment of autoimmune disease. In specific embodiments, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes are used for the diagnosis and/or treatment of systemic lupus erythematosus (SLE), rheumatoid arthritis (RA), Sjögren's Syndrome, idiopathic thrombocytic purpura (ITP) hemolytic anemia, myasthenia gravis, and/or IgA nephropathy.

[0020] In specific embodiments, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes are used for the diagnosis and/or treatment of atherosclerosis.

[0021] The present invention also provides pharmaceutical kits for the preparation of a Neutrokin-alpha conjugate or Neutrokin-alpha complex.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES

[0022] **Figure 1** shows a proposed scheme of a method of preparing a Neutrokin- α complex.

[0023] **Figure 2.** Effect of ^{131}I -labeled Neutrokin- α (lot TX1) on the survival of BCL1 tumor-bearing BALB/c mice. Survival curve expressed in terms of survival probability vs. time. Day 0 is the first day of tumor cell injection. Differences among the treatment groups were analyzed using the Log Rank Test for equality. Treatment with ^{131}I -labeled Neutrokin- α (LR131 in figure) at doses of either 11.9 or 15.3 mCi/kg (red and blue dotted lines, respectively) significantly prolonged survival ($p = 0.0162$ and $p = 0.0052$, respectively) compared with vehicle-treated controls (black solid line). In the group of mice that did not have BCL1 tumors but did receive 15.3 mCi/kg of ^{131}I -labeled Neutrokin- α , 10% of the mice died (yellow dashed line).

[0024] **Figure 3.** Effect of ^{131}I -labeled Neutrokin- α (lot TX2) on the survival of BCL1 tumor-bearing BALB/c mice. Survival curve expressed in terms of survival probability vs. time. Day 0 is the first day of tumor cell injection. Differences among the treatment groups were analyzed using the Log Rank Test for equality. Treatment with ^{131}I -labeled Neutrokin- α (LR131 in figure) at a dose of 17.5 mCi/kg (dashed line) significantly prolonged survival ($p = 0.0348$) compared with the vehicle-treated controls (black solid line). In the group of mice that did not have BCL1 tumors but did receive ^{131}I -labeled Neutrokin- α , 12.5% of the mice died (dotted line).

[0025] **Figure 4.** Effect of ^{131}I -labeled Neutrokin- α (lot TX3) on the survival of BCL1 tumor-bearing BALB/c mice. Survival curve expressed in terms of survival probability vs. time. Day 0 is the day the tumor cells were injected. Differences among the treatment groups were analyzed using the Log Rank Test for equality. Treatment with ^{131}I -labeled Neutrokin- α (LR131 in figure) at a dose of 37.7 mCi/kg (dashed line) significantly prolonged survival ($p = 0.0212$) compared to the vehicle-treated controls

- 8 -

(solid line). In the group of mice that did not have BCL1 tumors but did receive ^{131}I -labeled Neutrokin-alpha, 12.5% of the mice died (dotted line).

[0026] **Figure 5** shows a plasmid map of the pML124 vector. The sequence of this vector is shown in SEQ ID NO:14.

[0027] **Figure 6** shows a plasmid map of the pML124 vector containing the MBPss-Neutrokin-alpha fusion. The sequence of this vector is shown in SEQ ID NO:15.

DETAILED DESCRIPTION OF THE INVENTION

[0028] The restricted expression profile of Neutrokin-alpha receptors on lymphoid cells, and predominantly on B cells, makes Neutrokin-alpha an attractive vehicle for targeting therapies to lymphocytes, and to B lineage cells in particular. To date, three receptors for Neutrokin-alpha have been identified; namely BAFF-R (Locus ID: 115650, Refseq No. NM_052945, SEQ ID NO:5) TACI (Locus ID: 23495, Refseq No. NM_012452, SEQ ID NO:7), and BCMA (Locus ID: 608, Refseq No. NM_001192, SEQ ID NO:9). LocusID numbers refer to the NCBI Locus Link website (<http://www.ncbi.nlm.nih.gov/LocusLink/list.cgi>) and the "NM" Nos. are GenBank Refseq Nos. for the mRNAs encoding these receptors. Nucleotide sequences encoding these receptors are given in SEQ ID NOS: 4, 6, and 8, respectively.

[0029] Neutrokin-alpha receptor expression, defined functionally by the binding of biotinylated Neutrokin-alpha to cells, is found predominantly on B cells (see, *e.g.*, Moore *et al.*, *Science* 285:260-263 (1999), which is hereby incorporated by reference in its entirety), although TACI has also been reported to be expressed on activated T cells (Wang *et al.*, *Nature Immunol.* 2:577-8 (2001), which is hereby incorporated by reference in its entirety.) Furthermore, Neutrokin-alpha receptor expression is not observed in pre-B cells; rather, Neutrokin-alpha receptor expression becomes observable at the same stage in B cell development when surface Ig expression becomes

- 9 -

apparent (see, *e.g.*, Hsu *et al.*, *J. Immunol.* 168:5993-5996 (2002), which is hereby incorporated by reference in its entirety).

[0030] Receptors which bind proteins comprising Neutrokin-alpha or fragments or variants thereof may also be expressed on non-hematopoietic cells. In specific embodiments, receptors which bind proteins comprising Neutrokin-alpha or fragments or variants thereof (*e.g.*, Neutrokin-alpha heterotrimers (described below) comprising one or two APRIL monomers) are expressed on cells or cell lines of fibroblastic or epithelial lineage or having fibroblastic or epithelial morphology (*e.g.*, NIH-3T3 fibroblasts, A549 lung carcinoma cells, or HT-29 colorectal adenocarcinoma cells).

[0031] Additionally, Neutrokin-alpha receptor expression, defined functionally by the binding of biotinylated Neutrokin-alpha to cells, has been observed on multiple myeloma, Non-Hodgkin's lymphoma, and chronic lymphocytic leukemia primary tumor explants (see, *e.g.*, Briones *et al.*, *Experimental Hematology* 30:135-141 (2002) and Novak *et al.*, *Blood* 100:2973-2979 (2002), each of which is hereby incorporated by reference in its entirety) and on B lineage immortalized hematopoietic cell lines (*e.g.*, IM-9 (ATCC CCL-159); Reh (ATCC CRL-8286); ARH-77 (ATCC CRL-1621); Raji (ATCC-CCL-86); Namalwa (CRL-1432); and RPMI 8226 (ATCC CCL-155)).

[0032] Biodistribution studies of a Neutrokin-alpha protein radiolabeled with either iodine-125 (¹²⁵I) or indium-111 (¹¹¹In) injected into BALB/c mice illustrate that a Neutrokin-alpha has high in vivo targeting specificity for lymphoid tissues such as spleen and lymph nodes (See Example 1). ¹²⁵I- and ¹¹¹In-labeled Neutrokin-alpha proteins are used as indicators of the biodistribution of therapeutic Neutrokin-alpha proteins labeled with iodine-131 (¹³¹I)- or yttrium-90 (⁹⁰Y). Furthermore, administration of the ¹³¹I-labeled form of Neutrokin-alpha to mice bearing a B cell tumor has been shown to inhibit the growth of the tumor and prolong survival (Example 2). It is usually not possible or highly unfeasible to study the effects of ⁹⁰Y-labeled

- 10 -

Neutrokin- α in tumor bearing mice due to the quantity and strength of radiation involved compared to the body mass of a mouse.

[0033] Clearly, Neutrokin- α associated with a metal ion, such as a radionuclide, can be used as a therapy for B-cell malignancies. Such malignancies are responsive to radiation, and radiotherapy is an important part of the treatment plan for many patients with these diseases. Additionally, Neutrokin- α an agent comprising Neutrokin- α linked to a source of radiation can be used to treat other types of malignancies, (e.g. any type of cancer that metastasizes in the lymphatic system). Without being limited by this mechanism, cells in close proximity to cells bearing a Neutrokin- α receptor may also be killed by a Neutrokin- α protein linked to a source of radiation depending on the strength of the radioactive emission. Furthermore, an agent comprising Neutrokin- α linked to a source of radiation binds predominantly to B cells, so comparatively low doses of radiation will be effective at killing such cells.

[0034] A composition comprising Neutrokin- α associated with a radionuclide would also have application in the treatment of diseases associated with an aberrant number or function of cells expressing Neutrokin- α receptor, (e.g., lymphocytes, particularly B lymphocytes). In specific embodiments, Neutrokin- α conjugates and/or compositions of the invention have use in the treatment of autoimmune diseases, and in particular, of autoimmune diseases that involve autoantibodies.

[0035] The compositions of the invention generally comprise a Neutrokin- α protein, associated with either a chelator or a chelator and a metal ion. Herein a Neutrokin- α protein associated or linked, directly or indirectly, with a chelator is referred to as "Neutrokin- α conjugate." A "Neutrokin- α complex" herein refers to the association of a metal ion with a Neutrokin- α conjugate, preferably via coordination of the metal ion by the chelator. The following sections will describe the Neutrokin- α protein, the chelator and metal ion components of the Neutrokin- α conjugate and/or Neutrokin- α complex.

- 11 -

Neutrokin- α Conjugate

[0036] The present invention encompasses a Neutrokin- α conjugate, wherein said conjugate comprises a Neutrokin- α protein and a chelator. Such a conjugate has a number of uses, including diagnostic uses and therapeutic uses, described herein. The Neutrokin- α conjugate is also useful for preparing a Neutrokin- α complex as disclosed herein. The Neutrokin- α conjugate is a Neutrokin- α protein covalently bonded to a chelator. The following two sections describe in detail the nature of the Neutrokin- α protein, the chelator, and the association of the chelator with the Neutrokin- α protein.

[0037] The Neutrokin- α conjugate can be depicted as $NA-(Chel)_n$, wherein NA is the Neutrokin- α protein, Chel is the chelator, and n is the number of chelator molecules attached to said Neutrokin- α protein. In specific embodiments, each chelator molecule is directly attached to the Neutrokin- α protein. Therefore, if n is 3, the Neutrokin- α conjugate contains one Neutrokin- α protein with three chelator molecules, wherein each chelator molecule is bonded directly to said Neutrokin- α protein.

Neutrokin- α protein

[0038] The following section describes the Neutrokin- α protein component of the Neutrokin- α conjugates and Neutrokin- α complexes of the present invention. Nucleotides 147-1001 of SEQ ID NO:1 encode the protein shown in shown in Table 1 (SEQ ID NO:2). The nucleotide sequence of SEQ ID NO:1 was obtained by sequencing the HNEDU15 clone, which was deposited on October 22, 1996 at the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, and assigned ATCC Accession No. 97768. The deposited clone is contained in the pBluescript SK(-) plasmid (Stratagene, La Jolla, CA). The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the

International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

[0039] Neutrokin- α is a membrane protein which has an intracellular domain corresponding to amino acid 1-46 of SEQ ID NO:2, transmembrane domain (amino acids 47-72 of SEQ ID NO:2), and an extracellular domain (amino acid residues 73-285 of SEQ ID NO:2). Proteolytic cleavage of Neutrokin- α from the surface of the cell results in a mature, soluble form of the protein comprising amino acid residues 134-285 of SEQ ID NO:2.

[0040] In one embodiment, the Neutrokin- α protein is a polypeptide comprising or alternatively, consisting of, an amino acid sequence contained in SEQ ID NO:2, encoded by the cDNA contained in the ATCC Deposit No. 97768, or encoded by nucleic acids which hybridize (*e.g.*, under stringent hybridization conditions) to the nucleotide sequence contained in the ATCC Deposit No. 97768 or the complementary strand thereto. A protein fragment may be "free-standing" or within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region.

[0041] In preferred embodiments, the Neutrokin- α protein component of the Neutrokin- α conjugate or Neutrokin- α complex of the present invention corresponds to the extracellular domain of the polypeptide of SEQ ID NO:2 or a fragment or variant thereof. In other preferred embodiments, the Neutrokin- α protein component of the Neutrokin- α conjugate or Neutrokin- α complex of the present invention corresponds to the extracellular domain of the polypeptide encoded by the cDNA contained in ATCC Deposit No. 97768, or a fragment or variant thereof (*e.g.*, the mature, soluble form of the polypeptide encoded by the cDNA contained in ATCC Deposit No. 97768). Other forms of Neutrokin- α that can act as the Neutrokin- α protein component of the Neutrokin- α conjugate or Neutrokin- α complex of the present invention include the complete polypeptide encoded by the cDNA contained in ATCC Deposit No. 97768 including the intracellular, transmembrane and extracellular domains of the polypeptide encoded by the deposited cDNA, the mature, soluble polypeptide

- 13 -

encoded by the deposited cDNA, the extracellular domain minus the intracellular and transmembrane domains of the protein, the complete polypeptide of SEQ ID NO:2, the mature, soluble form of the Neutrokin- α protein (*e.g.*, amino acids 134-285 of SEQ ID NO:2 or SEQ ID NO:3), the extracellular domain of Neutrokin- α protein (amino acid residues 73-285 of SEQ ID NO:2) minus the intracellular and transmembrane domains, as well as polypeptides which have at least 80%, 85%, 90% similarity, more preferably at least 95% similarity, and still more preferably at least 96%, 97%, 98% or 99% similarity to those described above.

[0042] In another embodiment, a Neutrokin- α conjugate comprises a Neutrokin- α protein and a chelator, wherein said protein includes a polypeptide at least 80%, or at least 85% identical, more preferably at least 90% or 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to the polypeptide encoded by the deposited cDNA (ATCC Deposit No. 97768) or to the polypeptide of Tables 1 and 2 (SEQ ID NO:2 and SEQ ID NO:3, respectively), and also include portions of such polypeptides that are at least 30 amino acids and more preferably, at least 50 amino acids, in length.

[0043] By "% similarity" for two polypeptides, is intended a similarity score produced by comparing the amino acid sequences of the two polypeptides using the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711) and the default settings for determining similarity. Bestfit uses the local homology algorithm of Smith and Waterman (*Advances in Applied Mathematics* 2:482-489, 1981) to find the best segment of similarity between two sequences.

[0044] By a protein having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a Neutrokin- α protein is intended that the amino acid sequence of the protein is identical to the reference sequence except that the protein sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the Neutrokin- α protein. In other words, to obtain a protein having an

- 14 -

amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[0045] As a practical matter, whether any particular polypeptide or protein is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Tables 1 and 2, or fragments thereof, can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

[0046] In a specific embodiment, the identity between a reference (query) sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, is determined using the FASTDB computer program based on the algorithm of Brutlag *et al. Comp. App. Biosci.* 6:237-245 (1990). Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is

- 15 -

shorter. According to this embodiment, if the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction is made to the results to take into consideration the fact that the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. A determination of whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of this embodiment. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence. For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not

- 16 -

matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are made for the purposes of this embodiment.

[0047] Neutrokin alpha, like other members of the TNF family of ligands, forms biologically active trimers. Thus, in highly preferred embodiments, the Neutrokin-alpha protein component of a Neutrokin-alpha conjugate or Neutrokin-alpha complex is a trimer. Neutrokin-alpha may also form higher order structures (Liu *et al.*, *Cell* 108:383-394 (2002)). Thus, in some embodiments, a Neutrokin-alpha protein may comprise 3, 6, 9, 12, 15, 18, or 60 Neutrokin-alpha monomeric subunits. In another embodiment, the Neutrokin-alpha protein is a mature, soluble Neutrokin-alpha protein. The term "mature, soluble Neutrokin-alpha protein" refers to the portion of the extracellular domain of a Neutrokin-alpha protein that is cleaved from the membrane bound protein. In another embodiment, the Neutrokin-alpha protein is a human mature, soluble Neutrokin-alpha protein (*e.g.*, the protein of SEQ ID NO:3 or a Neutrokin-alpha protein consisting of a trimer of Neutrokin-alpha monomeric subunits wherein each subunit consists of amino acids 134-285 of the protein shown in Table 1 (SEQ ID NO:2)).

Table 1 Amino Acid Sequence of Full Length Human Neutrokin-alpha

1	MDDSTEREQS	RLTSCCLKRE	EMKLKECVSI	LPRKESPSVR	SSKDGKLLAA	TLLLALLSCC
61	LTVVSFYQVA	ALQGDLASLR	AELQGHAEK	LPAGAGAPKA	GLEEAPAVTA	GLKIFEPPAP
121	GEGNSSQNSR	NKRAVQGPEE	TVTQDCLQLI	ADSETPTIQK	GSYTFVPWLL	SFKRGSAL EE
181	KENKILVKET	GYFFIYGQVL	YTDKTYAMGH	LIQRKKVHVF	GDELSLVTLF	RCIQNMPETL
241	PNNSCYSAGI	AKLEEGDELQ	LAIPRENAQI	SLDGDVTFFG	ALKLL	

Table 2 Amino Acid Sequence of Mature Soluble Human Neutrokin-alpha

1		AVQGPEE	TVTQDCLQLI	ADSETPTIQK	GSYTFVPWLL	SFKRGSAL EE
61	KENKILVKET	GYFFIYGQVL	YTDKTYAMGH	LIQRKKVHVF	GDELSLVTLF	RCIQNMPETL
121	PNNSCYSAGI	AKLEEGDELQ	LAIPRENAQI	SLDGDVTFFG	ALKLL	

- 17 -

[0048] In the most preferred embodiments, the Neutrokin- α protein consists of a trimer of identical Neutrokin- α monomers each of which has an amino acid sequence that consists of the amino acid sequence shown in Table 2 (SEQ ID NO:3) which corresponds to amino acid residues 134-285 of the human Neutrokin- α protein shown in Table 1 (SEQ ID NO:2).

[0049] In other preferred embodiments, one or more of the individual monomeric units of a Neutrokin- α protein, preferably a Neutrokin- α trimer, may consist of a fragment or variant of the protein of SEQ ID NO:3. For example, a monomeric unit of the Neutrokin- α protein may consist of a protein that is at least 70% identical to the protein shown in SEQ ID NO:3. In other embodiments, a monomeric unit of the Neutrokin- α protein may consist of a protein that is at least 80% identical to the protein shown in SEQ ID NO:3. In preferred embodiments, a monomeric unit of the Neutrokin- α protein may consist of a protein that is at least 90% identical to the protein shown in SEQ ID NO:3. And in other preferred embodiments, a monomeric unit of the Neutrokin- α protein may consist of a protein that is at least 95% identical to the protein shown in SEQ ID NO:3. In other preferred embodiments, one or more of the individual monomeric units of a Neutrokin- α protein, preferably a Neutrokin- α trimer, may consist of Neutrokin- α protein from another species, *e.g.*, murine, canine, feline, or monkey Neutrokin- α . To be useful in the present invention, Neutrokin- α proteins must bind to one or more Neutrokin- α receptors, such as BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9). Neutrokin- α multimers comprising one or more monomers that consist of a fragment or variant of the protein of SEQ ID NO:3 may be tested for their ability to bind a Neutrokin- α receptor using any method known in the art including the method shown in Examples 11 and 12.

[0050] In other preferred embodiments, one or more of the individual monomeric units of a Neutrokin- α protein, preferably a Neutrokin- α trimer, comprise a fragment or variant of the protein of SEQ ID NO:3. For example, a monomeric unit of the Neutrokin- α protein may comprise a

- 18 -

protein that is at least 70% identical to the protein shown in SEQ ID NO:3. In other embodiments, a monomeric unit of the Neutrokin- α protein may comprise a protein that is at least 80% identical to the protein shown in SEQ ID NO:3. In preferred embodiments, a monomeric unit of the Neutrokin- α protein may comprise a protein that is at least 90% identical to the protein shown in SEQ ID NO:3. And in other preferred embodiments, a monomeric unit of the Neutrokin- α protein may comprise a protein that is at least 95% identical to the protein shown in SEQ ID NO:3. In other preferred embodiments, one or more of the individual monomeric units of a Neutrokin- α protein, preferably a Neutrokin- α trimer, may comprise a Neutrokin- α protein from another species, *e.g.*, murine, canine, feline, or monkey Neutrokin- α . To be useful in the present invention, Neutrokin- α proteins must bind to one or more Neutrokin- α receptors, such as BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9). Neutrokin- α proteins comprising one or more monomers that comprise a fragment or variant of the protein of SEQ ID NO:3 may be tested for their ability to bind a Neutrokin- α receptor using any method known in the art including the method shown in Examples 11 and 12.

[0051] In addition to forming homotrimers, Neutrokin- α proteins also form heterotrimers with APRIL, another TNF family ligand which is described, for example, in PCT International Publication Number WO97/33902 and in Hahne *et al.*, *J. Exp. Med.* 188:1185-1190 (1998), each of which is herein incorporated by reference in its entirety. The nucleotide and amino acid sequence of APRIL is given in GenBank Accession No. AF046888 (nucleotide) and AAC6132 (protein) and SEQ ID NOS: 10 and 11, respectively. Like Neutrokin- α , APRIL is processed into a mature, soluble protein which comprises 3 monomers each consisting of amino acids 105-250 of SEQ ID NO:11. Neutrokin- α and APRIL proteins can form heterotrimers wherein each heterotrimer comprises two Neutrokin- α monomers and one APRIL monomer. Alternatively, Neutrokin- α and APRIL proteins can form heterotrimers wherein each heterotrimer comprises

- 19 -

one Neutrokin- α monomer and two APRIL monomers. It is preferred that the Neutrokin- α monomer(s) in a Neutrokin- α /APRIL heterotrimer either comprise, or alternatively consist of, amino acid residues 134-285 of SEQ ID NO:2 or a fragment or variant thereof. It is preferred that the APRIL monomer(s) in a Neutrokin- α /APRIL heterotrimer either comprise, or alternatively consist of, amino acid residues 105-250 of SEQ ID NO:11 or a fragment or variant thereof. It is specifically contemplated that a Neutrokin- α /APRIL heterotrimers may be used as the Neutrokin- α protein component of the Neutrokin- α conjugates and Neutrokin- α complexes of the invention.

[0052] In other preferred embodiments, the Neutrokin- α protein comprises the amino acid residues shown in Table 1 (SEQ ID NO:2) or a fragment or variant of the protein shown in SEQ ID NO:2. Numerous functional variations of Neutrokin- α protein monomers and/or trimers can routinely be made by one of skill in the art. In the present application, a "functional" Neutrokin- α protein would be one that was capable of binding to one or more Neutrokin- α receptors. International Patent Application Publications WO98/17957, WO00/50597, and WO02/18620 each of which are hereby incorporated by reference in their entireties, describe numerous modifications that can be made to Neutrokin- α proteins. For example WO98/17957, WO00/50597, and WO02/18620 describe amino acid substitution, deletion, and addition mutations that can be made to Neutrokin- α protein as well as post translational modifications that may be made to Neutrokin- α either as a result of natural post-translational mechanisms or as a result of in vitro manipulation of the Neutrokin- α protein.

[0053] In additional embodiments, the Neutrokin- α protein component of a Neutrokin- α conjugate or Neutrokin- α complex comprises the predicted TNF-conserved domain of Neutrokin- α (amino acids 191 to 284 of SEQ ID NO:2).

[0054] To improve or alter the characteristics of Neutrokin- α proteins which are used in the Neutrokin- α conjugates or Neutrokin- α

- 20 -

complexes of the present invention, protein engineering may be employed. Recombinant DNA technology can be used to create novel mutant proteins, or "muteins," including single or multiple amino acid substitutions, deletions, additions, or fusion proteins. Such modified polypeptides can show, *e.g.*, enhanced activity or increased stability. In addition, they may be purified in higher yields and show better solubility than the corresponding natural polypeptide, at least under certain purification and storage conditions. For instance, for many proteins, including the extracellular domain or the mature form(s) of a secreted protein, it is known in the art that one or more amino acids may be deleted from the N-terminus or C-terminus without substantial loss of biological function. For instance, Ron *et al.*, *J. Biol. Chem.*, 268:2984-2988 (1993) reported modified KGF proteins that had heparin binding activity even if 3, 8, or 27 amino-terminal amino acid residues were missing.

[0055] In the present case, Neutrokine-alpha is a member of the TNF polypeptide family, with a conserved TNF family ligand domain spanning amino acid residues 191-284 of SEQ ID NO:2, deletions of N-terminal amino acids up to the Gly (G) residue at position 191 (in Table 1 (SEQ ID NO:2)) may retain some biological activity such as, for example, the ability to bind to a Neutrokine-alpha receptor or to stimulate lymphocyte (*e.g.*, B cell) proliferation, differentiation, activation, and/or survival to appropriate target cells. Polypeptides having further N-terminal deletions including the Gly (G) residue at position 191 would not be expected to retain biological activities because it is known that this residue in TNF-related polypeptides is in the beginning of the conserved domain required for biological activities. However, even if deletion of one or more amino acids from the N-terminus of a protein results in modification or loss of one or more biological functions of the protein, other functional activities may still be retained. Thus, the ability of the shortened protein to induce and/or bind to antibodies which recognize the complete protein or at least the extracellular domain of the protein generally will be retained when less than the majority of the residues of the complete or extracellular domain of the protein are removed from the N-

- 21 -

terminus. Whether a particular polypeptide lacking N-terminal residues of a complete protein retains such immunologic activities can readily be determined by routine methods described herein and/or otherwise known in the art.

[0056] Accordingly, the present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component has, for example, one or more residues deleted from the amino terminus of the amino acid sequence of the Neutrokin- α shown in Table 1 (SEQ ID NO:3), up to the glycine residue at position 191 (Gly-191 residue from the amino terminus). In particular, the present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively consists of, the amino acid sequence of residues n^1 -285 of SEQ ID NO:2, wherein n^1 is an integer in the range of the amino acid position of amino acid residues 2-190 of the amino acid sequence in SEQ ID NO:2. In specific embodiments, the Neutrokin- α conjugates and/or Neutrokin- α complexes of the present invention may comprise proteins comprising, or alternatively consisting of, an amino acid sequence selected from the group consisting of residues 2-285, 3-285, 4-285, 5-285, 6-285, 7-285, 8-285, 9-285, 10-285, 11-285, 12-285, 13-285, 14-285, 15-285, 16-285, 17-285, 18-285, 19-285, 20-285, 21-285, 22-285, 23-285, 24-285, 25-285, 26-285, 27-285, 28-285, 29-285, 30-285, 31-285, 32-285, 33-285, 34-285, 35-285, 36-285, 37-285, 38-285, 39-285, 40-285, 41-285, 42-285, 43-285, 44-285, 45-285, 46-285, 47-285, 48-285, 49-285, 50-285, 51-285, 52-285, 53-285, 54-285, 55-285, 56-285, 57-285, 58-285, 59-285, 60-285, 61-285, 62-285, 63-285, 64-285, 65-285, 66-285, 67-285, 68-285, 69-285, 70-285, 71-285, 72-285, 73-285, 74-285, 75-285, 76-285, 77-285, 78-285, 79-285, 80-285, 81-285, 82-285, 83-285, 84-285, 85-285, 86-285, 87-285, 88-285, 89-285, 90-285, 91-285, 92-285, 93-285, 94-285, 95-285, 96-285, 97-285, 98-285, 99-285, 100-285, 101-285, 102-285, 103-285, 104-285, 105-285, 106-285, 107-285, 108-285, 109-285, 110-285, 111-285, 112-285, 113-285, 114-285, 115-285, 116-

- 22 -

285, 117-285, 118-285, 119-285, 120-285, 121-285, 122-285, 123-285, 124-285, 125-285, 126-285, 127-285, 128-285, 129-285, 130-285, 131-285, 132-285, 133-285, 134-285, 135-285, 136-285, 137-285, 138-285, 139-285, 140-285, 141-285, 142-285, 143-285, 144-285, 145-285, 146-285, 147-285, 148-285, 149-285, 150-285, 151-285, 152-285, 153-285, 154-285, 155-285, 156-285, 157-285, 158-285, 159-285, 160-285, 161-285, 162-285, 163-285, 164-285, 165-285, 166-285, 167-285, 168-285, 169-285, 170-285, 171-285, 172-285, 173-285, 174-285, 175-285, 176-285, 177-285, 178-285, 179-285, 180-285, 181-285, 182-285, 183-285, 184-285, 185-285, 186-285, 187-285, 188-285, 189-285, and 190-285 of SEQ ID NO:2. The present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively, consists of, an amino acid sequence at least 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98% or 99% identical to a Neutrokin- α protein described above.

[0057] In specific embodiments, the present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively consists of, one of the following N-terminally deleted polypeptide fragments of Neutrokin- α : amino acid residues Ala-71 through Leu-285, amino acid residues Ala-81 through Leu-285, amino acid residues Leu-112 through Leu-285, amino acid residues Ala-134 through Leu-285, amino acid residues Leu-147 through Leu-285, and amino acid residues Gly-161 through Leu-285 of SEQ ID NO:2.

[0058] Similarly, many examples of biologically functional C-terminal deletion muteins are known. For instance, Interferon gamma shows up to ten times higher activities by deleting 8-10 amino acid residues from the carboxy terminus of the protein (Döbeli *et al.*, *J. Biotechnology* 7:199-216 (1988)). Since Neutrokin- α protein is a member of the TNF polypeptide family, deletions of C-terminal amino acids up to the leucine residue at position 284 are expected to retain most if not all biological activity such as, for example, ligand binding, the ability to stimulate lymphocyte (*e.g.*, B cell) proliferation,

- 23 -

differentiation, and/or activation, and modulation of cell replication. Polypeptides having deletions of up to about 10 additional C-terminal residues (*i.e.*, up to the glycine residue at position 274) also may retain some activity such as receptor binding, although such polypeptides would lack a portion of the conserved TNF domain which extends to about Leu-284 of SEQ ID NO:2. However, even if deletion of one or more amino acids from the C-terminus of a protein results in modification or loss of one or more biological functions of the protein, other functional activities may still be retained. Thus, the ability of the shortened protein to induce and/or bind to antibodies which recognize the complete or mature protein generally will be retained when less than the majority of the residues of the complete or mature protein are removed from the C-terminus. Whether a particular polypeptide lacking C-terminal residues of a complete protein retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art.

[0059] Accordingly, the present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component has one or more residues deleted from the carboxy terminus of the amino acid sequence of the Neutrokin- α protein shown in Tables 1 (SEQ ID NO:2), up to the glycine residue at position 274 (Gly-274). In particular, the present invention provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively consists of, the amino acid sequence of residues 1- m^1 of the amino acid sequence in SEQ ID NO:2, where m^1 is any integer in the range of the amino acid position of amino acid residues 274-284 in SEQ ID NO:2. More in particular, the invention provides a conjugate as described above wherein said Neutrokin- α protein comprises, or alternatively consists of, an amino acid sequence selected from the group consisting of residues 1-274, 1-275, 1-276, 1-277, 1-278, 1-279, 1-280, 1-281, 1-282, 1-283 and 1-284 of SEQ ID NO:2. The present invention is also directed to Neutrokin- α conjugates and/or Neutrokin- α complexes

- 24 -

wherein the Neutrokin- α protein component comprises, or alternatively, consists of, an amino acid sequence at least 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98% or 99% identical to the polypeptide sequence encoding the Neutrokin- α proteins described above.

[0060] The present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component has one or more residues deleted from both the amino and the carboxyl termini, which may be described generally as having residues n^1 - m^1 of SEQ ID NO:2, where n^1 and m^1 are integers as defined above.

[0061] In additional embodiments, the invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively consists of amino acid sequence of residues 134- m^2 of SEQ ID NO:2, where m^2 is an integer from 140 to 285, corresponding to the position of the amino acid residue in SEQ ID NO:2. For example, the invention provides further provides a conjugate as described above wherein said Neutrokin- α protein comprises, or alternatively consists of, an amino acid sequence selected from the group consisting of residues A-134 to Leu-285; A-134 to L-284; A-134 to K-283; A-134 to L-282; A-134 to A-281; A-134 to G-280; A-134 to F-279; A-134 to F-278; A-134 to T-277; A-134 to V-276; A-134 to D-275; A-134 to G-274; A-134 to D-273; A-134 to L-272; A-134 to S-271; A-134 to I-270; A-134 to Q-269; A-134 to A-268; A-134 to N-267; A-134 to E-266; A-134 to R-265; A-134 to P-264; A-134 to I-263; A-134 to A-262; A-134 to L-261; A-134 to Q-260; A-134 to L-259; A-134 to E-258; A-134 to D-257; A-134 to G-256; A-134 to E-255; A-134 to E-254; A-134 to L-253; A-134 to K-252; A-134 to A-251; A-134 to I-250; A-134 to G-249; A-134 to A-248; A-134 to S-247; A-134 to Y-246; A-134 to C-245; A-134 to S-244; A-134 to N-243; A-134 to N-242; A-134 to P-241; A-134 to L-240; A-134 to T-239; A-134 to E-238; A-134 to P-237; A-134 to M-236; A-134 to N-235; A-134 to Q-234; A-134 to I-233; A-134 to C-232; A-134 to R-231; A-134 to F-230; A-134 to L-229; A-134 to T-228; A-134 to V-227; A-134 to L-226; A-134 to S-225; A-134 to L-

- 25 -

224; A-134 to E-223; A-134 to D-222; A-134 to G-221; A-134 to F-220; A-134 to V-219; A-134 to H-218; A-134 to V-217; A-134 to K-216; A-134 to K-215; A-134 to R-214; A-134 to Q-213; A-134 to I-212; A-134 to L-211; A-134 to H-210; A-134 to G-209; A-134 to M-208; A-134 to A-207; A-134 to Y-206; A-134 to T-205; A-134 to K-204; A-134 to D-203; A-134 to T-202; A-134 to Y-201; A-134 to L-200; A-134 to V-199; A-134 to Q-198; A-134 to G-197; A-134 to Y-196; A-134 to I-195; A-134 to F-194; A-134 to F-193; A-134 to Y-192; A-134 to G-191; A-134 to T-190; A-134 to E-189; A-134 to K-188; A-134 to V-187; A-134 to L-186; A-134 to I-185; A-134 to K-184; A-134 to N-183; A-134 to E-182; A-134 to K-181; A-134 to E-180; A-134 to E-179; A-134 to L-178; A-134 to A-177; A-134 to S-176; A-134 to G-175; A-134 to R-174; A-134 to K-173; A-134 to F-172; A-134 to S-171; A-134 to L-170; A-134 to L-169; A-134 to W-168; A-134 to P-167; A-134 to V-166; A-134 to F-165; A-134 to T-164; A-134 to Y-163; A-134 to S-162; A-134 to G-161; A-134 to K-160; A-134 to Q-159; A-134 to I-158; A-134 to T-157; A-134 to P-156; A-134 to T-155; A-134 to E-154; A-134 to S-153; A-134 to D-152; A-134 to A-151; A-134 to I-150; A-134 to L-149; A-134 to Q-148; A-134 to L-147; A-134 to C-146; A-134 to D-145; A-134 to Q-144; A-134 to T-143; A-134 to V-142; A-134 to T-141; and A-134 to E-140 of SEQ ID NO:2. The present invention further provides Neutrokin- α conjugates and/or Neutrokin- α complexes wherein the Neutrokin- α protein component comprises, or alternatively, consists of, an amino acid sequence at least 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98% or 99% identical to a Neutrokin- α protein described above.

[0062] A variant is a protein that may include one or more amino acid substitutions, deletions, or additions, when compared to a reference protein, *e.g.*, SEQ ID NO:3 or SEQ ID NO:2. The substitutions, deletions, or additions may be the result of natural mutations, or human manipulation. Thus, a Neutrokin- α protein variant may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation. As indicated, any change is preferably of a minor nature, such

- 26 -

as conservative amino acid substitutions (see Table 3) that do not significantly affect the folding or activity of the protein, in this case the ability of the Neutrokin- α protein to bind a Neutrokin- α receptor.

TABLE 3. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

[0063] For example, site directed changes at the amino acid level of a Neutrokin- α protein can be made by replacing a particular amino acid with a conservative substitution. Preferred conservative substitution mutations of the Neutrokin- α amino acid sequence provided in Table 3 include: M1 replaced with A, G, I, L, S, T, or V; D2 replaced with E; D3 replaced with E; S4 replaced with A, G, I, L, T, M, or V; T5 replaced with A, G, I, L, S, M, or V; E6 replaced with D; R7 replaced with H, or K; E8 replaced with D; Q9 replaced with N; S10 replaced with A, G, I, L, T, M, or V; R11 replaced with H, or K; L12 replaced with A, G, I, S, T, M, or V; T13 replaced with A, G, I, L, S, M, or V; S14 replaced with A, G, I, L, T, M, or V; L16 replaced with A, G, I, S, T, M, or V; K17 replaced with H, or R; K18 replaced with H, or R;

- 27 -

R19 replaced with H, or K; E20 replaced with D; E21 replaced with D; M22 replaced with A, G, I, L, S, T, or V; K23 replaced with H, or R; L24 replaced with A, G, I, S, T, M, or V; K25 replaced with H, or R; E26 replaced with D; V28 replaced with A, G, I, L, S, T, or M; S29 replaced with A, G, I, L, T, M, or V; I30 replaced with A, G, L, S, T, M, or V; L31 replaced with A, G, I, S, T, M, or V; R33 replaced with H, or K; K34 replaced with H, or R; E35 replaced with D; S36 replaced with A, G, I, L, T, M, or V; S38 replaced with A, G, I, L, T, M, or V; V39 replaced with A, G, I, L, S, T, or M; R40 replaced with H, or K; S41 replaced with A, G, I, L, T, M, or V; S42 replaced with A, G, I, L, T, M, or V; K43 replaced with H, or R; D44 replaced with E; G45 replaced with A, I, L, S, T, M, or V; K46 replaced with H, or R; L47 replaced with A, G, I, S, T, M, or V; L48 replaced with A, G, I, S, T, M, or V; A49 replaced with G, I, L, S, T, M, or V; A50 replaced with G, I, L, S, T, M, or V; T51 replaced with A, G, I, L, S, M, or V; L52 replaced with A, G, I, S, T, M, or V; L53 replaced with A, G, I, S, T, M, or V; L54 replaced with A, G, I, S, T, M, or V; A55 replaced with G, I, L, S, T, M, or V; L56 replaced with A, G, I, S, T, M, or V; L57 replaced with A, G, I, S, T, M, or V; S58 replaced with A, G, I, L, T, M, or V; L61 replaced with A, G, I, S, T, M, or V; T62 replaced with A, G, I, L, S, M, or V; V63 replaced with A, G, I, L, S, T, or M; V64 replaced with A, G, I, L, S, T, or M; S65 replaced with A, G, I, L, T, M, or V; F66 replaced with W, or Y; Y67 replaced with F, or W; Q68 replaced with N; V69 replaced with A, G, I, L, S, T, or M; A70 replaced with G, I, L, S, T, M, or V; A71 replaced with G, I, L, S, T, M, or V; L72 replaced with A, G, I, S, T, M, or V; Q73 replaced with N; G74 replaced with A, I, L, S, T, M, or V; D75 replaced with E; L76 replaced with A, G, I, S, T, M, or V; A77 replaced with G, I, L, S, T, M, or V; S78 replaced with A, G, I, L, T, M, or V; L79 replaced with A, G, I, S, T, M, or V; R80 replaced with H, or K; A81 replaced with G, I, L, S, T, M, or V; E82 replaced with D; L83 replaced with A, G, I, S, T, M, or V; Q84 replaced with N; G85 replaced with A, I, L, S, T, M, or V; H86 replaced with K, or R; H87 replaced with K, or R; A88 replaced with G, I, L, S, T, M, or V; E89 replaced with D; K90 replaced with H, or R; L91

- 28 -

replaced with A, G, I, S, T, M, or V; A93 replaced with G, I, L, S, T, M, or V; G94 replaced with A, I, L, S, T, M, or V; A95 replaced with G, I, L, S, T, M, or V; G96 replaced with A, I, L, S, T, M, or V; A97 replaced with G, I, L, S, T, M, or V; K99 replaced with H, or R; A100 replaced with G, I, L, S, T, M, or V; G101 replaced with A, I, L, S, T, M, or V; L102 replaced with A, G, I, S, T, M, or V; E103 replaced with D; E104 replaced with D; A105 replaced with G, I, L, S, T, M, or V; A107 replaced with G, I, L, S, T, M, or V; V108 replaced with A, G, I, L, S, T, or M; T109 replaced with A, G, I, L, S, M, or V; A110 replaced with G, I, L, S, T, M, or V; G111 replaced with A, I, L, S, T, M, or V; L112 replaced with A, G, I, S, T, M, or V; K113 replaced with H, or R; I114 replaced with A, G, L, S, T, M, or V; F115 replaced with W, or Y; E116 replaced with D; A119 replaced with G, I, L, S, T, M, or V; G121 replaced with A, I, L, S, T, M, or V; E122 replaced with D; G123 replaced with A, I, L, S, T, M, or V; N124 replaced with Q; S125 replaced with A, G, I, L, T, M, or V; S126 replaced with A, G, I, L, T, M, or V; Q127 replaced with N; N128 replaced with Q; S129 replaced with A, G, I, L, T, M, or V; R130 replaced with H, or K; N131 replaced with Q; K132 replaced with H, or R; R133 replaced with H, or K; A134 replaced with G, I, L, S, T, M, or V; V135 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q136 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G137 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P138 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; E139 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E140 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T141 replaced with A, G, I, L, S, M, or V; V142 replaced with A, G, I, L, S, T, or M; T143 replaced with A, G, I, L, S, M, or V; Q144 replaced with N; D145 replaced with E; L147 replaced with A, G, I, S, T, M, or V; Q148 replaced with N; L149 replaced with A, G, I, S, T, M, or V; I150 replaced with A, G, L, S, T, M, or V; A151 replaced with G, I, L, S, T, M, or V; D152 replaced with E; S153 replaced with A, G, I, L, T, M, or V; E154 replaced with D; T155 replaced with A, G, I, L, S, M, or V; T157 replaced with A, G, I, L, S, M, or V; I158 replaced with A, G, L, S, T, M, or

- 29 -

V; Q159 replaced with N; K160 replaced with H, or R; G161 replaced with A, I, L, S, T, M, or V; S162 replaced with A, G, I, L, T, M, or V; Y163 replaced with F, or W; T164 replaced with A, G, I, L, S, M, or V; F165 replaced with W, or Y; V166 replaced with A, G, I, L, S, T, or M; W168 replaced with F, or Y; L169 replaced with A, G, I, S, T, M, or V; L170 replaced with A, G, I, S, T, M, or V; S171 replaced with A, G, I, L, T, M, or V; F172 replaced with W, or Y; K173 replaced with H, or R; R174 replaced with H, or K; G175 replaced with A, I, L, S, T, M, or V; S176 replaced with A, G, I, L, T, M, or V; A177 replaced with G, I, L, S, T, M, or V; L178 replaced with A, G, I, S, T, M, or V; E179 replaced with D; E180 replaced with D; K181 replaced with H, or R; E182 replaced with D; N183 replaced with Q; K184 replaced with H, or R; I185 replaced with A, G, L, S, T, M, or V; L186 replaced with A, G, I, S, T, M, or V; V187 replaced with A, G, I, L, S, T, or M; K188 replaced with H, or R; E189 replaced with D; T190 replaced with A, G, I, L, S, M, or V; G191 replaced with A, I, L, S, T, M, or V; Y192 replaced with F, or W; F193 replaced with W, or Y; F194 replaced with W, or Y; I195 replaced with A, G, L, S, T, M, or V; Y196 replaced with F, or W; G197 replaced with A, I, L, S, T, M, or V; Q198 replaced with N; V199 replaced with A, G, I, L, S, T, or M; L200 replaced with A, G, I, S, T, M, or V; Y201 replaced with F, or W; T202 replaced with A, G, I, L, S, M, or V; D203 replaced with E; K204 replaced with H, or R; T205 replaced with A, G, I, L, S, M, or V; Y206 replaced with F, or W; A207 replaced with G, I, L, S, T, M, or V; M208 replaced with A, G, I, L, S, T, or V; G209 replaced with A, I, L, S, T, M, or V; H210 replaced with K, or R; L211 replaced with A, G, I, S, T, M, or V; I212 replaced with A, G, L, S, T, M, or V; Q213 replaced with N; R214 replaced with H, or K; K215 replaced with H, or R; K216 replaced with H, or R; V217 replaced with A, G, I, L, S, T, or M; H218 replaced with K, or R; V219 replaced with A, G, I, L, S, T, or M; F220 replaced with W, or Y; G221 replaced with A, I, L, S, T, M, or V; D222 replaced with E; E223 replaced with D; L224 replaced with A, G, I, S, T, M, or V; S225 replaced with A, G, I, L, T, M, or V; L226 replaced with A, G, I, S, T, M, or V; V227 replaced with A, G, I, L, S, T, or M; T228

- 30 -

replaced with A, G, I, L, S, M, or V; L229 replaced with A, G, I, S, T, M, or V; F230 replaced with W, or Y; R231 replaced with H, or K; I233 replaced with A, G, L, S, T, M, or V; Q234 replaced with N; N235 replaced with Q; M236 replaced with A, G, I, L, S, T, or V; E238 replaced with D; T239 replaced with A, G, I, L, S, M, or V; L240 replaced with A, G, I, S, T, M, or V; N242 replaced with Q; N243 replaced with Q; S244 replaced with A, G, I, L, T, M, or V; Y246 replaced with F, or W; S247 replaced with A, G, I, L, T, M, or V; A248 replaced with G, I, L, S, T, M, or V; G249 replaced with A, I, L, S, T, M, or V; I250 replaced with A, G, L, S, T, M, or V; A251 replaced with G, I, L, S, T, M, or V; K252 replaced with H, or R; L253 replaced with A, G, I, S, T, M, or V; E254 replaced with D; E255 replaced with D; G256 replaced with A, I, L, S, T, M, or V; D257 replaced with E; E258 replaced with D; L259 replaced with A, G, I, S, T, M, or V; Q260 replaced with N; L261 replaced with A, G, I, S, T, M, or V; A262 replaced with G, I, L, S, T, M, or V; I263 replaced with A, G, L, S, T, M, or V; R265 replaced with H, or K; E266 replaced with D; N267 replaced with Q; A268 replaced with G, I, L, S, T, M, or V; Q269 replaced with N; I270 replaced with A, G, L, S, T, M, or V; S271 replaced with A, G, I, L, T, M, or V; L272 replaced with A, G, I, S, T, M, or V; D273 replaced with E; G274 replaced with A, I, L, S, T, M, or V; D275 replaced with E; V276 replaced with A, G, I, L, S, T, or M; T277 replaced with A, G, I, L, S, M, or V; F278 replaced with W, or Y; F279 replaced with W, or Y; G280 replaced with A, I, L, S, T, M, or V; A281 replaced with G, I, L, S, T, M, or V; L282 replaced with A, G, I, S, T, M, or V; K283 replaced with H, or R; L284 replaced with A, G, I, S, T, M, or V; and/or L285 replaced with A, G, I, S, T, M, or V. The resulting Neutrokin- α proteins may be routinely screened for Neutrokin- α functional activity and/or physical properties (such as, for example, the ability to bind one or more Neutrokin- α receptors and/or enhanced or reduced stability and/or solubility). The resulting Neutrokin- α protein variant may be used in a conjugate as described above.

- 31 -

- [0064] Of special interest are substitutions of charged amino acids with other charged or neutral amino acids which may produce proteins with highly desirable improved characteristics, such as less aggregation. Aggregation may not only reduce activity but also be problematic when preparing pharmaceutical formulations, because aggregates can be immunogenic (Pinckard *et al.*, *Clin. Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36: 838-845 (1987); Cleland *et al.*, *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).
- [0065] A partial, non-limiting, and non-exclusive list of residues of the Neutrokin- α protein sequence which may be targeted for mutation includes the following amino acid residues of the Neutrokin- α protein sequence as shown in Table 1 (SEQ ID NO:2): V-142; T-143; Q-144; D-145; C-146; L-147; Q-148; L-149; I-150; A-151; D-152; S-153; E-154; T-155; P-156; T-157; I-158; Q-159; and K-160.
- [0066] In another embodiment, a Neutrokin- α conjugate or Neutrokin- α complex of the invention comprises a Neutrokin- α fragment or variant, wherein said fragment or variant is a Neutrokin- α protein having an amino acid sequence containing one or more non-conservative substitutions of the amino acid sequence provided in SEQ ID NO:2. For example, non-conservative substitutions of the Neutrokin- α protein sequence provided in SEQ ID NO:2 include: M1 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D2 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; D3 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S4 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T5 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E6 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; R7 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E8 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; Q9 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; S10 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R11 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L12 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T13 replaced with D, E, H, K, R, N, Q, F,

- 32 -

W, Y, P, or C; S14 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C15 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; L16 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K17 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K18 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; R19 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E20 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E21 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; M22 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K23 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L24 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K25 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E26 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; C27 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; V28 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S29 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I30 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L31 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P32 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; R33 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K34 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E35 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S36 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P37 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; S38 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V39 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R40 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S41 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S42 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K43 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; D44 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G45 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K46 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L47 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L48 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A49 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A50

- 33 -

replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T51 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L52 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L53 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L54 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A55 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L56 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L57 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S58 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C59 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; C60 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; L61 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T62 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V63 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V64 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S65 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F66 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; Y67 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; Q68 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; V69 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A70 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A71 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L72 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q73 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G74 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D75 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L76 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A77 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S78 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L79 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R80 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A81 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E82 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L83 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q84 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G85 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; H86 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; H87 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A88

- 34 -

replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E89 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K90 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L91 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P92 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; A93 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G94 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A95 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G96 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A97 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P98 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; K99 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A100 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G101 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L102 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E103 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E104 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A105 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P106 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; A107 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V108 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T109 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A110 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G111 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L112 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K113 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; I114 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F115 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; E116 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; P117 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; P118 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; A119 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P120 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; G121 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E122 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G123 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; N124 replaced with D, E, H, K, R, A, G, I, L, S,

- 35 -

T, M, V, F, W, Y, P, or C; S125 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S126 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q127 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; N128 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; S129 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; R130 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; N131 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; K132 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; R133 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; A134 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V135 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q136 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; G137 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P138 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; E139 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E140 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T141 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V142 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T143 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q144 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; D145 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; C146 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; L147 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q148 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; L149 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I150 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A151 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D152 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; S153 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E154 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T155 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P156 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; T157 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I158 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q159 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; K160

- 36 -

replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G161 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S162 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y163 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; T164 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F165 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; V166 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P167 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; W168 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; L169 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L170 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S171 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F172 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; K173 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; R174 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G175 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S176 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A177 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L178 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E179 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E180 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K181 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E182 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; N183 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; K184 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; I185 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L186 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V187 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K188 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E189 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T190 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G191 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y192 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; F193 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; F194 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; I195 replaced with D, E, H, K, R, N, Q, F, W, Y, P,

- 37 -

or C; Y196 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; G197 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q198 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; V199 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L200 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y201 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; T202 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D203 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K204 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T205 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Y206 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; A207 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; M208 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G209 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; H210 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L211 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I212 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q213 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; R214 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K215 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; K216 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V217 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; H218 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V219 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F220 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; G221 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D222 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E223 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L224 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S225 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L226 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; V227 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T228 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L229 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F230 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; R231 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; C232 replaced with D, E, H, K,

- 38 -

R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; I233 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q234 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; N235 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; M236 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P237 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; E238 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; T239 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L240 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P241 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; N242 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; N243 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; S244 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; C245 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or P; Y246 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; S247 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A248 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; G249 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I250 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A251 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K252 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L253 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; E254 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E255 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G256 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D257 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E258 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L259 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q260 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; L261 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A262 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; I263 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; P264 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, or C; R265 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; E266 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; N267 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C;

- 39 -

A268 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; Q269 replaced with D, E, H, K, R, A, G, I, L, S, T, M, V, F, W, Y, P, or C; I270 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; S271 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L272 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D273 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; G274 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; D275 replaced with H, K, R, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; V276 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; T277 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; F278 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; F279 replaced with D, E, H, K, R, N, Q, A, G, I, L, S, T, M, V, P, or C; G280 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; A281 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; L282 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; K283 replaced with D, E, A, G, I, L, S, T, M, V, N, Q, F, W, Y, P, or C; L284 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C; and/or L285 replaced with D, E, H, K, R, N, Q, F, W, Y, P, or C. The resulting Neutrokin- α protein, Neutrokin- α conjugate, or Neutrokin- α complex of the invention may be routinely screened for Neutrokin- α functional activities and/or physical properties (such as, for example, the ability to bind one or more Neutrokin- α receptors and/or enhanced or reduced stability and/or solubility) described throughout the specification and known in the art. In one embodiment, the resulting Neutrokin- α protein, Neutrokin- α conjugate, or Neutrokin- α complex of the invention has either and increased or decreased Neutrokin- α functional activity while maintaining the ability to bind to a Neutrokin- α receptor and be internalized into a cell.

[0067] The resulting Neutrokin- α conjugate may be routinely screened for Neutrokin- α functional activities and/or physical properties (such as, for example, enhanced or reduced stability and/or solubility) described throughout the specification and known in the art.

[0068] A partial, non-limiting and non-exclusive list of residues of the Neutrokin- α protein sequence which may be targeted for mutation

- 40 -

includes one or more of the following amino acid residues of the Neutrokin- α protein sequence as shown in Table 1 (SEQ ID NO:2): V-142; T-143; Q-144; D-145; C-146; L-147; Q-148; L-149; I-150; A-151; D-152; S-153; E-154; T-155; P-156; T-157; I-158; Q-159; and K-160.

[0069] Thus, the invention also encompasses Neutrokin- α conjugates and Neutrokin- α complexes wherein the Neutrokin- α protein component has one or more amino acid residues deleted, added, or substituted to generate Neutrokin- α protein that is better suited for expression, scale up, etc., in the host cells chosen. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges; N-linked glycosylation sites can be altered or eliminated to achieve, for example, expression of a homogeneous product that is more easily recovered and purified from yeast hosts which are known to hyperglycosylate N-linked sites. To this end, a variety of amino acid substitutions at one or both of the first or third amino acid positions on any one or more of the glycosylation recognition sequences in the Neutrokin- α protein, and/or an amino acid deletion at the second position of any one or more such recognition sequences will prevent glycosylation of the Neutrokin- α at the modified tripeptide sequence (see, *e.g.*, Miyajimo *et al.*, *EMBO J* 5:1193-1197). By way of a non-limiting example, mutation of the serine at position 244 to alanine either singly or in combination with mutation of the asparagine at position 242 to glutamine abolishes glycosylation of the mature, soluble form of Neutrokin- α (amino acids 134-285) of SEQ ID NO:2) when expressed in the yeast *Pichia pastoris*. A mutant Neutrokin- α protein in which only the asparagine at position 242 is mutated to glutamine is still glycosylated when expressed in *Pichia pastoris*. In this mutant, the glycosylation event may be due to the activation or unmasking of an O-linked glycosylation site at serine 244.

[0070] Additionally, one or more of the amino acid residues of a Neutrokin- α protein (*e.g.*, arginine and lysine residues) may be deleted or substituted with another residue to eliminate undesired processing by proteases such as,

- 41 -

for example, furins or kexins. One possible result of such a mutation is that Neutrokin- α protein is not cleaved and released from the cell surface.

[0071] In a specific embodiment, Lys-132 and/or Arg-133 of the Neutrokin- α sequence shown in SEQ ID NO:2 is mutated to another amino acid residue, or deleted altogether, to prevent or diminish release of the soluble form of Neutrokin- α from cells expressing Neutrokin- α . In a more specific embodiment, Lys-132 of the Neutrokin- α sequence shown in SEQ ID NO:2 is mutated to Ala-132. In another, nonexclusive specific embodiment, Arg-133 of the Neutrokin- α sequence shown in SEQ ID NO:2 is mutated to Ala-133. These mutated proteins have uses such as, for example, in *ex vivo* therapy or gene therapy, to engineer cells expressing a Neutrokin- α polypeptide that is retained on the surface of the engineered cells.

[0072] In a specific embodiment, Cys-146 of the Neutrokin- α sequence shown in SEQ ID NO:1 or SEQ ID NO:2 is mutated to another amino acid residue, or deleted altogether, for example, to aid preventing or diminishing oligomerization of the mutant Neutrokin- α protein when expressed in an expression system. In a specific embodiment, Cys-146 is replaced with a serine amino acid residue.

[0073] In another specific embodiment, Cys-232 of the Neutrokin- α sequence shown in SEQ ID NO:1 or SEQ ID NO:2 is mutated to another amino acid residue, or deleted altogether, for example, to aid preventing or diminishing oligomerization of the mutant Neutrokin- α protein when expressed in an expression system (essentially as described in Example 1). In a specific embodiment, Cys-232 is replaced with a serine amino acid residue.

[0074] In yet another specific embodiment, Cys-245 of the Neutrokin- α sequence shown in SEQ ID NO:1 or SEQ ID NO:2 is mutated to another amino acid residue, or deleted altogether, for example, to aid preventing or diminishing oligomerization of the mutant Neutrokin- α protein when expressed in an expression system (essentially as described in Example 1). In a specific embodiment, Cys-245 is replaced with a serine amino acid residue.

- 42 -

[0075] As is known in the art, many peptides and proteins may contain carbohydrates attached to the peptide or protein. The Neutrokin- α protein used in the present invention may optionally contain one or more carbohydrates attached.

Vectors and Host Cells For Producing Neutrokin- α

[0076] The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, or which are otherwise engineered to produce the polypeptides of the invention, and the production of Neutrokin- α and/or Neutrokin- α SV polypeptides, or fragments thereof, by recombinant or synthetic techniques.

[0077] In one embodiment, the polynucleotides of the invention are joined to a vector (e.g., a cloning or expression vector). The vector may be, for example, a phage, plasmid, viral or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells. The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Introduction of the vector construct into the host cell can be effected by techniques known in the art which include, but are not limited to, calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986).

[0078] Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The

- 43 -

heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, for example, stabilization or simplified purification of expressed recombinant product.

[0079] In one embodiment, the DNA of the invention is operatively associated with an appropriate heterologous regulatory element (e.g., promoter or enhancer), such as, the phage lambda PL promoter, the *E. coli lac*, *trp*, *phoA*, and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan.

[0080] As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293 and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

[0081] The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. The host strain may be chosen which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide

- 44 -

may be controlled. Furthermore, different host cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the foreign protein expressed. Selection of appropriate vectors and promoters for expression in a host cell is a well-known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the host are routine skills in the art.

[0082] Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice. As a representative, but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well-known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Among vectors preferred for use in bacteria include pHE4-5 (ATCC Accession No. 209311; and variations thereof), pQE70, pQE60 and pQE-9, available from QIAGEN, Inc., *supra*; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not

- 45 -

limited to, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, pPIC9K, and PAO815 (all available from Invitrogen, Carlsbad, CA). Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL (available from Pharmacia). Other suitable vectors will be readily apparent to the skilled artisan.

[0083] Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

[0084] Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

[0085] In one embodiment, the yeast *Pichia pastoris* is used to express Neutrokin- α protein in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A main step in the methanol metabolism pathway is the oxidation of methanol to formaldehyde using O_2 . This reaction is catalyzed by the enzyme alcohol oxidase. In order to metabolize methanol as its sole carbon source, *Pichia pastoris* must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O_2 . Consequently, in a growth medium depending on methanol as a main carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1* gene comprises up to approximately 30% of the total soluble protein in *Pichia pastoris*. See, Ellis, S.B., *et al.*, *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J,

- 46 -

et al., *Yeast* 5:167-77 (1989); Tschopp, J.F., *et al.*, *Nucl. Acids Res.* 15:3859-76 (1987). Thus, a heterologous coding sequence, such as, for example, a Neutrokin- α or Neutrokin- α SV polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

[0086] In one example, the plasmid vector pPIC9K is used to express DNA encoding a Neutrokin- α or Neutrokin- α SV polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press, Totowa, NJ, 1998. This expression vector allows expression and secretion of a Neutrokin- α or Neutrokin- α SV protein of the invention by virtue of the strong *AOX1* promoter linked to the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal peptide (i.e., leader) located upstream of a multiple cloning site.

[0087] Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZ α , pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815, as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

[0088] In one embodiment, high-level expression of a heterologous coding sequence, such as, for example, a Neutrokin- α or Neutrokin- α SV polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an expression vector such as, for example, pGAPZ or pGAPZ α , and growing the yeast culture in the absence of methanol.

[0089] Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are *cis*-acting elements of DNA, usually about

- 47 -

from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

[0090] Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman (*Cell* 23:175 (1981)), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

[0091] In a specific embodiment, constructs designed to express a portion of the extracellular domain of the Neutrokin- α (e.g., amino acid residues Ala-134 through Leu-285) are preferred. One of skill in the art would be able to use the polynucleotide and polypeptide sequences provided as SEQ ID NO:1 and SEQ ID NO:2, respectively, or SEQ ID NO:18 and SEQ ID NO:19, respectively, to design polynucleotide primers to generate such an expression construct.

[0092] In another embodiment, constructs designed to express the entire predicted extracellular domain of the Neutrokin- α (i.e., amino acid residues Gln-73 through Leu-285) are preferred. One of skill in the art would be able to use the polynucleotide and polypeptide sequences provided as SEQ ID NO:1 and SEQ ID NO:2, respectively, or SEQ ID NO:18 and SEQ ID NO:19, respectively, to design polynucleotide primers to generate such an expression construct.

[0093] In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and

- 48 -

immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., Neutrokin- α coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with Neutrokin- α polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous Neutrokin- α polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous Neutrokin- α polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

[0094] The host cells described *infra* can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, cell-free translation systems can also be employed to produce the polypeptides of the invention using RNAs derived from the DNA constructs of the present invention.

[0095] The polypeptide of the invention may be expressed or synthesized in a modified form, such as a fusion protein (comprising the polypeptide joined via a peptide bond to a heterologous protein sequence (of a different protein)), and may include not only secretion signals, but also additional heterologous functional regions. Such a fusion protein can be made by ligating polynucleotides of the invention and the desired nucleic acid sequence encoding the desired amino acid sequence to each other, by methods known in the art, in the proper reading frame, and expressing the fusion protein product by methods known in the art. Alternatively, such a fusion protein can be made by protein synthetic techniques, e.g., by use of a peptide synthesizer. Thus, for instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability

- 49 -

and persistence in the host cell, during purification, or during subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art.

[0096] In one embodiment, polynucleotides encoding Neutrokin- α and/or Neutrokin- α SV polypeptides of the invention may be fused to signal sequences which will direct the localization of a protein of the invention to particular compartments of a prokaryotic or eukaryotic cell and/or direct the secretion of a protein of the invention from a prokaryotic or eukaryotic cell. For example, in *E. coli*, one may wish to direct the expression of the protein to the periplasmic space. Examples of signal sequences or proteins (or fragments thereof) to which the polypeptides of the invention may be fused in order to direct the expression of the polypeptide to the periplasmic space of bacteria include, but are not limited to, the *pelB* signal sequence, the maltose binding protein (MBP) signal sequence, MBP, the *ompA* signal sequence, the signal sequence of the periplasmic *E. coli* heat-labile enterotoxin B-subunit, and the signal sequence of alkaline phosphatase. Several vectors are commercially available for the construction of fusion proteins which will direct the localization of a protein, such as the pMAL series of vectors (particularly the pMAL-p series) available from New England Biolabs. In a specific embodiment, polynucleotides encoding Neutrokin- α and/or Neutrokin- α SV polypeptides of the invention may be fused to the *pelB* pectate lyase signal sequence to increase the efficiency of expression and purification of such polypeptides in Gram-negative bacteria. *See*, U.S. Patent Nos. 5,576,195 and 5,846,818, the contents of which are herein incorporated by reference in their entireties.

[0097] Examples of signal peptides that may be fused to a polypeptide of the invention in order to direct its secretion in mammalian cells include, but are

- 50 -

not limited to, the MPIF-1 signal sequence (amino acids 1-21 of GenBank Accession number AAB51134), the stanniocalcin signal sequence (MLQNSAVLLLLVISASA, SEQ ID NO:45), and a consensus signal sequence (MPTWAWWLFLVLLLALWAPARG, SEQ ID NO:46). A suitable signal sequence that may be used in conjunction with baculoviral expression systems is the gp67 signal sequence, (amino acids 1-19 of GenBank Accession Number AAA72759).

[0098] A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to stabilize and purify proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5 has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett et al., *J. Molecular Recognition* 8:52-58 (1995) and K. Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995).

[0099] Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include

- 51 -

an initial modified methionine residue, in some cases as a result of host-mediated processes.

[0100] Neutrokin- α protein includes naturally purified Neutrokin- α , Neutrokin- α of chemical synthetic procedures, and Neutrokin- α produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells (see, *e.g.*, Example 3). Depending upon the host employed in a recombinant production procedure, the Neutrokin- α protein may be glycosylated or may be non-glycosylated. In addition, Neutrokin- α may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

[0101] As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention (*e.g.*, the soluble mature form of human Neutrokin- α) can be fused to heterologous polypeptide sequences. For example, Neutrokin- α (including fragments or variants thereof), may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof, resulting in chimeric polypeptides. By way of another non-limiting example, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) may be fused with albumin (including but not limited to recombinant human serum albumin or fragments or variants thereof (see, *e.g.*, U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)). In a preferred embodiment, Neutrokin- α polypeptides (including fragments or variants thereof) are fused with the mature form of human serum albumin (*i.e.*, amino acids 1 – 585 of human serum albumin as shown in Figures 1 and 2 of EP Patent 0 322 094) which is herein incorporated by reference in its entirety. In another preferred embodiment, Neutrokin- α polypeptides (including fragments or variants thereof) are fused with polypeptide fragments comprising, or alternatively consisting of, amino acid residues 1-x of human serum albumin, where x is an

- 52 -

integer from 1 to 585 and the albumin fragment has human serum albumin activity. In another preferred embodiment, Neutrokin- α polypeptides (including fragments or variants thereof) are fused with polypeptide fragments comprising, or alternatively consisting of, amino acid residues 1-z of human serum albumin, where z is an integer from 369 to 419, as described in U.S. Patent 5,766,883 herein incorporated by reference in its entirety. Neutrokin- α polypeptides p (including fragments or variants thereof) may be fused to either the N- or C-terminal end of the heterologous protein (e.g., immunoglobulin Fc polypeptide or human serum albumin polypeptide).

[0102] Such fusion proteins as those described above may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves

- 53 -

as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

[0103] Neutrokin- α can be chemically synthesized using techniques known in the art (*e.g.*, see Creighton, 1983, *Proteins: Structures and Molecular Principles*, W.H. Freeman & Co., N.Y.; and Hunkapiller, M., *et al.*, *Nature* 310:105-111 (1984)). For example, a peptide corresponding to a fragment of the Neutrokin- α can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the Neutrokin- α sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, C α -methyl amino acids, N α -methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

[0104] Neutrokin- α can also be differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc.

[0105] Additional post-translational modifications suitable for Neutrokin- α include, for example, N-linked or O-linked carbohydrate chains,

- 54 -

processing of N-terminal or C-terminal ends, attachment of chemical moieties to the amino acid backbone, chemical modifications of *N*-linked or *O*-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell expression. Neutrokin- α may also be modified with a detectable label, such as an enzymatic, fluorescent, radioisotopic, or affinity label to allow for detection and isolation of the protein.

[0106] The Neutrokin- α protein as used in the present invention is preferably provided in an isolated form, and preferably are substantially purified. By "isolated" is intended Neutrokin- α protein removed from its native environment. Thus, for example, a protein produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention. By way of example, a protein that is "substantially purified" is meant a protein that is substantially free from substances that limit its effect or produce undesired side effects. Usually, a substantially purified protein will be at least 90% pure. A recombinantly produced version of the Neutrokin- α protein can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

[0107] Neutrokin- α can be routinely produced, recovered and purified by methods known in the art (see, e.g., Example 5 and U.S. Application No. 10/270,487 which is hereby incorporated by reference in its entirety). Methods that can be used to purify Neutrokin- α include, but are not limited to, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

[0108] Also suitable are chemically modified derivatives of Neutrokin- α which may provide additional advantages such as increased solubility, stability, and *in vivo* or *in vitro* circulating time of the polypeptide, or

- 55 -

decreased immunogenicity (see, *e.g.*, U. S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The Neutrokin- α protein may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

[0109] The polymer may be of any molecular weight and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (*e.g.*, the duration of sustained release desired, the effects, if any, on biological activity, the ease in handling, the degree or lack of antigenicity, and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an average molecular weight of about 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10,000, 10,500, 11,000, 11,500, 12,000, 12,500, 13,000, 13,500, 14,000, 14,500, 15,000, 15,500, 16,000, 16,500, 17,000, 17,500, 18,000, 18,500, 19,000, 19,500, 20,000, 25,000, 30,000, 35,000, 40,000, 50,000, 55,000, 60,000, 65,000, 70,000, 75,000, 80,000, 85,000, 90,000, 95,000, or 100,000 kDa.

[0110] As noted above, the polyethylene glycol may have a branched structure. Branched polyethylene glycols are described, for example, in U.S. Patent No. 5,643,575; Morpurgo *et al.*, *Appl. Biochem. Biotechnol.* 56:59-72 (1996); Vorobjev *et al.*, *Nucleosides Nucleotides* 18:2745-2750 (1999); and Caliceti *et al.*, *Bioconjug. Chem.* 10:638-646 (1999), the disclosures of each of which are incorporated herein by reference.

[0111] The polyethylene glycol molecules or other chemical moieties should be attached to the protein with consideration of effects on functional or

- 56 -

antigenic domains of the protein. Furthermore, the polyethylene glycol molecules or other chemical moieties should be attached to the protein with consideration of possible effects on the chelator molecule of said Neutrokin- α conjugate or Neutrokin- α complex. There are a number of attachment methods available to those skilled in the art, *e.g.*, EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik *et al.*, Exp. Hematol. 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include, for example, lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues, glutamic acid residues, and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

[0112] As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to a proteins via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (*e.g.*, lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the protein or to more than one type of amino acid residue (*e.g.*, lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein.

[0113] One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (or peptide)

- 57 -

molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

[0114] As indicated above, pegylation of Neutrokin- α may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein either directly or by an intervening linker. Linkerless systems for attaching polyethylene glycol to proteins are described in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304 (1992); Francis *et al.*, *Intern. J. Hematol.* 68:1-18 (1998); U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466, the disclosures of each of which are incorporated herein by reference.

[0115] One system for attaching polyethylene glycol directly to amino acid residues of proteins without an intervening linker employs tresylated MPEG, which is produced by the modification of monomethoxy polyethylene glycol (MPEG) using tresylchloride ($\text{ClSO}_2\text{CH}_2\text{CF}_3$). Upon reaction of protein with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein. Thus, the invention includes protein-polyethylene glycol conjugates produced by reacting proteins of the invention with a polyethylene glycol molecule having a 2,2,2-trifluoroethanesulphonyl group.

[0116] Polyethylene glycol can also be attached to a protein using any of a number of different intervening linkers. For example, U.S. Patent No. 5,612,460, the entire disclosure of which is incorporated herein by reference,

- 58 -

discloses urethane linkers for connecting polyethylene glycol to a protein. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the protein by a linker can also be produced by reaction of a protein with a compound such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG-p-nitrophenolcarbonate, and various MPEG-succinate derivatives. A number additional polyethylene glycol derivatives and reaction chemistries for attaching polyethylene glycol to a protein are described in WO 98/32466, the entire disclosure of which is incorporated herein by reference. Pegylated Neutrokin- α produced using the reaction chemistries set out herein are included as being suitable within the scope of the present invention.

[0117] The number of polyethylene glycol moieties attached to each Neutrokin- α (*i.e.*, the degree of substitution) may also vary. For example, the pegylated proteins of the invention may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304 (1992).

Neutrokin- α Conjugate:

Chelator and Association of Chelator with Neutrokin- α Protein

[0118] Chelator molecules that may be used in the Neutrokin- α complexes and Neutrokin- α conjugates of the present invention are known in the art. For example, see Subramanian, R. and Meares, C.F., "Bifunctional Chelating Agents for Radiometal-labeled monoclonal

Antibodies," in *Cancer Imaging with Radiolabeled Antibodies* (D. M. Goldenberg, Ed.) Kluwer Academic Publications, Boston; Saji, H., "Targeted delivery of radiolabeled imaging and therapeutic agents: bifunctional radiopharmaceuticals," *Crit. Rev. Ther. Drug Carrier Syst.* 16:209-244 (1999); Srivastava S.C. and Mease R.C., "Progress in research on ligands, nuclides and techniques for labeling monoclonal antibodies," *Int. J. Rad. Appl. Instrum. B* 18:589-603 (1991); and Liu, S. and Edwards, D.S., "Bifunctional chelators for therapeutic lanthanide radiopharmaceuticals," *Bioconjug. Chem.* 12:7-34 (2001). Any chelator which can be covalently bound to said Neutrokin- α protein may be used according to the present invention. The chelator may further comprise a linker moiety that connects the chelating moiety to the Neutrokin- α protein.

[0119] In one embodiment, the chelator is an acyclic chelator such as diethylene triamine-N,N,N',N'',N''-pentaacetic acid (DPTA), analogues of DPTA, and derivatives of DPTA. As non-limiting examples, the chelator may be 2-(p-isothiocyanatobenzyl)-6-methyldiethylenetriaminepentaacetic acid (1B4M-DPTA, also known as MX-DTPA), 2-methyl-6-(rho-nitrobenzyl)-1,4,7-triazaheptane-N,N,N',N'',N''-pentaacetic acid (nitro-1B4M-DTPA or nitro-MX-DTPA); 2-(p-isothiocyanatobenzyl)-cyclohexyldiethylenetriaminepentaacetic acid (CHX-DTPA), or N-[2-amino-3-(rho-nitrophenyl)propyl]-trans-cyclohexane-1,2-diamine-N,N',N''-pentaacetic acid (nitro-CHX-A-DTPA).

[0120] In another embodiment, the chelator is an acyclic terpyridine chelator such as 6,6''-bis[[N,N,N',N''-tetra(carboxymethyl)amino]methyl]-4'-(3-amino-4-methoxyphenyl)-2,2':6',2''-terpyridine (TMT-amine).

[0121] In a specific embodiment, Neutrokin- α proteins of the invention are attached to macrocyclic chelators useful for conjugating radiometal ions, including but not limited to, ^{111}In , ^{177}Lu , ^{90}Y , ^{166}Ho , and ^{153}Sm , to polypeptides. In a specific embodiment, the radiometal ion which associates with the macrocyclic chelators attached to Neutrokin- α proteins of the invention is ^{111}In . In another preferred embodiment, the radiometal ion which

- 60 -

associates with the macrocyclic chelator attached to Neutrokin- α proteins of the invention is ^{90}Y . In specific embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA). In other specific embodiments, the DOTA is attached to the Neutrokin- α and/or Neutrokin- α SV polypeptide of the invention via a linker molecule. Examples of linker molecules useful for conjugating DOTA to a polypeptide are commonly known in the art - see, for example, DeNardo *et al.*, *Clin. Cancer Res.* 4(10):2483-90, 1998; Peterson *et al.*, *Bioconjug. Chem.* 10(4):553-7, 1999; and Zimmerman *et al.*, *Nucl. Med. Biol.* 26(8):943-50, 1999 which are hereby incorporated by reference in their entirety. In addition, U.S. Patents 5,652,361 and 5,756,065, which disclose chelating agents that may be conjugated to antibodies, and methods for making and using them, are hereby incorporated by reference in their entirety. Though U.S. Patent Nos. 5,652,361 and 5,756,065 focus on conjugating chelating agents to antibodies, one skilled in the art could readily adapt the method disclosed therein in order to conjugate chelating agents to other polypeptides.

[0122] Bifunctional chelators based on macrocyclic ligands in which conjugation is via an activated arm, or functional group, attached to the carbon backbone of the ligand can be employed as described by M. Moi *et al.*, *J. Amer. Chem. Soc.* 49:2639 (1989) (2-*p*-nitrobenzyl-1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid); S. V. Deshpande *et al.*, *J. Nucl. Med.* 31:473 (1990); G. Ruser *et al.*, *Bioconj. Chem.* 1:345 (1990); C. J. Broan *et al.*, *J. C. S. Chem. Comm.* 23:1739 (1990); and C. J. Anderson *et al.*, *J. Nucl. Med.* 36:850 (1995).

[0123] In one embodiment, the chelator is a macrocyclic chelator, such as polyazamacrocyclic chelators, optionally containing one or more carboxy, amino, hydroxamate, phosphonate, or phosphate groups. In another embodiment, the chelator is a chelator selected from the group consisting of DOTA, analogues of DOTA, and derivatives of DOTA.


[0124] In one embodiment, suitable chelator molecules include DOXA (1-oxa-4,7,10-triazacyclododecanetriacetic acid), NOTA

- 61 -

(1,4,7-triazacyclononanetriacetic acid), TETA (1,4,8,11-tetraazacyclotetradecanetetraacetic acid), and THT (4'-(3-amino-4-methoxy-phenyl)-6,6"-bis(N',N'-dicarboxymethyl-N-methylhydrazino)-2,2':6',2"-terpyridine), and analogs and derivatives thereof. See, *e.g.*, Ohmono *et al.*, *J. Med. Chem.* 35: 157-162 (1992); Kung *et al.*, *J. Nucl. Med.* 25: 326-332 (1984); Jurisson *et al.*, *Chem. Rev.* 93:1137-1156 (1993); and U.S. Patent No. 5,367,080. Other suitable chelators include chelating agents disclosed in U.S. Patent Nos. 4,647,447; 4,687,659; 4,885,363; EP-A-71564; WO89/00557; and EP-A-232751.

[0125] In another embodiment, suitable macrocyclic carboxylic acid chelators which can be used in the present invention include 1,4,7,10-tetraazacyclododecane-*N,N',N'',N'''*-tetraacetic acid (DOTA); 1,4,8,12-tetraazacyclopentadecane-*N,N',N'',N'''*-tetraacetic acid (15N4); 1,4,7-triazacyclononane-*N,N',N''*-triacetic acid (9N3); 1,5,9-triazacyclododecane-*N,N',N''*-triacetic acid (12N3); and 6-bromoacetamido-benzyl-1,4,8,11-tetraazacyclotetradecane-*N,N',N'',N'''*-tetraacetic acid (BAT).

[0126] In another embodiment, the chelator is a chelator having the Formula *I*:



wherein

each Q is independently hydrogen or (CHR⁵)_pCO₂R, preferably -CH₂-CO₂R, more preferably -CH₂COOH;

Q¹ is hydrogen or (CHR⁵)_wCO₂R, preferably CO₂R, more preferably COOH;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl, preferably hydrogen or C₁₋₄ alkyl, more preferably hydrogen;

with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen;

- 62 -

each R^5 independently is hydrogen, C_1 - C_4 alkyl or $-(C_1$ - C_2 alkyl)phenyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1, preferably 0;

m is an integer from 0 to 10 inclusive, preferably 0 to 1, more preferably 0;

p is 1 or 2, preferably 1;

r is 0 or 1, preferably 0;

w is 0 or 1, preferably 0;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato;

R^3 is selected from the group consisting of C_1 - C_4 alkoxy, $-OCH_2CO_2H$, hydroxy and hydrogen, preferably C_1 - C_4 alkoxy, more preferably methoxy;

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato;


with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof. Such a compound is described in U.S. Patent No. 5,652,361.

[0127] With reference to the chelator of Formula I, it is understood that the chelator portion of the Neutrokin- α conjugate of the present invention is formed from a molecule of Formula I. Accordingly, in one embodiment, the Neutrokin- α protein is bonded to an appropriate functional group on a compound of Formula I. For example, in one embodiment, the functional group designated R^2 is reacted with a Neutrokin- α protein to form a Neutrokin- α conjugate. By way of example, in one embodiment, R^2 is an isothiocyanato and reacts with a hydroxyl group on a serine residue of a

- 63 -

Neutrokin- α protein. In another embodiment, the functional group designated R^4 is reacted with a Neutrokin- α protein to form a Neutrokin- α conjugate.


[0128] In another specific embodiment, the chelator of said conjugate is formed from α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (MeO-DOTA-NCS), which has the structure of Formula **II**:

**II**

wherein "NCS" denotes a isothiocyanato group. A pharmaceutically acceptable salt or ester of α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid may also be used.

[0129] When forming the Neutrokin- α conjugate using a chelator as described above, the Neutrokin- α protein reacts with a functional group on the chelator to form a covalent bond and thus form the conjugate. With reference to a chelator according to Formula **I**, the Neutrokin- α protein will preferentially react with a functional group selected from the group consisting of amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl. Thus, the Neutrokin- α protein will preferentially form a bond with a functional group of either R^2 or R^4 . Thus, in accordance with the description of a chelator according to Formula **I**, in one embodiment, a Neutrokin- α conjugate has the structure of Formula **III**:

- 64 -



III

wherein n^1 is an integer from 1 to about 30, in another embodiment from 1 to about 20, in yet another embodiment 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, in yet another embodiment from about 1 to about 5, or in another embodiment about 1;

Q, Q^1 , R, R^2 , R^3 , R^5 , X, Y, n, m, p, r, and w are defined as above; and
NA is a Neutrokin- α protein;


wherein R^6 is a functional group resulting from the reaction of an amino acid residue of NA with a functional group selected from the group defined for R^4 above, *i.e.*, consisting of amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably isothiocyanato.

[0130] In one embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula **III**, wherein said NA is a mature, soluble Neutrokin- α protein or a fragment or variant thereof. In another embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula **III**, wherein said NA is a human mature, soluble Neutrokin- α protein or a fragment or variant thereof. In another embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula **III**, wherein said NA comprises, or alternatively consists of the sequence shown in Table 2 or a fragment or variant thereof.

[0131] With reference to a chelator according to Formula **II**, the Neutrokin- α protein, in one embodiment, reacts with an isothiocyanato group of the chelator. Thus, the Neutrokin- α protein, in one embodiment, forms a bond with the isothiocyanato group to form a thiocarbamate group or a

- 65 -

thiourea group, preferably a thiourea group. Thus, in accordance with the description of a chelator according to Formula *II*, a particular Neutrokin- α conjugate has the structure of Formula *IV*:

**IV**

or a pharmaceutically acceptable salt thereof, wherein NA is a Neutrokin- α protein;

n^1 is an integer from 1 to about 30, preferably from about 1 to about 20, preferably from about 1 to about 5, most preferably about 1; and

N' is a nitrogen atom from an amino acid residue, preferably the N-terminal amino acid residue or a lysine residue, of the Neutrokin- α protein.

[0132] In one embodiment, the Neutrokin- α protein of Formula *IV* is any Neutrokin- α protein as described herein. In one embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula *IV*, wherein said NA is a mature, soluble Neutrokin- α protein or a fragment or variant thereof. In another embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula *IV*, wherein said NA is a human mature, soluble Neutrokin- α protein (*i.e.*, amino acids 134-285 of SEQ ID NO:2) or a fragment or variant thereof. In another embodiment, the Neutrokin- α conjugate of the present invention has the structure of Formula *IV*, wherein said NA comprises the sequence shown in Table 1 or Table 2 or a fragment or variant thereof.

- 66 -

[0133] It is understood that other functional groups, as described above, may be used to link the chelator to the Neutrokin- α protein.

[0134] As used herein, "pharmaceutically acceptable salt" means any salt of a compound of Formulae *I-IV* which is sufficiently non-toxic to be useful in therapy or diagnosis of mammals. Thus, the salts are useful in accordance with this invention. Representative of those salts, which are formed by standard reactions, from both organic and inorganic sources include, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, palmoic, mucic, glutamic, d-camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, steric, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic acids and other suitable acids. Also included are salts formed by standard reactions from both organic and inorganic sources such as ammonium, alkali metal ions, alkaline earth metal ions, and other similar ions. Particularly preferred are the salts of a chelator where the salt is potassium, sodium, ammonium, or mixtures thereof.

[0135] Of course, the free acid of the compounds and conjugates of Formulae *I-IV* may be used, also the protonated form of the compounds, for example when the carboxylate is protonated and/or the nitrogen atoms are protonated, *e.g.*, when the HCl salt is formed.

[0136] The Neutrokin- α conjugate may contain more than an average of one chelator molecule per monomer of Neutrokin- α . In one embodiment, the Neutrokin- α conjugate contains about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, or 35 chelator molecules per monomer of Neutrokin- α . In another embodiment, the Neutrokin- α conjugate contains about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 chelator molecules per monomer of Neutrokin- α protein. In another embodiment, the Neutrokin- α conjugate contains about 1, 2, 3, 4, or 5 chelator molecules per monomer of Neutrokin- α protein. In another embodiment, the Neutrokin- α conjugate contains an average of about 1 chelator molecule per monomer of Neutrokin- α . In

- 67 -

another embodiment, the Neutrokin- α conjugate contains about 1.1 molecules per monomer of Neutrokin- α protein.

[0137] The Neutrokin- α conjugate may contain more than an average of one chelator molecule per monomer of Neutrokin- α . In one embodiment, the Neutrokin- α conjugate contains at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, or 35 chelator molecules per monomer of Neutrokin- α . In another embodiment, the Neutrokin- α conjugate contains at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 chelator molecules per monomer of Neutrokin- α protein. In another embodiment, the Neutrokin- α conjugate contains at least 1, 2, 3, 4, or 5 chelator molecules per monomer of Neutrokin- α protein. In another embodiment, the Neutrokin- α conjugate contains an average of at least 1 chelator molecule per monomer of Neutrokin- α .

[0138] The Neutrokin- α conjugate, as described above, comprises a Neutrokin- α protein and a chelator molecule covalently bonded together. The covalent bond between the protein and chelator is formed between an atom of the protein and an atom of the chelator. The atom of the protein that forms the covalent bond can be any suitable atom of the protein, such as a carbon, nitrogen, oxygen, and sulfur. Certain functional groups on the protein are preferred to form the covalent bond with the chelator. Such groups include an amino group at the terminus of the protein; a carboxy group at the terminus of the protein; an amino group on a lysine residue; a carboxy group on an aspartame or glutamate residue; a guanidino group on an arginine residue; a thiol group on a cysteine residue; a hydroxy group on a serine residue or threonine residue; an imidazole group of a histidine residue; a hydroxy group of a tyrosine; a hydroxy group of a tryptophan group; and an amide group of an asparagine or glutamine residue.

[0139] In one embodiment, the chelator is bonded to a lysine residue of the Neutrokin- α protein. In another embodiment, the chelator is bonded to one or more residues selected from the group consisting of LYS160, LYS173, LYS181, LYS184, LYS188, LYS204, LYS215, LYS216, LYS252, and LYS283.

- 68 -

- [0140] In another embodiment, the chelator is bonded to ALA134.
- [0141] In another embodiment, the chelator is bonded to an N-terminal amino acid of the Neutrokin- α protein.
- [0142] In another embodiment, the chelator is bonded to the C-terminus amino acid of the Neutrokin- α protein.
- [0143] Accordingly, in one embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS160. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS173. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS181. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS184. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS188. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS204. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS215. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS216. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS252. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to LYS283. In another embodiment, the Neutrokin- α conjugate of the present invention is a molecule according to Formula **IV**, wherein the chelator molecule is bonded to ALA134.

- 69 -

[0144] In a further embodiment, the Neutrokin- α conjugate, existing as a trimer, has a differential conjugation pattern. That is, each monomer of the trimer has a chelator molecule attached to a different amino acid. By way of example, in one embodiment, a Neutrokin- α conjugate trimer has one monomer with a chelator attached to LYS252, has another monomer with a chelator attached to LYS 283, and a third monomer with a chelator attached to ALA134.

[0145] As described earlier, Neutrokin- α can exist as a monomer protein or as multiple subunits associated together. In one embodiment, the Neutrokin- α conjugate is any of the specific embodiments described above wherein the conjugate is in the monomer form. In another embodiment, the Neutrokin- α conjugate is any of the specific embodiments described above wherein the conjugate is in the trimer form.

Neutrokin- α Complex

[0146] An additional embodiment of the present invention is directed to a Neutrokin- α complex comprising a Neutrokin- α conjugate and a metal ion, wherein said metal ion is associated with the chelator moiety of said Neutrokin- α conjugate. Specific embodiments of a Neutrokin- α complex of the invention include a complex comprising a Neutrokin- α conjugate, as set forth herein, and a metal ion. Herein "metal ion" refers to any ion that can associate with the chelator moiety of the invention as described herein.

Metal Ion

[0147] Any metal ion that associates with the chelating moiety of said Neutrokin- α conjugate may be used. Such a metal ion includes a metal ion selected from the group consisting of Ac, Ag, At, Au, Bi, Ce, Co, Cr, Cu, Dy, Er, Eu, Fe, Ga, Gd, Hg, Ho, In, La, Lu, Mn, Mo, Nd, Ni, Os, Pb, Pd, Pm,

- 70 -

Pr, Pt, Rb, Re, Rh, Ru, Sb, Sc, Si, Sm, Sn, Sr, Tb, Tc, Tl, Tm, V, W, Y, and Yb.

[0148] In one embodiment of the present invention, the metal ion of the complex is a radionuclide. Radionuclides useful in the Neutrokinine-alpha complex are known in the art. The radionuclides useful in the present invention include, but are not limited to, gamma-emitters, positron-emitters, x-ray emitters, fluorescence-emitters, beta-emitters, alpha-emitters, auger electron emitters, and electron and neutron-capturing agents. In one embodiment, a gamma-emitter, positron-emitter, x-ray emitter, and/or fluorescence-emitter is used for localization and/or therapy. In another embodiment, a beta-emitter, alpha-emitter, or an electron-capturing or neutron-capturing agent, such as boron or uranium, is used for therapy.

[0149] The radionuclide used in the complex of the present invention may be suitable for therapeutic, diagnostic, or both therapeutic and diagnostic purposes. Examples of appropriate metals include Ag, At, Au, Bi, Cu, Ga, Ho, In, Lu, Pb, Pd, Pm, Pr, Rb, Re, Rh, Sc, Sr, Tc, Tl, Y, and Yb. Examples of the radionuclide used for diagnostic purposes include Fe, Gd, ^{111}In , ^{67}Ga , or ^{68}Ga . In another embodiment, the radionuclide used for diagnostic purposes is ^{111}In , or ^{67}Ga . Examples of the radionuclide used for therapeutic purposes include ^{166}Ho , ^{165}Dy , ^{90}Y , $^{115\text{m}}\text{In}$, ^{52}Fe , or ^{72}Ga . In one embodiment, the radionuclide used for diagnostic purposes is ^{166}Ho or ^{90}Y . Examples of the radionuclides used for both therapeutic and diagnostic purposes include ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{175}Yb , or ^{47}Sc . In one embodiment, the radionuclide is ^{153}Sm , ^{177}Lu , ^{175}Yb , or ^{159}Gd .

[0150] Preferred metal radionuclides include, but are not limited to, ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{111}In , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{97}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{95}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{203}Pb and ^{141}Ce .

[0151] In a particular embodiment, the metal ion of the Neutrokinine-alpha complex is selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac .

- 71 -

- [0152] Moreover, according to the present invention, γ -emitting radionuclides, such as ^{99m}Tc , ^{111}In , ^{67}Ga , and ^{169}Yb may be used for diagnostic imaging, while complexes of β -emitters, such as ^{67}Cu , ^{111}Ag , ^{186}Re , and ^{90}Y are useful for the applications in tumor therapy. Also other useful radionuclides include γ -emitters, such as ^{99m}Tc , ^{111}In , ^{67}Ga , and ^{169}Yb , and β -emitters, such as ^{67}Cu , ^{111}Ag , ^{186}Re , ^{188}Re and ^{90}Y , as well as other radionuclides of interest such as ^{211}At , ^{212}Bi , ^{177}Lu , ^{86}Rb , ^{105}Rh , ^{153}Sm , ^{198}Au , ^{149}Pm , ^{85}Sr , ^{142}Pr , ^{214}Pb , ^{109}Pd , ^{166}Ho , ^{208}Tl , and ^{44}Sc .
- [0153] In another embodiment, paramagnetic metal ions that may be used according to the present invention include ions of transition and lanthanide metal, such as metals having atomic numbers of 21-29, 42, 43, 44, or 57-71, in particular ions of Cr, V, Mn, Fe, Co, Ni, Cu, La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, and Lu. The paramagnetic metals used in the composition for magnetic resonance imaging include the elements having atomic numbers of 22 to 29, 42, 44 and 58-70.
- [0154] In another embodiment, fluorescent metal ions that may be used according to the present invention include lanthanides, in particular La, Ce, Pr, Nd, Pm, Sm, Eu, Gd, Tb, Dy, Ho, Er, Tm, Yb, and Lu.
- [0155] In another embodiment, heavy metal-containing reporters that may be used according to the present invention may include atoms of Mo, Bi, Si, and W.
- [0156] In an additional embodiment, the Neutrokin- α complex of the invention comprises a metal ion selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac , and a conjugate according to Formula IV. In another embodiment, the Neutrokin- α complex of the invention comprises a metal ion selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac , and a conjugate according to Formula IV, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 or a fragment or variant thereof.
- [0157] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{90}Y and a conjugate according to Formula IV, wherein

- 72 -

said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0158] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{111}In and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0159] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{177}Lu and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0160] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{166}Ho and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0161] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{215}Bi and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0162] In an additional embodiment, the Neutrokin- α complex of the invention comprises ^{166}Ho and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

Method of Preparing Conjugate

[0163] An additional embodiment of the present invention is directed to a method of preparing a Neutrokin- α conjugate. The general procedure for preparing a Neutrokin- α conjugate comprises reacting a Neutrokin- α protein with a chelator. A suitable chelator will be able to react with the Neutrokin- α protein to form a covalent bond between the chelator and the Neutrokin- α protein.

- 73 -

[0164] Any chelator which is able to form a covalent bond with the Neutrokin- α protein may be used to form the Neutrokin- α conjugate. Such chelators are described herein or are otherwise known in the art, as are methods for preparing and/or attaching such chelators.


[0165] In one embodiment, a chelator to be used in the method according to the present invention is an activated chelator. The activated chelator contains a functional group that will readily react with a functional group on the protein, thereby forming a covalent bond between the protein and the chelator. Such functional groups are known in the art.

[0166] A suitable reactive functional group is a group that will react directly with carboxy, aldehyde, amine, alcohol, or sulfhydryl group on the Neutrokin- α protein. Such groups include, for example, active halogen containing groups including, for example, chloromethylphenyl groups and chloroacetyl ($\text{ClCH}_2\text{C}(=\text{O})-$) groups; activated 2-(leaving group substituted)-ethylsulfonyl and ethylcarbonyl groups such as 2-chloroethylsulfonyl and 2-chloroethylcarbonyl; vinylsulfonyl; vinylcarbonyl; epoxy; isocyanato; isothiocyanato; aldehyde; aziridine; succinimidoxycarbonyl; activated acyl groups such as carboxylic acid halides; and mixed anhydrides.

[0167] A chelator which can be used in the present method includes, for example, any specific chelator as described above for the Neutrokin- α conjugate. Moreover, in another embodiment, the chelator used in the method of the invention is a chelator which can be used to prepare any specific Neutrokin- α conjugate as described above. In one embodiment, the chelator is an activated chelator selected from the group consisting of DOTA, analogues of DOTA, and derivatives of DOTA, wherein each of said DOTA, analogues of DOTA, and derivatives of DOTA contains a suitable activating group which enables the chelator molecule to be covalently bonded to the Neutrokin- α protein.

[0168] In one embodiment, the activated chelator for use in preparing a conjugate according to the present invention is an activated chelator having the Formula:

- 74 -



wherein

each Q is independently hydrogen or (CHR⁵)_pCO₂R, preferably -CH₂-CO₂R, more preferably -CH₂CO₂H;

Q¹ is hydrogen or (CHR⁵)_wCO₂R, preferably CO₂R, more preferably COOH;

each R independently is hydrogen, benzyl or C₁-C₄ alkyl; preferably hydrogen or C₁-C₄ alkyl, more preferably hydrogen;

with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen;

each R⁵ independently is hydrogen, C₁-C₄ alkyl or -(C₁-C₂ alkyl)phenyl;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1; preferably 0

m is an integer from 0 to 10 inclusive, preferably 0 to 1, more preferably 0;

p is 1 or 2, preferably 1;

r is 0 or 1, preferably 0;

w is 0 or 1, preferably 0;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R² is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato;


- 75 -

R^3 is selected from the group consisting of C_1 - C_4 alkoxy, $-OCH_2CO_2H$, hydroxy and hydrogen, preferably C_1 - C_4 alkoxy, more preferably methoxy;

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof. Such a compound is described in U.S. Patent No. 5,652,361.

In another embodiment, the activated chelator for use in preparing a conjugate according to the present invention is α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, which has the following structure:



wherein "NCS" denotes an isothiocyanato group, and which is also known as MeO-DOTA-NCS. A salt or ester of α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid may also be used.

[0169] In a specific embodiment of the invention, the Neutrokin- α protein is trimer of Neutrokin- α monomeric subunits wherein each Neutrokin- α subunit consists of amino acids 134-285 of human Neutrokin- α and having the sequence listed in Table 1 (SEQ ID NO:2). In additional specific embodiments, the Neutrokin- α protein used in the method of the invention is any one of the specific Neutrokin- α proteins described above. In additional specific embodiments, the Neutrokin- α conjugate prepared according to the present method is any one of the specific Neutrokin- α conjugates described above. In particular, the Neutrokin-

- 76 -

alpha protein used in the present method is in the form of a trimer, as described above.

[0170] In one embodiment, the method of preparing a Neutrokin- α conjugate comprises mixing, agitating, or preparing a solution, said solution comprising a Neutrokin- α protein and a chelator. In an additional embodiment, the method of preparing a Neutrokin- α conjugate comprises mixing, agitating, or preparing a solution comprising a Neutrokin- α protein, a chelator, citrate buffer, HEPES buffer, and sterile water. In an additional embodiment, the solution further comprises NaOH. In an additional embodiment, the solution has a pH of about 8.5.

[0171] In an additional embodiment, the method of preparing the conjugate comprises a) mixing, agitating, or preparing a solution comprising a Neutrokin- α protein and a chelator, at a temperature of about 0 °C to about 50 °C for about 0.5 hours to about 24 hours, wherein said solution has a pH of about 8.0 to about 9.0; and b) optionally adding a quenching agent.

[0172] In one embodiment of the above process, the solution is mixed, agitated or prepared at a temperature of about 0 °C to about 50 °C. In another embodiment of the above process, the solution is mixed, agitated or prepared at a temperature of about 20 °C to about 30 °C. The temperature at which the solution is mixed, agitated, or prepared can be about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 °C.

[0173] Any suitable chelator as described above may be used in an embodiment of the present process. Chelators as described herein may be purchased for example, from Dow Chemical Company, (Midland, Michigan).

[0174] When making a Neutrokin- α conjugate it is important to consider the ratio of the number of moles of chelator molecules in the reaction compared to the number of moles of sites to which the chelator molecule may attach in the Neutrokin- α protein (herein referred to as the molar ratio of chelator to chelator bonding sites). The molar ratio of chelator to chelator bonding sites used in the preparation of the conjugate can vary depending on the number of chelator bonding sites in the Neutrokin- α protein

- 77 -

component of the Neutrokin- α conjugate. For example, if the Neutrokin- α protein component is a trimer of Neutrokin α monomers, wherein each monomer consists of the Neutrokin- α protein of SEQ ID NO:3, and the chelator is bonding to either the N terminus or to Lysine residues in the Neutrokin- α trimer, and it is desired that the Neutrokin- α conjugate product comprises, on average, 3 chelator molecules on the trimeric Neutrokin- α protein, it would be desirable to use a molar ratio of chelator to chelator binding sites in Neutrokin- α protein between 12:1 and 10:1 in the reaction (see, *e.g.*, Example 5).

[0175] In specific embodiments, molar ratio of chelator to chelator binding sites in Neutrokin- α protein is less than or equal to 1000:1. In other specific embodiments, molar ratio of chelator to chelator binding sites in Neutrokin- α protein is less than or equal to 100:1. In preferred embodiments, molar ratio of chelator to chelator binding sites in Neutrokin- α protein is 20:1, 15:1, 12:1, 11:1, 10:1 or 5:1.

[0176] In the final Neutrokin- α conjugate product, the molar ratio of chelator to Neutrokin- α protein monomer can be from about 10:1 to about 1:10. In one embodiment, the molar ratio of chelator to Neutrokin- α protein monomer may be 5:1 or about 1:5; In a preferred embodiment, from about 1:3 to about 3:1. In another preferred embodiment, the molar ratio of chelator to Neutrokin- α protein monomer in the final Neutrokin- α conjugate product, is about 1:1. In another embodiment, the molar ratio of chelator to Neutrokin- α protein monomer in the final conjugate product is about 1.1:1. The molar ratio of chelator to Neutrokin- α protein monomers refers to the ratio of the number of chelator molecules to the number of Neutrokin- α protein molecules (monomers) in the product. As is known in the art, Neutrokin- α proteins can form bound oligomers, for example trimers. Thus, according to the present invention, each Neutrokin- α trimer is equal to three Neutrokin- α protein monomers when calculating the ratio of chelator to Neutrokin- α protein monomers.

- 78 -

[0177] When preparing a Neutrokin- α conjugate, the solution can be mixed, agitated, or allowed to stand for a variable number of hours, depending upon a number of variables, such factors including the identity of the chelator, the identity of the Neutrokin- α protein, the temperature of the reaction, the pH, the identity of the buffer, the molar ratio of the reactants, the purity of the available reagents, the presence of a catalyst or activator, and other factors which would be evident to, and routinely manipulated by, one of skill in the art to achieve a desired result. In one embodiment of the above process, the solution comprising a Neutrokin- α protein and a chelator is mixed, agitated, or allowed to stand for about 0.5 hours to about 24 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 1 hour to about 20 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 2 hour to about 10 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 3 hour to about 5 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 4 hours. In other embodiments, the solution is mixed, agitated, or allowed to stand for about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 hours.

[0178] The pH of the solution used to prepare the Neutrokin- α conjugate can vary. In one embodiment, the pH is about 6, 7, 8, 9, or 10. In another embodiment, said solution has a pH of about 7 to about 10. In another embodiment, said solution has a pH of about 7.5 to about 9.5. In another embodiment, said solution has a pH of about 8 to about 9. In another embodiment, said solution has a pH of about 8.5.

[0179] The solution used to prepare the Neutrokin- α conjugate may further comprise a buffer. Buffers are well-known in the art and may be routinely applied to maintain the desired pH of the solutions used in making and/or using the Neutrokin- α conjugates and/or Neutrokin- α complexes of the invention. Suitable buffers for use in the preparation of a Neutrokin- α include, for example, those described below for the method of preparing a Neutrokin- α conjugate. In one embodiment, the buffer is a

- 79 -

citrate buffer or an acetate buffer. In another embodiment, the buffer includes an acetate buffer having a concentration of about 1 to about 50 mM and having a NaCl concentration of about 1 to about 500 mM. In another embodiment, the buffer includes an acetate buffer having a concentration of about 10 mM and having a NaCl concentration of about 140 mM. Suitable acetate buffers include acetate buffers having a concentration of about 1, 20, 25, 50, 75, 100, 200, 250, 300, 400, or 500mM. Suitable buffers and solutions include those having a NaCl concentration of about 1, 50, 75, 100, 125, 140, 150, 175, 200, 225, 250, 275, 300, 350, 400, 450, or 500 mM. An additional suitable buffer is a HEPES buffer, in particular a HEPES buffer having a concentration of about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 mM. In an additional embodiment, the solution comprises a HEPES buffer having a concentration of about 50 mM. In another embodiment, a citrate buffer is used, having a concentration of about 1 to about 100 mM, or about 1, 2, 5, 7, 10 15, or 20 mM.

[0180] The concentration of the Neutrokin- α protein used in the method of the invention may vary. In one embodiment, the concentration of Neutrokin- α protein in the solution is from about 0.1 mg/mL to about 10 mg/mL. In another embodiment, the Neutrokin- α concentration is about 0.5 mg/mL to about 5 mg/mL. In other specific embodiments, the Neutrokin- α concentration is about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, or 2.5 mg/mL. In another specific embodiment, the concentration of Neutrokin- α protein in the solution is about 1 mg/mL to about 2 mg/mL.

[0181] A reagent which facilitates the formation of the conjugate can also be used according to the methods of the invention. Such reagents typically activate a chelator so that it reacts more readily with a protein. Alternatively, the reagent may activate the protein to react more readily with the chelator. Examples of such reagents are known in the art and include dicyclohexylcarbodiimide (DCC), diethylazodicarboxylate (DEAD), and diisopropylazodicarboxylate (DIAD).

- 80 -

[0182] According to the method of the present invention, a quenching agent may optionally be added to the reaction solution after the Neutrokin- α conjugate is prepared to a sufficient degree. Suitable quenching agents are known in the art. In one embodiment, the quenching agent is glycine. In a specific embodiment, a solution comprising a glycine buffer, e.g., glycine HCl buffer, is used as a quenching agent. By way of example, after the reaction solution comprising the Neutrokin- α protein and the chelator has been mixed, agitated, or allowed to stand for a sufficient amount of time, e.g., 1-10 hours, 3-5 hours, or 1-3 hours, a solution comprising a glycine buffer is added to the reaction solution to stop the reaction. The reaction solution is then allowed to stand for an additional amount of time, e.g., about 0.25, 0.5, 0.75, 1, or 1.5 hours.

[0183] In a specific embodiment, the method of preparing a Neutrokin- α conjugate comprises mixing, agitating, or preparing a solution comprising:

a Neutrokin- α protein, wherein said protein comprises, or alternatively consists of a trimer of Neutrokin- α proteins, each consisting of amino acids 134-285 of SEQ ID NO: 2 and has a concentration of about 1 to about 2 mg/mL;

a MeO-DOTA-NCS in an amount such that the ratio MeO-DOTA-NCS molecules to lysine residues in a Neutrokin- α protein is about 12 to 1;

a citrate buffer having a concentration of about 0.5 to about 20 mM;

NaCl having a concentration of about 1 to about 200 mM;

HEPES buffer having a concentration of about 10 to about 500 mM;

and

sterile water.

[0184] In an additional embodiment, the method of the present invention comprises:

preparing a first solution comprising a Neutrokin- α protein and a citrate buffer;

- 81 -

adding a second solution comprising a HEPES buffer to said first solution;

adding a third solution comprising a chelator, for example MeO-DOTA-NCS, and NaOH to said first solution;

optionally adjusting the pH of said first solution to about 8.5;

mixing, agitating, or allowing to stand said first solution at about 25 °C for about 3 to about 5 hours; and

optionally, adding quenching agent, *e.g.*, a glycine buffer, to said first solution.

[0185] In a further embodiment, the mixing, agitating, or allowing to stand step is performed for about 4.5 to about 5.5 hours. In a further embodiment, the reaction solution can be incubated for an additional period of time after addition of the quenching agent.

[0186] In a further embodiment, the process of the invention is used to prepare a conjugate according to Formula IV.

[0187] The method of preparing a Neutrokin- α conjugate according to the present invention further optionally comprises a process of purifying said conjugate. A number of known methods may be used to purifying said protein conjugate; including but not limited to chromatography.

[0188] In one embodiment, the purifying step uses normal flow filtration or tangential flow filtration. In another embodiment, the process for purifying the conjugate according to the present invention is a diafiltration method (see, for example, Example 5).

[0189] In one embodiment, the Neutrokin- α conjugate is purified by HPLC, *e.g.*, reverse phase HPLC.

Method of Preparing Complex

[0190] An additional embodiment of the present invention is directed to a method of preparing a Neutrokin- α complex. In one embodiment, a Neutrokin- α complex may be prepared by reacting a Neutrokin- α conjugate, as described herein, with a metal ion, such as a radionuclide, which

- 82 -

is able to associate noncovalently with the conjugate. In another embodiment, the metal ion associates noncovalently with the chelator moiety of said Neutrokin- α conjugate. The reaction between the Neutrokin- α complex and the metal ion may occur in solution, such as in a suitable buffered solution.

[0191] In one embodiment, a method for preparing a Neutrokin- α complex comprises reacting a Neutrokin- α conjugate with a radionuclide. Such a method comprises mixing, agitating, or preparing a solution comprising a Neutrokin- α conjugate and a radionuclide. In one embodiment, said solution further comprises a buffer.

[0192] In another embodiment, the Neutrokin- α conjugate used in the present method is any specific Neutrokin- α conjugate as described above. In a particular embodiment, the Neutrokin- α conjugate comprises a Neutrokin- α protein comprising, or alternatively consisting of, amino acids 134-285 of SEQ ID NO:2 and chelator formed from MeO-DOTA-NCS.

[0193] In another embodiment, a method for preparing a Neutrokin- α complex comprises reacting a Neutrokin- α protein with a chelator complexed with a metal ion. In this embodiment, a chelator is first complexed with a metal ion according to known procedures in the art. For example, see U.S. Patent No. 5,654,361. The chelator can be an activated chelator as described above. After the chelator is complexed with the metal ion and thereby forming a chelator-metal ion complex, the Neutrokin- α protein is reacted with the chelator-metal ion complex, using a procedure as described above for preparing a Neutrokin- α conjugate or using another procedure known in the art. According to the method, the solution comprising the Neutrokin- α protein and the chelator-metal ion complex is mixed, agitated, or prepared, thereby forming said Neutrokin- α complex.

[0194] When preparing a Neutrokin- α complex, the solution can be mixed, agitated, or allowed to stand for a variable number of hours, depending upon a number of variables, such factors including the identity of the chelator, the identity of the Neutrokin- α protein, the identity of the metal ion, the

- 83 -

temperature of the reaction, the pH, the identity of the buffer, the molar ratio of the reactants, the purity of the available reagents, the presence of a catalyst, or activator, and other factors which would be evident to one of skill in the art. The solution comprising a Neutrokin- α conjugate and a metal ion can be mixed, agitated, or allowed to stand for any amount of time that permits sufficient formation of the Neutrokin- α complex. In one embodiment of the above process, the solution is mixed, agitated, or allowed to stand for about 0.5 minutes to about 24 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 1 hour to about 20 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 2 hours to about 10 hours. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 1 minute to about 1 hour. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 1 minute to about 30 minutes. In another embodiment, the solution is mixed, agitated, or allowed to stand for about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 55 minutes.

[0195] The pH of the solution used to prepare the Neutrokin- α complex of the invention can vary. In one embodiment, the pH is about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. In another embodiment, said solution has a pH of about 4 to about 8. In another embodiment, said solution has a pH of about 5 to about 7. In another embodiment, said solution has a pH of about 5.5 to about 7. In another embodiment, said solution has a pH of about 6.5.

[0196] The solution used to prepare the Neutrokin- α complex may further comprise a buffer. Buffers that may be used according to the present invention are well-known in the art. Suitable buffers for use in the preparation of a Neutrokin- α complex include, but are not limited to, citrate, acetate, phosphate, carbonate, diphosphate, glycyl-glycine-piperazine-2HCl-NaOH; MES-NaOH-NaCl; TRIS-malic acid-NaOH; MES-NaOH; ACES-NaOH-NaCl; BES-NaOH-NaCl; MOPS-NaOH-NaCl; TES-NaOH-NaCl; MOPS-KOH; HEPES-NaOH-NaCl; TRIS-HCl; HEPPSO-NaOH; TAPS-NaOH-NaCl; HEPPS (EPPS)-NaOH; citric acid-disodiumhydrogenphosphate; boric

- 84 -

acid-citric acid-potassium dihydrogen phosphate-Diethyl- barbituric acid-NaOH; citric acid-sodium citrate; sodium acetate-acetic acid; potassium hydrogenphthalate-NaOH; cacodylic acid sodium salt-HCl; potassium dihydrogen phosphate-disodium hydrogenphosphate; potassium dihydrogen-phosphate-NaOH; sodium dihydrogen phosphate- disodium hydrogen phosphate; imidazole-HCl; sodium tetraborate-boric acid; 2-amino-2-methyl-1,3-propanediol-HCl; diethanolamine-HCl; potassium chloride-boric acid-NaOH; boric acid-NaOH-KCl; glycine-NaOH; sodium bicarbonate, and sodium carbonate-sodium hydrogen carbonate.

[0197] In specific embodiments, the solution used to prepare the Neutrokinine-alpha complex may comprise sodium acetate and sodium bicarbonate. In specific embodiments, the solution used to prepare the Neutrokinine-alpha complex may comprise sodium acetate in a concentration range of about 50 mM to about 300mM, preferably 200-250 mM, and sodium bicarbonate in a concentration range of about 50 mM to about 300mM, preferably 50-100mM. In specific embodiments, the solution used to prepare the Neutrokinine-alpha complex may comprise about 220 mM sodium acetate and about 75 mM sodium bicarbonate. In specific embodiments, the solution used to prepare the Neutrokinine-alpha complex may comprise 275 mM sodium acetate and 377 mM sodium bicarbonate in a ratio of about 1 to 10 to about 10 to 1. In specific embodiments, the solution used to prepare the Neutrokinine-alpha complex may further comprise 275 mM sodium acetate and 377 mM sodium bicarbonate in a ratio of about 4 to 1.

[0198] The present invention encompasses varying the molar ratio of metal ions to chelator moieties on the Neutrokinine-alpha conjugate in the preparation of a Neutrokinine-alpha complex wherein a Neutrokinine-alpha conjugate is reacted with a metal ion. In one embodiment, the ratio of metal ions to chelator moieties in Neutrokinine-alpha conjugate is less than or equal to 1:100. In another embodiment, the ratio of metal ions to chelator moieties in Neutrokinine-alpha conjugate is less than or equal to 1:80. In another embodiment, the ratio of metal ions to chelator moieties in Neutrokinine-alpha

conjugate is less than or equal to 1:50. In a preferred embodiment, the ratio of metal ions to chelator moieties in Neutrokin- α conjugate is exactly or approximately 1:30, 1:25, 1:20, 1:15, 1:10, 1:5 or 1:1. In a specific preferred embodiment, the ratio of metal ions to chelator moieties in Neutrokin- α conjugate is exactly or approximately 1:20.

[0199] In certain embodiments, preparation of a Neutrokin- α complex wherein a Neutrokin- α protein is reacted with a chelator-metal ion complex, the molar ratio of chelator-metal ion complex to chelator binding sites in Neutrokin- α protein is less than or equal to 1000:1. In other specific embodiments, molar ratio of chelator-metal ion complex to chelator binding sites in Neutrokin- α protein is less than or equal to 100:1. In preferred embodiments, molar ratio of chelator-metal ion complex to chelator binding sites in Neutrokin- α protein is 20:1, 15:1, 12:1, 11:1, 10:1 or 5:1.

[0200] Radionuclides which can be used in the present invention are known in the art, many of which are described and listed above and are available commercially. Additionally, several known methods can be used to prepare the radionuclides for use in the present invention. In one embodiment, the metal ion is in the form of a salt, for example a chloride salt. Such salts are known in the art. In another embodiment, the metal ion salt is yttrium chloride or yttrium acetate.

[0201] When preparing the complex of the present invention according to one embodiment of the invention, other metal ions which could compete for complexation with the chelator are not present in significant amounts in the solution. For example, when preparing a Neutrokin- α complex comprising ^{90}Y , Fe^{3+} is not present in the solution in a significant amount or concentration. As used with reference to the one or more competing metal ions in the present process, the phrase "significant amount or concentration" refers to an amount or concentration which significantly interferes, retards, delays, inhibits, or prevents preparation of the Neutrokin- α complex.

[0202] The method of preparing a Neutrokin- α complex may further comprise a step of removing excess metal ion from the reaction solution

- 86 -

and/or the Neutrokin- α complex. In one embodiment, such a step comprises adding a secondary chelating agent which is able to complex with excess metal ion in the solution. Such a secondary chelating agent may include one or more known chelating agents, for example, DTPA, EDTA, and MeO-DOTA-glycine. In one embodiment, MeO-DOTA-glycine is used as a secondary chelating agent. In another embodiment, DTPA is used as a secondary chelating agent.

[0203] The method of preparing a Neutrokin- α complex may further comprise a step of removing excess metal ion from the reaction solution comprising the chelator-metal ion complex. In one embodiment, after the chelator-metal ion complex is formed and is in solution, a secondary chelating agent which is able to complex with excess metal ion is added. The solution is then eluted through a DEAE-cellulose anion exchange resin (Sigma Chemical Co., St. Louis, MO), which has been converted to acetate form to purify the neutral species chelator-metal ion complex from the charged species, *i.e.*, secondary chelator-metal ion complex. The purified chelator-metal ion complex is then used to prepare the Neutrokin- α complex as described herein.

[0204] The secondary chelating agent can be added to the reaction mixture as a solid or as a solution. In one embodiment, the secondary chelating agent is added in a buffered solution. In another embodiment of the present invention, a buffer comprising an acetate buffer having a concentration of about 1 to about 50 mM, preferably about 10 mM, having a NaCl concentration of about 1 to about 500 mM, preferably about 140 mM, having a HSA concentration of about 1% to about 20%, preferably about 7% to about 8%, more preferably about 7.5%, having a pH of about 3-8, preferably about 6, and having a DTPA concentration of about 0.01 mM to about 100 mM, preferably about 1 mM, is added to the reaction solution after the Neutrokin- α complex is formed to a satisfactory level of completion.

[0205] In one embodiment, a solution, having a NaCl concentration of about 1 to about 500 mM, preferably about 140 mM, having a pH of about 3-8,

- 87 -

preferably about 6, and having a DTPA concentration of about 0.01 mM to about 100 mM, preferably about 2 mM, and comprising from about 0% to 20% sodium ascorbate, preferably 7-10% sodium ascorbate, is added to the reaction solution after the Neutrokin- α complex is formed to a satisfactory level of completion.

[0206] In one embodiment, the secondary chelating agent is added about 5 minutes after the formation of the Neutrokin- α complex. In other embodiments, the secondary chelating agent is added about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, or 60 minutes after formation of the Neutrokin- α complex. In other embodiments, the secondary chelating agent is added at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, or 60 minutes after formation of the Neutrokin- α complex.

[0207] In another embodiment, the step of removing excess metal ion from the reaction solution and/or the Neutrokin- α complex comprises subjecting the reaction solution and/or the Neutrokin- α complex to centrifugation. Subjecting the reaction solution and/or the Neutrokin- α complex to centrifugation can remove non-chelated metal ions.

[0208] In another embodiment, the step of removing excess metal ion from the reaction solution and/or the Neutrokin- α complex comprises washing the excess metal ion from the reaction solution and/or the Neutrokin- α complex with a buffered solution. The washing may be repeated one or more times, as is necessary. In one embodiment, the washing is repeated two or three times. In one embodiment, the washing is repeated at least two or three times.

[0209] In one embodiment of the present method, two or more methods of removing excess metal ion are used in combination to remove excess metal ion from the step of removing excess metal ion from the reaction solution and/or the Neutrokin- α complex.

[0210] In another embodiment of the present method, for radiopharmaceutical and radiotherapy applications, the Neutrokin- α complex is prepared from a metal in an oxidation state different from that of the desired complex. In this

- 88 -

case, either a reducing agent or an oxidizing agent, depending on the oxidation state of the metal used and the oxidation state of the desired final product, is added to the reaction mixture to bring the metal to the desired oxidation state. The oxidant or reductant can be used to form an intermediate complex in the desired oxidation state but with labile ligands. These labile ligands can then be displaced by the desired chelating moiety of the present invention. In another embodiment, the labile ligands are added to the reaction mixture along with the reductant or oxidant and the desired ligand to achieve the change to the desired oxidation state and chelation to the desired metal in a single step.

[0211] In a specific embodiment of the present invention, the method of preparing a Neutrokin- α complex comprises mixing, agitating, or preparing a solution comprising a Neutrokin- α conjugate according to Formula *IV* and metal ion selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac , wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0212] In an additional embodiment, the method of preparing a Neutrokin- α complex comprises mixing, agitating, or preparing a solution comprising ^{111}In and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0213] In an additional embodiment, the method of preparing a Neutrokin- α complex comprises mixing, agitating, or preparing a solution comprising ^{177}Lu and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0214] In an additional embodiment, the method of preparing a Neutrokin- α complex comprises mixing, agitating, or preparing a solution comprising ^{90}Y and a conjugate according to Formula *IV*, wherein said Neutrokin- α protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

- 89 -

[0215] In an additional embodiment, the method of preparing a Neutrokin-alpha complex comprises mixing, agitating, or preparing a solution comprising ^{215}Bi and a conjugate according to Formula *IV*, wherein said Neutrokin-alpha protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0216] In an additional embodiment, the method of preparing a Neutrokin-alpha complex comprises mixing, agitating, or preparing a solution comprising ^{166}Ho and a conjugate according to Formula *IV*, wherein said Neutrokin-alpha protein comprises, or alternatively consists of, amino acids 134-285 of the sequence shown in Table 1 (SEQ ID NO:2).

[0217] In a further embodiment of the present invention, the method of preparing a Neutrokin-alpha complex comprises:

- preparing a first solution comprising a Neutrokin-alpha conjugate;
- adding to said first solution a second solution comprising a metal ion capable of complexing with said Neutrokin-alpha conjugate;
- mixing, agitating, or allowing to stand said first solution; and
- optionally, adding a secondary chelating agent, such as DTPA, to complex with uncomplexed metal ion.

[0218] In an additional embodiment, the method of preparing a Neutrokin-alpha complex comprises:

- combining a first solution, second solution, and a third solution, wherein
- said first solution comprises a Neutrokin-alpha conjugate,
- said second solution comprises an acetate buffer, and
- said third solution comprises $^{90}\text{YCl}_3$;
- mixing, agitating, or allowing to stand the combined solutions; and
- adding to the combined solution a fourth solution, said fourth solution comprising MeO-DOTA-NCS, human serum albumin (HSA), acetate buffer, and NaCl.

Compositions

[0219] A further embodiment of the present invention is a composition comprising the conjugate or complex as described above. In one embodiment, a composition according to the present invention comprises a Neutrokin- α conjugate and an acceptable carrier, preferably a pharmaceutically acceptable carrier. Suitable acceptable carriers include any liquid or solvent in which the conjugate or complex can be dissolved or suspended. Suitable pharmaceutically acceptable carriers are well-known in the art. A composition comprising a Neutrokin- α conjugate may further comprise a metal ion, preferably a radionuclide.

[0220] In an additional embodiment, a composition according to the present invention comprises a Neutrokin- α complex and an acceptable carrier, preferably a pharmaceutically acceptable carrier. Suitable acceptable carriers include any liquid or solvent in which the conjugate or complex can be dissolved or suspended. Suitable pharmaceutically acceptable carriers are known in the art.

[0221] In an additional embodiment, a composition according to the present invention comprises a Neutrokin- α conjugate, a metal ion, and an acceptable carrier, preferably a pharmaceutically acceptable carrier. Suitable acceptable carriers include any liquid or solvent in which the conjugate or complex can be dissolved or suspended. Suitable pharmaceutically acceptable carriers are known in the art.

[0222] The concentration of the Neutrokin- α conjugate or Neutrokin- α complex in the composition of the present invention can vary. For example, the concentration of Neutrokin- α conjugate or complex can be from about 0.1 $\mu\text{g/mL}$ to about 100 mg/mL , preferably from about 0.1 mg/mL to about 10 mg/mL , more preferably less than 4 mg/mL . In specific embodiments, the concentration of Neutrokin- α conjugate or Neutrokin- α complex in the composition is exactly or approximately 0.2 mg/mL . In specific embodiments, the concentration of Neutrokin- α conjugate or

- 91 -

Neutrokin- α complex in the composition is exactly or approximately 2.0 mg/mL.

[0223] For compositions containing a radionuclide, either all or some chelator moieties may contain a metal ion. The amount of radionuclide incorporated into a Neutrokin- α complex or chelator-metal ion complex may be calculated as described in Example 6 and compared to expected values. For example, it may be desirable to design the Neutrokin- α conjugate of the invention such that it comprises an excess number of chelator moieties compared to the number of metal ions that are desired to be incorporated into the final Neutrokin- α complex. The advantage of this approach would be that the excess number of chelators would ensure that all metal ions are complexed into Neutrokin- α complexes thereby eliminating the need for steps to remove excess metal ions in the preparation of Neutrokin- α complexes. In a specific embodiment it is contemplated that approximately every chelator moiety in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In another specific embodiment, it is contemplated that approximately 1 out of every 3 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In another specific embodiment, it is contemplated that approximately 1 out of every 5 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In another specific embodiment, it is contemplated that approximately 1 out of every 10 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific and preferred embodiment, it is contemplated that approximately 1 out of every 20 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific embodiment, it is contemplated that approximately 1 out of every 50 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific embodiment, it is contemplated that approximately 1 out of every 100 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion.

[0224] In another specific embodiment, it is contemplated that at least 1 out of every 3 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In another specific embodiment, it is contemplated that at least 1 out of every 5 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In another specific embodiment, it is contemplated that at least 1 out of every 10 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific and preferred embodiment, it is contemplated that at least 1 out of every 20 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific embodiment, it is contemplated that at least 1 out of every 50 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion. In a specific embodiment, it is contemplated that at least 1 out of every 100 chelator moieties in a solution of Neutrokin- α chelator becomes complexed with a metal ion.

[0225] The conjugates of this invention, and in some instances the complexes of this invention, may be employed as a formulation. The formulation comprises a Neutrokin- α conjugate or Neutrokin- α complex and a physiologically acceptable carrier, excipient, or vehicle therefore. Thus, the formulation may consist of a physiologically acceptable carrier with a Neutrokin- α complex (Neutrokin- α protein + chelator moiety(ies) + metal ion), Neutrokin- α conjugate (Neutrokin- α protein + chelator moiety(ies)) or Neutrokin- α protein. The methods for preparing such formulations are well known. The formulation may be in the form of a suspension, injectable solution, or other suitable formulation. Physiologically acceptable suspending media, with or without adjuvants, may be used.

[0226] A composition described herein optionally further comprises one or more known stabilizers. The use of a stabilizer is particularly useful in compositions comprising a radionuclide. As is known in the art, one concern when administering any radiopharmaceutical is the potential for radiolytic degradation of the organic molecule present in the formulation, which may

- 93 -

alter the biodistribution of the radioisotope or result in toxic by-products. Neither of these events is desirable. When high amounts of radioactivity are needed, there is the increased potential for radiation damage to the organic molecule. This degradation is more likely to occur when therapeutic radionuclides are used which are designed to deliver high radiation doses.

[0227] Examples of stabilizers that are suitable for use in the present compositions and methods include free radical inhibitors, such as benzyl chloride and ascorbic acid. See, *e.g.*, H. Ikebuchi *et al.*, *Radioisotopes* 26:451-457 (1977); B. J. Floor *et al.*, *J. Pharm. Sci.* 74:197-200 (1985); A. Rego, *et al.*, *J. Pharm. Sci.* 71:1219-23 (1982); and U.S. Patent Nos. 5,843,396 and 5,384,113.

[0228] An additional embodiment of the present invention is a pharmaceutical composition comprising a Neutrokin- α conjugate and one or more pharmaceutically acceptable carriers. A preferred composition of the present invention is a pharmaceutical composition comprising a Neutrokin- α conjugate selected from a preferred group of Neutrokin- α conjugates as defined above, and one or more pharmaceutically acceptable excipients. A pharmaceutical composition that comprises Neutrokin- α conjugate may be formulated, as is well known in the art.

[0229] An additional embodiment of the present invention is a pharmaceutical composition comprising a Neutrokin- α complex and one or more pharmaceutically acceptable carriers. A preferred composition of the present invention is a pharmaceutical composition comprising a Neutrokin- α complex selected from a preferred group of Neutrokin- α complexes as defined above, and one or more pharmaceutically acceptable excipients. A pharmaceutical composition that comprises Neutrokin- α complex may be formulated, as is well known in the art.

[0230] In specific embodiments, a Neutrokin- α conjugate or Neutrokin- α complex is formulated in a solution comprising 10 mM sodium acetate and 140 mM sodium chloride, pH 6.0. In other specific embodiments, a Neutrokin- α conjugate or Neutrokin- α complex is formulated in a

- 94 -

solution comprising 10 mM sodium acetate and 140 mM sodium chloride, 3-5% ascorbate, pH 6.0. Optionally, the formulation may also comprise 1 mM of the chelator diethylene triamine-N,N,N',N'',N''-pentaacetic acid (DTPA).

[0231] The pharmaceutical composition of the invention can be administered to any animal that can experience the beneficial effects of the Neutrokin- α conjugate or the Neutrokin α complex of the invention. Foremost among such animals are humans, although the invention is not intended to be so limited.

[0232] The pharmaceutical composition of the present invention can be administered by any means that achieves its intended purpose. For example, administration can be by subcutaneous, intravenous, intramuscular, intraperitoneal, buccal, or ocular routes, rectally, parenterally, intrasystemically, intravaginally, topically (as by powders, ointments, drops or transdermal patch), or as an oral or nasal spray. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

[0233] In addition to the Neutrokin- α conjugate or Neutrokin- α complex, the new pharmaceutical preparation can contain one or more suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations that can be used pharmaceutically.

[0234] The pharmaceutical preparation of the present invention is manufactured in a manner that is, itself, known, for example, by means of conventional mixing, granulating, dragée-making, dissolving, or lyophilizing processes. Thus, a pharmaceutical preparation for oral use can be obtained by combining the Neutrokin- α complex or the Neutrokin- α conjugate with one or more solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

[0235] Pharmaceutical excipients are well known in the art. *See, e.g.*, Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., USA. Suitable excipients include fillers such as saccharides, for example, lactose or sucrose, mannitol or sorbitol; cellulose preparations, and calcium phosphates, for example, tricalcium phosphate or calcium hydrogen phosphate; as well as binders, such as, starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents can be added, such as the above- mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as, sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragée cores are provided with suitable coatings that, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions can be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol, and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations, such as, acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or pigments can be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

[0236] Other pharmaceutical preparations which can be used orally include push- fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as, glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules that may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as, fatty oils or liquid paraffin. In addition, stabilizers may be added.

- 96 -

[0237] A Neutrokin- α conjugate, Neutrokin- α complex, or Neutrokin- α composition according to the present invention may also be administered parenterally as an injectable dosage form in a physiologically acceptable diluent such as sterile liquids or mixtures thereof, including water, saline, aqueous dextrose and other pharmaceutically acceptable sugar solutions, alcohols such as ethanol, isopropanol, or hexadecyl alcohol, glycols such as propylene glycol and polyethylene glycol, glycerol ketals such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers such as poly(ethyleneglycol)400, a pharmaceutically acceptable oil, fatty acid, fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or detergent, suspending agent such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, an emulsifying agent or pharmaceutical adjuvants. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage. It is also preferable that the composition is preserved against the contaminating action of micro-organisms such as bacteria and fungi.

[0238] Additional, suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts, alkaline solutions, and cyclodextrin inclusion complexes. Especially preferred alkaline salts are ammonium salts prepared, for example, with Tris, choline hydroxide, Bis-Tris propane, *N*-methylglucamine, or arginine.

[0239] In an additional embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile, isotonic, aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in

- 97 -

unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0240] Pharmaceutically acceptable oils which are useful in the formulation herein include, for example, those of petroleum, animal, vegetable, or synthetic origin, including peanut oil, soybean oil, sesame oil, cottonseed oil, olive oil, sunflower oil, petrolatum, and mineral oil. Fatty acids which may be used include, for example, oleic acid, stearic acid, and isostearic acid, while the fatty acid esters useful herein include ethyl oleate and isopropyl myristate. Suitable soaps include, for example, fatty acid alkali metal, ammonium, and triethanolamine salts. Acceptable detergents include cationic detergents and anionic detergents. Suitable cationic detergents include, for example, dimethyl dialkyl ammonium halides, alkyl pyridinium halides, and alkylamine acetates. Suitable anionic detergents include, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether and monoglyceride sulfates, and sulfosuccinates. Useful non-ionic detergents include fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers. Amphoteric detergents include alkyl-beta-aminopropionates and 2-alkylimidazoline quaternary salts, and mixtures thereof.

[0241] The parenteral compositions of this invention preferably will contain from about 0.1 mg/mL to about 10 mg/mL of the Neutrokin- α conjugate or Neutrokin- α complex as described herein in solution. In specific embodiments, the concentration of Neutrokin- α conjugate or Neutrokin- α complex in the parenteral composition is approximately 0.2 mg/mL. In specific embodiments, the concentration of Neutrokin- α conjugate or Neutrokin- α complex in the parenteral composition is approximately 2.0

mg/mL. The parenteral formulations in the form of sterile injectable solutions or suspensions will also preferably contain from about 0.05% to about 5% suspending agent in an isotonic medium. Buffers and preservatives may be added. A suitable surfactant may also be added. These surfactants may include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate, and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

[0242] In addition, suspensions of the active compounds as appropriate oily injection suspensions can be administered. Suitable lipophilic solvents or vehicles include, for example, fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or polyethylene glycol-400. Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

[0243] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

[0244] Suspensions, in addition to the active compounds, may contain suspending agents such as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

- 99 -

- [0245] Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical administration, including those for inhalation, may be prepared as a dry powder which may be pressurized or non-pressurized. In nonpressurized powder compositions, the active ingredients in finely divided form may be used in admixture with a larger-sized pharmaceutically acceptable inert carrier comprising particles having a size, for example, of up to 100 micrometers in diameter. Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 micrometers.
- [0246] Alternatively, the composition may be pressurized and contain a compressed gas, such as nitrogen or a liquefied gas propellant. The liquefied propellant medium and indeed the total composition are preferably such that the active ingredients do not dissolve therein to any substantial extent. The pressurized composition may also contain a surface-active agent. The surface-active agent may be a liquid or solid non-ionic surface-active agent or may be a solid anionic surface-active agent. It is preferred to use the solid anionic surface-active agent in the form of a sodium salt.
- [0247] A further form of topical administration is to the eye. The compounds and compositions of the present invention are delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the compounds are maintained in contact with the ocular surface for a sufficient time period to allow the compounds to penetrate the corneal and internal regions of the eye, as for example the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an encapsulating material.
- [0248] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of the present invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room

- 100 -

temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the drugs.

[0249] Another embodiment of the present invention is a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex and a diagnostically suitable carrier.

[0250] One or more lyophilization aids may be used in the preparation a Neutrokin- α conjugate, a Neutrokin- α complex, or compositions thereof. Such lyophilization aids include but are not limited to mannitol, lactose, sorbitol, dextran, Ficoll, and polyvinylpyrrolidone(PVP).

[0251] One or more stabilization aids may be used in the preparation a Neutrokin- α conjugate, a Neutrokin- α complex, or compositions thereof. Such stabilization aids include but are not limited to ascorbic acid, cysteine, monothioglycerol, sodium bisulfite, sodium metabisulfite, gentisic acid, and inositol.

[0252] One or more solubilization aids may be used in the preparation a Neutrokin- α conjugate, a Neutrokin- α complex, or compositions thereof. Such solubilization aids include but are not limited to ethanol, glycerin, polyethylene glycol, propylene glycol, polysorbates and lecithin.

[0253] One or more bacteriostats may be used in the preparation a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex. Such bacteriostats include but are not limited to benzyl alcohol, benzalkonium chloride, chlorbutanol, and methyl, propyl or butyl paraben.

[0254] In specific embodiment, the composition of the present invention is formulated using the BEMATM BioErodible Mucoadhesive System, MCATM MucoCutaneous Absorption System, SMPTM Solvent MicroParticle System, or BCPTM BioCompatible Polymer System of Atrix Laboratories, Inc. (Fort Collins, Colorado).

[0255] Sustained-release compositions also include liposomally entrapped compositions of the invention (see generally, Langer, Science 249:1527-1533 (1990); Treat *et al.*, in "Liposomes in the Therapy of Infectious Disease and Cancer," Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and

353-365 (1989)). Liposomes comprising a Neutrokin- α complex or conjugate may be prepared by methods known per se: DE 3,218,121; Epstein *et al.*, *Proc. Natl. Acad. Sci. USA* 82:3688-3692 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal therapy.

[0256] In another embodiment, a sustained release composition of the invention includes crystal formulations known in the art.

[0257] In yet an additional embodiment, a composition of the invention is delivered by way of a pump (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald *et al.*, *Surgery* 88:507 (1980); Saudek *et al.*, *N. Engl. J. Med.* 321:574 (1989)).

[0258] Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

[0259] The composition comprising the Neutrokin- α conjugate or Neutrokin- α complex to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (*e.g.*, 0.2 micron membranes). A therapeutic Neutrokin- α conjugate or Neutrokin- α complex compositions generally is placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0260] Pharmaceutical compositions containing Neutrokin- α conjugate or Neutrokin- α complex of the invention may be administered orally, rectally, parenterally, subcutaneously, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray (*e.g.*, via inhalation of a vapor or powder). In one embodiment, "pharmaceutically acceptable carrier" means a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or

- 102 -

formulation auxiliary of any type. In a specific embodiment, "pharmaceutically acceptable" means approved by a regulatory agency of the federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly humans. Nonlimiting examples of suitable pharmaceutical carriers according to this embodiment are provided in "Remington's Pharmaceutical Sciences" by E.W. Martin, and include sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can be employed as liquid carriers, particularly for injectable solutions. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

[0261] The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[0262] In one embodiment, a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex is administered subcutaneously.

[0263] In another embodiment, a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex is administered intravenously.

[0264] A composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, *e.g.*, films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

[0265] Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. *et al.*, *Biopolymers* 22:547-556 (1983)), poly-(2-hydroxyethyl methacrylate) (R. Langer *et al.*, *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer *et al.*, *id.*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

[0266] In another embodiment, a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex is formulated in a biodegradable, polymeric drug delivery system, for example as described in U.S. Patent Nos. 4,938,763; 5,278,201; 5,278,202; 5,324,519; 5,340,849; and 5,487,897 and in International Publication Numbers WO01/35929, WO00/24374, and WO00/06117 which are hereby incorporated by reference in their entirety. In specific preferred embodiments, the composition comprising a Neutrokin- α conjugate or Neutrokin- α complexes formulated using the ATRIGEL® Biodegradable System of Atrix Laboratories, Inc. (Fort Collins, Colorado). In other specific embodiments, a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex is formulated using the ProLease® sustained release system available from Alkermes, Inc. (Cambridge, MA).

[0267] Examples of biodegradable polymers which can be used in the formulation of a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex, include but are not limited to, polylactides, polyglycolides, polycaprolactones, polyanhydrides, polyamides, polyurethanes, polyesteramides, polyorthoesters, polydioxanones, polyacetals, polyketals, polycarbonates, polyorthocarbonates, polyphosphazenes, polyhydroxybutyrates, polyhydroxyvalerates, polyalkylene oxalates, polyalkylene succinates, poly(malic acid), poly(amino acids), poly(methyl vinyl ether), poly(maleic anhydride), polyvinylpyrrolidone, polyethylene glycol, polyhydroxycellulose, chitin, chitosan, and copolymers, terpolymers, or combinations or mixtures of the above materials. The preferred polymers

- 104 -

are those that have a lower degree of crystallization and are more hydrophobic. These polymers and copolymers are more soluble in the biocompatible solvents than the highly crystalline polymers such as polyglycolide and chitin which also have a high degree of hydrogen-bonding. Preferred materials with the desired solubility parameters are the polylactides, polycaprolactones, and copolymers of these with glycolide in which there are more amorphous regions to enhance solubility. In specific preferred embodiments, the biodegradable polymers which can be used in the formulation of a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex are poly(lactide-co-glycolides). Polymer properties such as molecular weight, hydrophobicity, and lactide/glycolide ratio may be modified to obtain the desired Neutrokin- α conjugate or Neutrokin- α complex release profile (See, *e.g.*, Ravivarapu *et al.*, Journal of Pharmaceutical Sciences 89:732-741 (2000), which is hereby incorporated by reference in its entirety).

[0268] It is also preferred that the solvent for the biodegradable polymer be non-toxic, water miscible, and otherwise biocompatible. Examples of such solvents include, but are not limited to, N-methyl-2-pyrrolidone, 2-pyrrolidone, C₂ to C₆ alkanols, C₁ to C₁₅ alcohols, diols, triols, and tetraols such as ethanol, glycerine propylene glycol, butanol; C₃ to C₁₅ alkyl ketones such as acetone, diethyl ketone and methyl ethyl ketone; C₃ to C₁₅ esters such as methyl acetate, ethyl acetate, ethyl lactate; alkyl ketones such as methyl ethyl ketone, C₁ to C₁₅ amides such as dimethylformamide, dimethylacetamide and caprolactam; C₃ to C₂₀ ethers such as tetrahydrofuran, or solketal; tweens, triacetin, propylene carbonate, decylmethylsulfoxide, dimethyl sulfoxide, oleic acid, 1-dodecylazacycloheptan-2-one. Other preferred solvents are benzyl alcohol, benzyl benzoate, dipropylene glycol, tributyrin, ethyl oleate, glycerin, glycofural, isopropyl myristate, isopropyl palmitate, oleic acid, polyethylene glycol, propylene carbonate, and triethyl citrate. The most preferred solvents are N-methyl-2-pyrrolidone, 2-pyrrolidone, dimethyl

- 105 -

sulfoxide, triacetin, and propylene carbonate because of the solvating ability and their compatibility.

[0269] Additionally, formulations comprising a Neutrokin- α conjugate or a Neutrokin- α complex and a biodegradable polymer may also include release-rate modification agents and/or pore-forming agents. Examples of release-rate modification agents include, but are not limited to, fatty acids, triglycerides, other like hydrophobic compounds, organic solvents, plasticizing compounds and hydrophilic compounds. Suitable release rate modification agents include, for example, esters of mono-, di-, and tricarboxylic acids, such as 2-ethoxyethyl acetate, methyl acetate, ethyl acetate, diethyl phthalate, dimethyl phthalate, dibutyl phthalate, dimethyl adipate, dimethyl succinate, dimethyl oxalate, dimethyl citrate, triethyl citrate, acetyl tributyl citrate, acetyl triethyl citrate, glycerol triacetate, di-(*n*-butyl) sebecate, and the like; polyhydroxy alcohols, such as propylene glycol, polyethylene glycol, glycerin, sorbitol, and the like; fatty acids; triesters of glycerol, such as triglycerides, epoxidized soybean oil, and other epoxidized vegetable oils; sterols, such as cholesterol; alcohols, such as C6-C12 alkanols, 2-ethoxyethanol, and the like. The release rate modification agent may be used singly or in combination with other such agents. Suitable combinations of release rate modification agents include, but are not limited to, glycerin/propylene glycol, sorbitol/glycerine, ethylene oxide/propylene oxide, butylene glycol/adipic acid, and the like. Preferred release rate modification agents include, but are not limited to, dimethyl citrate, triethyl citrate, ethyl heptanoate, glycerin, and hexanediol. Suitable pore-forming agents that may be used in the polymer composition include, but are not limited to, sugars such as sucrose and dextrose, salts such as sodium chloride and sodium carbonate, polymers such as hydroxylpropylcellulose, carboxymethylcellulose, polyethylene glycol, and polyvinylpyrrolidone. Solid crystals that will provide a defined pore size, such as salt or sugar, are preferred.

[0270] For parenteral administration, in one embodiment, a Neutrokin- α conjugate or a Neutrokin- α complex is formulated generally by mixing it

- 106 -

at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, *i.e.*, one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

[0271] Generally, the formulations are prepared by contacting a Neutrokine-alpha conjugate or a Neutrokine-alpha complex uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

[0272] The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, *e.g.*, polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, sucrose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; preservatives, such as cresol, phenol, chlorobutanol, benzyl alcohol and parabens, and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

[0273] In a specific embodiment, a composition of the invention comprises, about between 0.1 mg/mL and 20mg/mL of a Neutrokine-alpha conjugate or a

- 107 -

Neutrokin- α complex wherein the Neutrokin- α protein of said conjugate or complex comprises amino acid residues 134-285 of SEQ ID NO:2, 10.0 mM sodium citrate, 140.0 mM sodium chloride, 8.7mM HEPES, 4% (w/v) sodium ascorbate, and 3.3% (w/v) Gentran-40. In a specific embodiment, a composition of the invention comprises, between 1mg/mL and 10mg/mL of a Neutrokin- α conjugate or a Neutrokin- α complex wherein the Neutrokin- α protein of said conjugate or complex comprises amino acid residues 134-285 of SEQ ID NO:2, 10.0 mM sodium citrate, 140.0 mM sodium chloride, 8.7mM HEPES, 4% (w/v) sodium ascorbate, and 3.3% (w/v) Gentran-40. In a specific embodiment, a composition of the invention comprises, between 2mg/mL and 8mg/mL of a Neutrokin- α conjugate or a Neutrokin- α complex wherein the Neutrokin- α protein of said conjugate or complex comprises amino acid residues 134-285 of SEQ ID NO:2, 10.0 mM sodium citrate, 140.0 mM sodium chloride, 8.7mM HEPES, 4% (w/v) sodium ascorbate, and 3.3% (w/v) Gentran-40. In a specific embodiment, a composition of the invention comprises, between 3mg/mL and 6mg/mL of a Neutrokin- α conjugate or a Neutrokin- α complex wherein the Neutrokin- α protein of said conjugate or complex comprises amino acid residues 134-285 of SEQ ID NO:2, 10.0 mM sodium citrate, 140.0 mM sodium chloride, 8.7mM HEPES, 4% (w/v) sodium ascorbate, and 3.3% (w/v) Gentran-40. The above described compositions may be used as pharmaceutical compositions.

[0274] A composition of the invention may be administered alone or in combination with other therapeutic agents, including but not limited to, chemotherapeutic agents, antibiotics, antivirals, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, immunosuppressants, and cytokines. Combinations may be administered either concomitantly, *e.g.*, as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but

- 108 -

simultaneously, *e.g.*, as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

[0275] In one embodiment, the compositions of the invention are administered in combination with anti-TNF-alpha antibodies such as infliximab (also known as Remicade™, Centocor, Inc.).

[0276] In one embodiment, the compositions of the invention are administered in combination with ENBREL™ (Etanercept).

[0277] In one embodiment, the compositions of the invention are administered in combination with anti-CD20 antibodies such as Ibritumomab Tiuxetan (Zevalin™) or rituximab (Rituxan™).

[0278] In one embodiment, the compositions of the invention are administered in combination with anti-TRAIL-R1 and/or TRAIL-R2 antibodies, for example those described in WO02/97033 and WO02/79377, which are hereby incorporated by reference in their entireties.

[0279] In one embodiment, a composition of the invention is administered in combination with one or more other members of the TNF family. TNF, TNF-related, or TNF-like molecules that may be administered with the compositions of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), AIM-II (International Publication No. WO 97/34911), APRIL (J. Exp. Med. 188(6):1185-1190), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and Neutrokine-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO

- 109 -

98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12.

[0280] In another embodiment, a composition of the invention is administered in combination with CD40 ligand (CD40L), a soluble form of CD40L (*e.g.*, AVREND™), biologically active fragments, variants, or derivatives of CD40L, anti-CD40L antibodies (*e.g.*, agonistic or antagonistic antibodies), and/or anti-CD40 antibodies (*e.g.*, agonistic or antagonistic antibodies).

Demonstration of Therapeutic or Prophylactic Activity

[0281] The Neutrokin- α conjugate, Neutrokin- α complex, and compositions thereof of the invention are preferably tested *in vitro*, and then *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, *in vitro* assays which can be used to determine whether administration of a specific compound is indicated include *in vitro* cell culture assays in which a patient tissue sample is grown in culture and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

[0282] Furthermore, a number of cell lines can be used to test for the binding of a Neutrokin- α conjugate or Neutrokin- α complex of the invention. Suitable cell lines that may be used to test this binding include, for example: IM-9 (ATCC CCL-159); Reh (ATCC CRL-8286); ARH-77 (ATCC CRL-1621); Raji (ATCC CCL-86); Namalwa (CRL-1432); RPMI 8226

- 110 -

(ATCC CCL-155). Additionally, one could use other cell lines (*e.g.*, CHO, NS0) transfected with expression constructs for receptors for Neutrokin-alpha. Suitable expression constructs include expression constructs which encode receptors for Neutrokin-alpha, *e.g.*, BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9). A Neutrokin-alpha conjugate or Neutrokin-alpha complex of the invention will bind to a receptor capable of binding a Neutrokin-alpha protein. Cells exposed to Neutrokin-alpha complexes of the invention may further be assessed for viability.

Diagnostic Uses of Neutrokin-Alpha Conjugates and Neutrokin-Alpha Complexes

[0283] Neutrokin-alpha receptors are expressed primarily on B cells (see beginning of Therapeutic Uses of Neutrokin-Alpha Conjugates and Neutrokin-Alpha Complexes Section below). Herein, Neutrokin-alpha receptors refer not only to Neutrokin-alpha receptors such as BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9, but also to allelic variants, conserved variants, synthetic fragments and/or biologically processed fragments of Neutrokin-alpha receptors that are capable of binding Neutrokin-alpha.

[0284] Accordingly, in one embodiment, a Neutrokin-alpha conjugate or Neutrokin-alpha complex of the invention is used to quantitate or qualitate concentrations of B lineage cells expressing Neutrokin-alpha receptor on their cell surfaces (*e.g.*, normal B cells as well as B cell related leukemias or lymphomas).

[0285] In one embodiment, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention may be used in diagnostic applications comprising detecting Neutrokin-alpha receptor expression (BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9)). Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention may also be used to determine the structure and/or temporal, tissue, cellular, or subcellular location of Neutrokin-alpha receptors and/or to determine the

- 111 -

activity of signalling pathways associated with Neutrokin-alpha. These diagnostic assays may be performed *in vitro*, such as, for example, on blood samples or biopsy tissue, or *in vivo*, using techniques described herein or otherwise known in the art.

[0286] In specific embodiments, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention may be used to diagnose cancers, particularly lymphocytic cancers. For example, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention may be used to detect cancer cells that express Neutrokin-alpha receptors. Diagnosis of cancer may be made on the basis of increased or decreased expression of Neutrokin-alpha receptors on cancer cells compared to non-cancerous Neutrokin-alpha receptor expressing cells (normal B cells). Alternatively, diagnosis of cancer may be made on the basis of combined expression of Neutrokin-alpha receptor expression, at normal or aberrant levels, in combination with expression of other markers that are diagnostic of cancer.

[0287] As a non-limiting example, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention, may be used to determine the presence or absence of Neutrokin-alpha receptors on cells from a patient that is a candidate for treatment with the Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention. In specific embodiments, the patients are cancer patients, particularly cancer patients with lymphocytic cancers, such as multiple myeloma and Non-Hodgkin's lymphoma. Expression of Neutrokin-alpha receptors on cancer cells from such cancer patients would indicate that therapy with Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention is likely to be effective, whereas the absence of detectable expression would suggest the patient probably would not benefit from therapy with Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention.

[0288] As another non-limiting example, Neutrokin-alpha conjugates and/or Neutrokin-alpha complexes of the invention may be used to localize and/or

- 112 -

quantitate Neutrokin- α receptor expressing cells, *e.g.*, multiple myeloma cells, non-Hodgkin's lymphoma, or chronic lymphocytic leukemia cells.

[0289] In one embodiment, a Neutrokin- α conjugate or Neutrokin- α complex of the invention is used to diagnose or monitor an individual having an immunodeficiency. Examples of immunodeficiencies that may be diagnosed or monitored with a Neutrokin- α conjugate or Neutrokin- α complex of the invention, include, but are not limited to, common variable immunodeficiency (CVID) and diseases characterized by deficiencies in one or more classes or subclasses of immunoglobulin (*e.g.*, selective IgA deficiency, ataxia telangiectasia, X-linked agammaglobulinemia, severe combined immunodeficiency (SCID), Wiskott-Aldrich syndrome, and hyper IgM syndrome)

[0290] Diagnosis of an immunodeficiency may be made on the basis of increased or decreased expression of Neutrokin- α receptors on known Neutrokin- α receptor expressing cells (*e.g.*, lymphocytes, particularly B cells) compared to Neutrokin- α receptor expressing cells from patients without an immunodeficiency. Immunodeficiencies may be characterized by decreased levels of Neutrokin- α receptor expression, suggesting the immunodeficiency may be the result of decreased or absent Neutrokin- α receptor activity. Alternatively, increased levels of Neutrokin- α receptor expression may also be indicative of an immunodeficiency if, for example, Neutrokin- α receptors are only partially functional and therefore, more receptors are necessary to deliver Neutrokin- α receptor-mediated signaling. Thus, diagnosis of an immunodeficiency may be made on the basis of increased or decreased expression of Neutrokin- α receptors compared to Neutrokin- α receptor expression in a patient without the immunodeficiency. Alternatively, diagnosis of an immunodeficiency may be made on the basis of expression of Neutrokin- α receptor expression, at normal or aberrant levels, in combination with expression of other markers that are diagnostic of an immunodeficiency (*e.g.* cell surface or serum immunoglobulin expression).

- 113 -

[0291] In another embodiment, a Neutrokinine-alpha conjugate or Neutrokinine-alpha complex of the invention is used to diagnose or monitor an individual having an autoimmune disease or disorder, particularly an autoimmune disease associated with the production of autoantibodies. Examples of autoimmune diseases that may be diagnosed or monitored with a Neutrokinine-alpha conjugate or Neutrokinine-alpha complex of the invention, include, but are not limited to, rheumatoid arthritis, systemic lupus erythematosus (SLE), and Sjögren's syndrome.

[0292] Diagnosis of an autoimmune disease may be made on the basis of increased or decreased expression of Neutrokinine-alpha receptors on known Neutrokinine-alpha receptor expressing cells (*e.g.*, lymphocytes, particularly B cells) compared to Neutrokinine-alpha receptor expressing cells from patients without an autoimmune disease. Autoimmune diseases may be characterized by increased levels of Neutrokinine-alpha receptor expression, suggesting the autoimmune disease may be the result of excess Neutrokinine-alpha receptor activity. Alternatively, decreased levels of Neutrokinine-alpha receptor expression may also be indicative of an autoimmune disease, if for example, Neutrokinine-alpha receptors are more easily activated to signal or even are constitutively active in the absence of ligand, and therefore, fewer receptors are necessary to deliver Neutrokinine-alpha receptor-mediated signaling. Thus, diagnosis of an autoimmune disease may be made on the basis of increased or decreased expression of Neutrokinine-alpha receptors compared to Neutrokinine-alpha receptor expression in a patient without the autoimmune disease. Alternatively, diagnosis of an autoimmune disease may be made on the basis of expression of Neutrokinine-alpha receptor expression, at normal or aberrant levels, in combination with expression of other markers that are diagnostic of an autoimmune disease (*e.g.*, cell surface or serum immunoglobulin expression).

[0293] Any means described herein or otherwise known in the art may be applied to detect Neutrokinine-alpha receptors (*e.g.*, FACS analysis or ELISA detection of Neutrokinine-alpha proteins of the invention. Additionally,

- 114 -

hybridization or PCR detection of Neutrokin- α receptor polynucleotides of the invention (BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and/or BCMA (SEQ ID NO:9) and to determine the expression profile of Neutrokin- α polynucleotides and/or polypeptides of the invention in a biological sample.

[0294] By analyzing or determining the expression level of the gene encoding the Neutrokin- α receptor is intended qualitatively or quantitatively measuring or estimating the level of the Neutrokin- α receptor polypeptide in a first biological sample either directly (*e.g.*, by determining or estimating absolute protein level) or relatively (*e.g.*, by comparing to the Neutrokin- α receptor present in a second biological sample). In one embodiment, the Neutrokin- α receptor level in the first biological sample is measured or estimated and compared to a standard Neutrokin- α receptor level, the standard being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder of the immune system. As will be appreciated in the art, once a standard Neutrokin- α receptor level is known, it can be used repeatedly as a standard for comparison.

[0295] By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source. As indicated, biological samples include body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid), immune system tissue, and other tissue sources. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art.

[0296] The Neutrokin- α conjugate or complex of the present invention may, additionally, be employed histologically, as in immunofluorescence, immunoelectron microscopy, or non-immunological assays, for *in situ* detection of Neutrokin- α receptors. *In situ* detection may be accomplished by removing a histological specimen from a patient, and applying thereto a labeled Neutrokin- α conjugate or Neutrokin- α

- 115 -

complex of the present invention. The Neutrokin- α conjugate or Neutrokin- α complex is preferably applied by overlaying the Neutrokin- α conjugate or Neutrokin- α complex onto a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of Neutrokin- α receptors, but also their distribution in the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such *in situ* detection.

[0297] Immunoassays and non-immunoassays for Neutrokin- α receptor products or conserved variants or peptide fragments thereof will typically comprise incubating a sample, such as a biological fluid, a tissue extract, freshly harvested cells, or lysates of cells which have been incubated in cell culture, in the presence of Neutrokin- α conjugate or Neutrokin- α complex capable of identifying Neutrokin- α receptors, and detecting the bound Neutrokin- α conjugate or Neutrokin- α complex by any of a number of techniques well-known in the art.

[0298] The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier such as nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the Neutrokin- α conjugate or Neutrokin- α complex. The solid phase support may then be washed with the buffer a second time to remove unbound Neutrokin- α conjugate or Neutrokin- α complex. Optionally, the Neutrokin- α conjugate or Neutrokin- α complex is subsequently labeled. The amount of bound label on solid support may then be detected by conventional means.

[0299] By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for

- 116 -

the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

[0300] The binding activity of a given lot Neutrokin- α conjugate or Neutrokin- α complex may be determined according to well-known methods. Those skilled in the art will be able to determine operative and optimal assay conditions for each determination by employing routine experimentation. In particular, cell lines as described above may be used.

[0301] In addition to assaying Neutrokin- α receptor levels in a biological sample obtained from an individual, Neutrokin- α receptors can also be detected *in vivo* by imaging. For example, in one embodiment of the invention, a Neutrokin- α conjugate or Neutrokin- α complex is used to image B cell lymphomas. In another embodiment, Neutrokin- α conjugate or Neutrokin- α complex is used to image lymphomas (*e.g.*, monocyte and B cell lymphomas).

[0302] Labels or markers for *in vivo* imaging of Neutrokin- α conjugate or Neutrokin- α complex include those detectable by X-radiography, NMR, MRI, CAT-scans or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject.

[0303] *In vivo* tumor imaging is described in S.W. Burchiel *et al.*, "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

- 117 -

[0304] Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the Neutrokin- α conjugate or Neutrokin- α complex, it is possible to detect Neutrokin- α receptors through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., *Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques*, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by means including, but not limited to, a gamma counter, a scintillation counter, or autoradiography.

[0305] The Neutrokin- α conjugate or Neutrokin- α complex can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series.

Therapeutic Uses of Neutrokin- α Conjugates and Neutrokin- α Complexes

[0306] Neutrokin- α receptor expression, defined functionally by the binding of biotinylated Neutrokin- α to cells, is found predominantly on B cells (see, *e.g.*, Moore *et al.*, *Science* 285:260-263 (1999), which is hereby incorporated by reference in its entirety), although TACI has also been reported to be expressed on activated T cells (Wang *et al.*, *Nature Immunol.* 2:577-8 (2001), which is hereby incorporated by reference in its entirety.) Furthermore, Neutrokin- α receptor expression is not observed in pre-B cells, rather Neutrokin- α receptor expression becomes observable at the same stage in B cell development when surface Ig expression becomes apparent (see, *e.g.*, Hsu *et al.*, *J. Immunol.* 168:5993-6 (2002), which is hereby incorporated by reference in its entirety).

[0307] Receptors which bind proteins comprising Neutrokin- α or fragments or variants thereof may also be expressed on non-hematopoietic cells. In specific embodiments, receptors which bind proteins comprising Neutrokin- α or fragments or variants thereof (*e.g.*, Neutrokin- α heterotimers (described below) comprising one or two APRIL monomers) are

- 118 -

expressed on cells or cell lines of fibroblastic or epithelial lineage or having fibroblastic or epithelial morphology (*e.g.*, NIH-3T3 fibroblasts, A549 lung carcinoma cells or HT-29 colorectal adenocarcinoma cells).

[0308] Additionally, Neutrokin- α receptor expression, defined functionally by the binding of biotinylated Neutrokin- α to cells, has been observed on multiple myeloma, Non-Hodgkin's lymphoma, and chronic lymphocytic leukemia primary tumor explants (see, *e.g.*, Briones *et al.*, *Experimental Hematology* 30:135-141 (2002) and Novak *et al.*, *Blood* 100:2973-2979 (2002), each of which is hereby incorporated by reference in its entirety) and on B lineage immortalized hematopoietic cell lines (*e.g.*, IM-9 (ATCC CCL-159); Reh (ATCC CRL-8286); ARH-77 (ATCC CRL-1621); Raji (ATCC-CCL-86); Namalwa (CRL-1432); RPMI 8226 (ATCC CCL-155)).

[0309] The restricted expression profile of Neutrokin- α receptor expression described above makes Neutrokin- α an attractive vehicle for targeting therapies to lymphocytes, and to B lineage cells in particular.

[0310] Biodistribution studies of a Neutrokin- α complex injected into BALB/c mice illustrate that a Neutrokin- α conjugate or a Neutrokin- α complex has high *in vivo* targeting specificity for lymphoid tissues such as spleen and lymph nodes (See Example 1). Thus in a specific embodiment, the invention provides a method for the specific destruction or disablement of lymphoid tissue (*e.g.*, lymph nodes and spleen) comprising administering a Neutrokin- α conjugate or a Neutrokin- α complex. In a preferred embodiment, the lymphoid tissue is not permanently destroyed but rather is temporarily disabled, (*e.g.*, cells of hematopoietic lineage in lymphoid tissues are destroyed/killed while a Neutrokin- α conjugate or a Neutrokin- α complex is administered, but these populations recover once administration of the Neutrokin- α conjugate or the Neutrokin- α complex is stopped.)

[0311] The Neutrokin- α conjugate, Neutrokin- α complex, or composition of the present invention is useful as a therapeutic agent for the treatment of diseases and conditions associated with cells that express

- 119 -

Neutrokin- α receptors. Because Neutrokin- α receptors are known to be expressed predominantly on B cells, it is a preferred embodiment of the present invention that Neutrokin- α conjugates or complexes are useful for treating diseases involving B cells or B cell activity.

[0312] TACI is also known to be expressed by T cells, specifically activated T cells. Therefore, in other embodiments, Neutrokin- α conjugates or complexes may also be useful for treating diseases involving T cells or T cell activity.

[0313] The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a Neutrokin- α protein and/or a receptor for the Neutrokin- α (*e.g.*, BAFF-R, TACI, BCMA,) includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. The conjugate or complex of the invention may also be used to target and kill cells which bind Neutrokin- α .

[0314] In particular, the Neutrokin- α conjugate or Neutrokin- α complex are administered as a form of radiotherapy. In one embodiment, the method of the present invention comprises administering to a subject a composition comprising a Neutrokin- α complex and a pharmaceutically acceptable carrier, for radiotherapeutic treatment of said subject wherein the Neutrokin- α complex is administered in an amount sufficient for the needed treatment. In another embodiment, the method further comprises monitoring said patient. In another embodiment, the subject is a mammal. In a further embodiment, the subject is a human.

[0315] In another embodiment, the method of the present invention comprises administering to a subject a composition comprising a Neutrokin- α conjugate and a pharmaceutically acceptable carrier, for therapeutic treatment of said subject wherein the Neutrokin- α conjugate is administered in an amount sufficient for the needed treatment. In another embodiment, the method further comprises monitoring said patient. In another embodiment, the subject is a mammal. In a further embodiment, the subject is a human.

- 120 -

[0316] In one embodiment, the invention provides a method of target compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex to Neutrokin- α receptor expressing cells, such as, for example, B cells or T cells. A Neutrokin- α conjugate or a Neutrokin- α complex of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

[0317] In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering a Neutrokin- α conjugate or a Neutrokin- α complex that is associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein or cytotoxin to or into the targeted cell.

[0318] In another embodiment, the invention provides for a method of killing lymphocytes, comprising, or alternatively consisting of, contacting a Neutrokin- α conjugate or a Neutrokin- α complex with lymphocytes. In a specific embodiment, the method of killing lymphocytes, comprises, or alternatively consists of, administering to an animal in which such killing is desired, a Neutrokin- α conjugate or a Neutrokin- α complex in an amount effective to kill lymphocytes. Lymphocytes include, but are not limited to, healthy and diseased cells as found present in an animal, preferably a mammal and most preferably a human, or as isolated from an animal, transformed cells, cell lines derived from the above listed cell types, and cell cultures derived from the above listed cell types. Lymphocytes may be found or isolated in, for example, different developmental stages or in resting, activated or anergic states. In preferred embodiments, the lymphocytes are B lineage cells.

[0319] In another embodiment, the invention provides a method for the specific destruction (*i.e.*, killing) of cells (*e.g.*, the destruction of cancer cells with lymphocyte phenotypes) by administering a Neutrokin- α conjugate or a Neutrokin- α complex in which such destruction of cells is desired.

- 121 -

In one embodiment, the cells targeted for destruction express Neutrokin-alpha receptors on their surface. In another embodiment, the cells targeted for destruction are in proximity to cells that express Neutrokin-alpha receptors on their surface. In preferred embodiments the cells are B lineage tumors are B lineage cells.

[0320] In another embodiment, the invention provides a method for the specific destruction of lymphocytes (*e.g.*, the destruction of tumor cells) by administering a Neutrokin-alpha conjugate or a Neutrokin-alpha complex in association with toxins or cytotoxic prodrugs.

[0321] In a specific embodiment, the invention provides a method for the specific destruction of cells of B cell lineage (*e.g.*, B cell related leukemias or lymphomas) by administering a Neutrokin-alpha conjugate or a Neutrokin-alpha complex and a toxin or cytotoxic prodrug.

[0322] In one embodiment, the invention provides methods and compositions for inhibiting or reducing proliferation of lymphocytes, comprising, or alternatively consisting of, contacting an effective amount of a Neutrokin-alpha conjugate or a Neutrokin-alpha complex with lymphocytes, wherein the effective amount of a Neutrokin-alpha conjugate or a Neutrokin-alpha complex inhibits or reduces proliferation of lymphocytes. In another embodiment, the invention provides methods and compositions for inhibiting or reducing proliferation of lymphocytes comprising, or alternatively consisting of, administering to an animal in which such inhibition or reduction is desired, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex in an amount effective to inhibit or reduce B cell proliferation. In preferred embodiments, the lymphocytes are B cells.

[0323] B cell proliferation is most commonly assayed in the art by measuring tritiated-thymidine incorporation. This and other assays are commonly known in the art and could be routinely adapted for the use of determining the effect of the Neutrokin-alpha conjugate or the Neutrokin-alpha complex on B cell proliferation.

- 122 -

[0324] In one embodiment, the invention provides methods and compositions for decreasing lifespan of lymphocytes, comprising, or alternatively consisting of, contacting an effective amount of a Neutrokin- α conjugate or a Neutrokin- α complex with lymphocytes, wherein the effective amount of the Neutrokin- α conjugate or the Neutrokin- α complex inhibits or reduces lifespan of lymphocytes. In another embodiment, the lymphocytes are B cells.

[0325] B cell life span may be measured using or routinely modify techniques known in the art. In one example, B cell lifespan is measured *in vivo* may be measured by 5-bromo-2'-deoxyuridine (BrdU) labeling experiments which are well known to one skilled in the art. BrdU is a thymidine analogue that gets incorporated into the DNA of dividing cells. Cells containing BrdU in their DNA can be detected using, for example fluorescently labeled anti-BrdU antibody and flow cytometry. Briefly, an animal is injected with BrdU in an amount sufficient to label developing B cells. Then, a sample of B cells is withdrawn from the animal, for example, from peripheral blood, and analyzed for the percentage of cells that contain BrdU. Such an analysis performed at several time points can be used to calculate the half life of B cells. Alternatively, B cell survival may be measured *in vitro*. For example B cells may be cultured under conditions where proliferation does not occur, (for example the media should contain no reagents that crosslink the immunoglobulin receptor, such as anti-IgM antibodies) for a period of time (usually 2-4 days). At the end of this time, the percent of surviving cells is determined, using for instance, the vital dye Trypan Blue, or by staining cells with propidium iodide or any other agent designed to specifically stain apoptotic cells and analyzing the percentage of cells stained using flow cytometry. One could perform this experiment under several conditions, such as B cells treated with Neutrokin- α conjugates or complexes and untreated B cells or B cells treated with an unlabelled form of Neutrokin- α protein (*e.g.* a Neutrokin- α trimer consisting of three subunits each comprising amino acids 134-285 of SEQ ID NO:2) in order to determine the

- 123 -

effects of Neutrokin- α proteins on B cell survival. These and other methods for determining B cell lifespan are commonly known in the art.

Treatment of Cancer

[0326] The present invention provides methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex useful in the treatment of cancer, *e.g.* by destroying the cancerous cells and/or by inhibiting the growth, progression, and/or metastasis of cancer cells. In a preferred embodiment, the present invention provides methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex useful in the treatment of lymphoma. Use of Neutrokin- α conjugates and Neutrokin- α complexes is superior to conventional radiotherapy because Neutrokin- α radiotherapy is targeted at Neutrokin- α receptor expressing cells rather than a non-specific population of cells such as rapidly dividing cells. Furthermore, the fact that Neutrokin- α receptors are not expressed on the earlier stages of lymphocyte development (pre-lymphocytes such as pre B-cells) suggests that patients treated with Neutrokin- α conjugates and/or Neutrokin- α complexes will be able to reconstitute the cells of their immune system faster than a patient that had undergone conventional radiotherapy.

[0327] In a preferred embodiment, the present invention provides methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex useful in the treatment of non-Hodgkin's lymphoma. In particular embodiments, the non-Hodgkin's lymphoma may be a diffuse large cell, mantle cell, marginal zone, or follicular lymphoma. In preferred embodiments, Neutrokin α complexes of the invention comprising radiometal ions that emit beta-particles, *e.g.*, ^{90}Y , are used to treat solid tumors such as lymphoma.

[0328] In a preferred embodiment, the present invention provides methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex useful in the treatment of chronic lymphocytic leukemia.

- 124 -

- [0329] In a preferred embodiment, the present invention provides methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex useful in the treatment of multiple myeloma.
- [0330] In other preferred embodiments, Neutrokin α complexes of the invention comprising radiometal ions that emit α -particles or auger electrons, *e.g.*, ^{111}In (auger electron emitter), are used to treat diffuse cancers such as chronic lymphocytic leukemia and multiple myeloma.
- [0331] A non-limiting example of a Neutrokin- α conjugate or a Neutrokin- α complex of the invention that can be administered to a cancer patient is a Neutrokin- α conjugate or complex of Formula *IV* which binds to a Neutrokin- α receptor.
- [0332] Additional cancers that may be treated with methods and compositions comprising a Neutrokin- α conjugate or a Neutrokin- α complex include B cell malignancies such as acute lymphocytic leukemia (ALL), plasmacytomas, Burkitt's lymphoma, and EBV-transformed diseases.
- [0333] Neutrokin- α conjugates and/or a Neutrokin- α complexes may be useful in the treatment of Chronic Myelogenous Leukemia by decreasing the involvement of B cells and Ig associated with this disease.
- [0334] Because Neutrokin- α protein, *e.g.*, amino acids 134-285 of SEQ ID NO:2, is able to stimulate B cell proliferation, activation and survival, administration of Neutrokin- α conjugates may induce proliferation of normal and cancerous B cells, thus making it more susceptible to Neutrokin- α complexes and/or other anti-neoplastic agents (*e.g.*, chemotherapy and radiation therapy). For example, multiple myeloma is a slowly dividing disease and is thus refractory to virtually all anti-neoplastic regimens. If these cells were induced to proliferate more rapidly their susceptibility profile would likely change.
- [0335] The possibility that receptors for proteins comprising Neutrokin- α or fragments or variants thereof (*e.g.*, Neutrokin- α heterotimers (described below) comprising one or two APRIL monomers) are expressed on cells or cell lines of fibroblastic or epithelial lineage or having fibroblastic or

- 125 -

epithelial morphology, indicates that Neutrokin- α conjugates and/or Neutrokin- α complexes of the invention may be used to diagnose or treat cancers of cells of fibroblastic or epithelial lineage or having fibroblastic or epithelial morphology (*e.g.*, skin cancer, lung cancer, or colorectal cancer).

[0336] Additionally, because the energy from certain radioactive decay events can span more than a single cell diameter, Neutrokin- α conjugates and/or Neutrokin- α complexes of the invention may be used to kill cells, *e.g.*, cancerous cells, in close proximity to cells that express Neutrokin α receptors. In specific embodiments, Neutrokin- α conjugates and/or Neutrokin- α complexes of the invention may be used to prevent the metastasis of cancers through the lymphatic system. Cancerous cells that spread through the lymphatic system are likely to be in close proximity to cells that express receptors for Neutrokin- α (*e.g.*, B cells that are numerous in the lymph, lymph nodes and spleen). Thus, decay events from a Neutrokin- α complex of the invention associated with a B cell may be able to kill the cell expressing Neutrokin- α receptors, but also the cancer cell in close proximity to said cell expressing Neutrokin- α receptors, thereby preventing or inhibiting metastasis of said cancer through the lymphatic system.

[0337] The present invention further encompasses methods and compositions for killing cells bearing Neutrokin- α receptors and/or cells in close proximity to cells bearing Neutrokin- α receptors, comprising, or alternatively consisting of, contacting a Neutrokin- α conjugate and/or a Neutrokin- α complex of the invention with cells bearing Neutrokin- α receptors. In preferred embodiments, the cells bearing Neutrokin- α receptors are B cells.

[0338] The present invention further encompasses methods and compositions for killing cells bearing Neutrokin- α receptors or cells in close proximity to cells bearing a Neutrokin- α receptor, comprising, or alternatively consisting of, administering to an animal in which such killing is desired, a Neutrokin- α conjugate and/or a Neutrokin- α complex in an amount

- 126 -

effective to kill cells bearing Neutrokin- α receptors and/or cells in close proximity to cells bearing a Neutrokin- α receptor. In preferred embodiments, the cells bearing Neutrokin- α receptors are B cells.

Treatment of Autoimmune disease

- [0339] Elevated levels of Neutrokin- α protein correlate with autoimmune disease (Zhang *et al.*, *The Journal of Immunology* 166:6-10 (2001); Cheema *et al.*, *Arthritis and Rheumatism* 44:1313-1319 (2001), Groom *et al.*, *Journal of Clinical Investigation* 109:59-68 (2002); and Vaux, *The Journal of Clinical Investigation* 109:17-18 (2002)). Additionally, it has been demonstrated that that elevated levels of Neutrokin- α are relevant to the pathology of autoimmune disease because a Neutrokin- α antagonist (TACI-Fc) is able to ameliorate the symptoms of autoimmunity in mouse models of autoimmune disease (Gross *et al.*, *Nature*, 404:995-999 (2000)).
- [0340] Elevated levels of soluble Neutrokin- α have been observed in the serum of patients with Systemic Lupus Erythematosus (SLE). In comparing the sera of 150 SLE patients with that of 38 control individuals, it was found that most of the SLE patients had more than 5 ng/mL of serum Neutrokin- α , more than 30% of SLE patients had levels greater than 10 ng/mL, and approximately 10% of SLE patients had serum Neutrokin- α levels greater than 20 ng/mL. In contrast, the majority of normal controls had Neutrokin- α levels less than 5 ng/mL, and less than 10% had levels higher than 10 ng/mL. The elevated levels of Neutrokin- α protein in sera is present in the soluble form and has biologic activity as assayed by the ability to stimulate anti-IgM treated B cells *in vitro*. SLE patients with more than 15 ng/mL serum Neutrokin- α were also found to have elevated levels of anti-dsDNA antibodies compared to both normal controls and SLE patients with less than 5 ng/mL of serum Neutrokin- α (Zhang *et al.*, *The Journal of Immunology* 166:6-10 (2001)).

- 127 -

[0341] In addition, the serum of two subgroups of patients which were positive for anti-nuclear antibodies (ANA+) but did not meet the formal requirements of the American College of Rheumatology (ACR) for classification of SLE were analyzed for Neutrokin- α levels. The first subgroup of sera was ANA+ sera that came from patients who did not present with the clinical impression of SLE. This group had only slightly elevated levels of Neutrokin- α (~9 ng/mL Neutrokin- α). The second subgroup however, which was ANA+ sera from patients who presented with the clinical impression of SLE, had significantly increased Neutrokin- α levels (~15 ng/mL) (See, Zhang *et al.*, *The Journal of Immunology*, 166:6-10) (2001). These results suggest that an elevated level of Neutrokin- α precedes the formal fulfillment of the ACR criteria. The ACR criteria are described in Tan, E.M., *et al.*, *Arthritis and Rheumatism* 25:1271 - 1277 (1982).

[0342] Therefore, in one embodiment of the present invention, a Neutrokin- α conjugate or a Neutrokin- α complex may be used to treat or ameliorate autoimmune disease. In a specific embodiment a Neutrokin- α conjugate or a Neutrokin- α complex may be used to treat or ameliorate autoimmune diseases associated with autoantibody production.

[0343] The present invention further encompasses methods and compositions for treating or ameliorating an autoimmune disease, comprising, or alternatively consisting of, contacting a Neutrokin- α conjugate and/or a Neutrokin- α complex of the invention with cells bearing Neutrokin- α receptors. In preferred embodiments, the cells bearing Neutrokin- α receptors are B cells.

[0344] The present invention further encompasses methods and compositions for treating or ameliorating an autoimmune disease, comprising, or alternatively consisting of, administering to an animal in which such treatment is desired, a Neutrokin- α conjugate and/or a Neutrokin- α complex in an amount effective to treat or ameliorate an autoimmune disease.

- 128 -

[0345] Autoantibody production is common to several autoimmune diseases and contributes to tissue destruction and exacerbation of disease. Autoantibodies can also lead to the occurrence of immune complex deposition complications and lead to many symptoms of systemic lupus erythematosus, including kidney failure, neuralgic symptoms and death. Decreasing or preventing antibody production would be beneficial in the treatment of autoimmune diseases such as myasthenia gravis and rheumatoid arthritis. For example, B cells have been shown to play a role in the secretion of arthritogenic immunoglobulins in rheumatoid arthritis, (Korganow *et al.*, Immunity 10:451-61, 1999). One way to achieve the inhibition or abolition of autoantibody production, as well as of antibody production in general, is via the destruction or ablation of antibody secreting cells using a Neutrokin-alpha conjugate or a Neutrokin-alpha complex of the present invention.

[0346] In another preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate systemic lupus erythematosus.

[0347] In a preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate rheumatoid arthritis.

[0348] In another preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate Sjögren's syndrome.

[0349] In another preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate idiopathic thrombocytopenic purpura (ITP).

[0350] In another preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate IgA nephropathy.

[0351] In another preferred embodiment, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex may be used to treat or ameliorate Myasthenia gravis.

- 129 -

[0352] In another preferred embodiment, a Neutrokin- α conjugate or a Neutrokin- α complex may be used to treat or ameliorate vasculitis.

[0353] A non-limiting example of a Neutrokin- α conjugate or a Neutrokin- α complex of the invention that can be administered to an individual with an autoimmune disease is a Neutrokin- α conjugate or complex of Formula *IV* which binds to a Neutrokin- α receptor.

[0354] Neutrokin- α conjugates and/or a Neutrokin- α complexes may be useful in the treatment of chronic hypergammaglobulinemia evident in such diseases as monoclonal gammopathy of undetermined significance (MGUS), Waldenstrom's disease, related idiopathic monoclonal gammopathies, and plasmacytomas.

[0355] More generally, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.*, IgM, IgG, and/or IgA production), comprising, or alternatively consisting of, contacting an effective amount of Neutrokin- α complex or Neutrokin- α conjugate with lymphocytes, wherein the effective amount of Neutrokin- α complex or Neutrokin- α conjugate inhibits or reduces immunoglobulin production.

[0356] In specific embodiments, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.*, IgM, IgG, and/or IgA production) in response to T cell dependent antigens, comprising, or alternatively consisting of, contacting an effective amount of Neutrokin- α complex or Neutrokin- α conjugate with lymphocytes, wherein the effective amount of Neutrokin- α complex or Neutrokin- α conjugate inhibits or reduces immunoglobulin production in response to T cell dependent antigens.

[0357] In specific embodiments, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.* IgM, IgG, and/or IgA production) in response to T cell independent antigens, comprising, or alternatively consisting of, contacting an effective amount of Neutrokin- α complex or Neutrokin- α conjugate with lymphocytes, wherein the effective amount of Neutrokin- α complex or Neutrokin-

- 130 -

alpha conjugate inhibits or reduces immunoglobulin production in response to T cell independent antigens.

[0358] In another embodiment, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.*, IgM, IgG, and/or IgA production), comprising, or alternatively consisting of, administering to an animal in which such inhibition or reduction is desired, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex in an amount effective to inhibit or reduce immunoglobulin production.

[0359] In another embodiment, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.* IgM, IgG, and/or IgA production) in response to T cell dependent antigens, comprising, or alternatively consisting of, administering to an animal in which such inhibition or reduction is desired, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex in an amount effective to inhibit or reduce immunoglobulin production in response to T cell dependent antigens.

[0360] In another embodiment, the invention provides methods and compositions for inhibiting or reducing immunoglobulin production (*e.g.* IgM, IgG, and/or IgA production) in response to T cell independent antigens, comprising, or alternatively consisting of, administering to an animal in which such inhibition or reduction is desired, a Neutrokin-alpha conjugate or a Neutrokin-alpha complex in an amount effective to inhibit or reduce immunoglobulin production in response to T cell independent antigens.

[0361] Determinations of immunoglobulin levels are most often performed by comparing the level of immunoglobulin in a sample to a standard containing a known amount of immunoglobulin using ELISA assays. Determination of immunoglobulin levels in a given sample, can readily be determined using ELISA or other methods known in the art.

[0362] Additional autoimmune disorders and conditions associated with these disorders that may be treated, prevented, and/or diagnosed with the a Neutrokin-alpha conjugate or a Neutrokin-alpha complex include, but are not limited to, autoimmune hemolytic anemia, autoimmune neutropenia,

- 131 -

autoimmune neonatal thrombocytopenia, idiopathic thrombocytopenia purpura, autoimmunocytopenia, hemolytic anemia, antiphospholipid syndrome, dermatitis, allergic encephalomyelitis, myocarditis, relapsing polychondritis, rheumatic heart disease, glomerulonephritis (*e.g.*, IgA nephropathy), dense deposit disease, Multiple Sclerosis, Neuritis, Uveitis Ophthalmia, Polyendocrinopathies, Purpura (*e.g.*, Henloch-Scoenlein purpura), Reiter's Disease, Stiff-Man Syndrome, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, gluten sensitive enetropathy, insulin dependent diabetes mellitus, discoid lupus, and autoimmune inflammatory eye disease.

[0363] Additional autoimmune disorders that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, autoimmune thyroiditis, hypothyroidism (*i.e.*, Hashimoto's thyroiditis) (often characterized, *e.g.*, by cell-mediated and humoral thyroid cytotoxicity), systemic lupus erythematosus (often characterized, *e.g.*, by circulating and locally generated immune complexes), Goodpasture's syndrome (often characterized, *e.g.*, by anti-basement membrane antibodies), Pemphigus (often characterized, *e.g.*, by epidermal acantholytic antibodies), Receptor autoimmunities such as, for example, (a) Graves' Disease (often characterized, *e.g.*, by TSH receptor antibodies), (b) Myasthenia Gravis (often characterized, *e.g.*, by acetylcholine receptor antibodies), and (c) insulin resistance (often characterized, *e.g.*, by insulin receptor antibodies), autoimmune hemolytic anemia (often characterized, *e.g.*, by phagocytosis of antibody-sensitized RBCs), autoimmune thrombocytopenic purpura (often characterized, *e.g.*, by phagocytosis of antibody-sensitized platelets).

[0364] Additional autoimmune disorders that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, rheumatoid arthritis (often characterized, *e.g.*, by immune complexes in joints), schleroderma with anti-collagen antibodies (often characterized, *e.g.*, by nucleolar and other nuclear antibodies), mixed connective tissue disease (often characterized, *e.g.*, by antibodies to

- 132 -

extractable nuclear antigens (*e.g.*, ribonucleoprotein)), polymyositis/dermatomyositis (often characterized, *e.g.*, by nonhistone ANA), pernicious anemia (often characterized, *e.g.*, by antiparietal cell, microsomes, and intrinsic factor antibodies), idiopathic Addison's disease (often characterized, *e.g.*, by humoral and cell-mediated adrenal cytotoxicity, infertility (often characterized, *e.g.*, by antispermatozoal antibodies), glomerulonephritis (often characterized, *e.g.*, by glomerular basement membrane antibodies or immune complexes) such as primary glomerulonephritis and IgA nephropathy, bullous pemphigoid (often characterized, *e.g.*, by IgG and complement in basement membrane), Sjogren's syndrome (often characterized, *e.g.*, by multiple tissue antibodies, and/or a specific nonhistone ANA (SS-B)), diabetes mellitus (often characterized, *e.g.*, by cell-mediated and humoral islet cell antibodies), and adrenergic drug resistance (including adrenergic drug resistance with asthma or cystic fibrosis) (often characterized, *e.g.*, by beta-adrenergic receptor antibodies).

[0365] Additional autoimmune disorders that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, chronic active hepatitis (often characterized, *e.g.*, by smooth muscle antibodies), primary biliary cirrhosis (often characterized, *e.g.*, by mitochondrial antibodies), other endocrine gland failure (often characterized, *e.g.*, by specific tissue antibodies in some cases), vitiligo (often characterized, *e.g.*, by melanocyte antibodies), vasculitis (often characterized, *e.g.*, by Ig and complement in vessel walls and/or low serum complement), post-MI (often characterized, *e.g.*, by myocardial antibodies), cardiomy syndrome (often characterized, *e.g.*, by myocardial antibodies), urticaria (often characterized, *e.g.*, by IgG and IgM antibodies to IgE), atopic dermatitis (often characterized, *e.g.*, by IgG and IgM antibodies to IgE), asthma (often characterized, *e.g.*, by IgG and IgM antibodies to IgE), inflammatory myopathies, and many other inflammatory, granulomatous, degenerative, and atrophic disorders.

[0366] In another embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated,

- 133 -

prevented, and/or diagnosed using a Neutrokin- α conjugate or a Neutrokin- α complex of the invention.

Additional uses

[0367] The present invention further encompasses methods and compositions for treating or diseases and disorders, particularly those described herein, comprising, or alternatively consisting of, contacting a Neutrokin- α conjugate and/or a Neutrokin- α complex of the invention with cells bearing Neutrokin- α receptors. In preferred embodiments, the cells bearing Neutrokin- α receptors are B cells.

[0368] The present invention further encompasses methods and compositions for treating or ameliorating diseases and disorders, particularly those described herein, comprising, or alternatively consisting of, administering to an animal in which such treatment is desired, a Neutrokin- α conjugate and/or a Neutrokin- α complex in an amount effective to treat or ameliorate an autoimmune disease.

[0369] Neutrokin- α conjugates and/or Neutrokin- α complexes may also be useful for the treatment of atherosclerosis.

[0370] Neutrokin- α conjugates and/or Neutrokin- α complexes may also be useful for the treatment of asthma and other chronic airway diseases such as bronchitis and emphysema.

[0371] A Neutrokin- α conjugate or a Neutrokin- α complex may be used to modulate IgE concentrations *in vitro* or *in vivo*.

[0372] Additionally, a Neutrokin- α conjugate or a Neutrokin- α complex may be used to treat, prevent, and/or diagnose IgE-mediated allergic reactions. Such allergic reactions include, but are not limited to, asthma, rhinitis, and eczema.

[0373] Neutrokin- α conjugates and/or a Neutrokin- α complexes may be useful as an immunosuppressive agent.

[0374] A Neutrokin- α conjugate or a Neutrokin- α complex may be used to treat, prevent, and/or diagnose various immune system-related

- 134 -

disorders and/or conditions associated with these disorders, in mammals, preferably humans. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a Neutrokin- α conjugate or a Neutrokin- α complex that can inhibit an immune response, particularly the proliferation of B cells and/or the production of immunoglobulins, may be an effective therapy in treating and/or preventing autoimmune disorders. Thus, in preferred embodiments, a Neutrokin- α conjugate or a Neutrokin- α complex of the invention is used to treat, prevent, and/or diagnose an autoimmune disorder.

[0375] In another embodiment, AIDS is treated, prevented, and/or diagnosed using a Neutrokin- α conjugate or a Neutrokin- α complex.

[0376] In another embodiment, HIV infection is treated, prevented, and/or diagnosed using a Neutrokin- α conjugate or a Neutrokin- α complex.

[0377] A Neutrokin- α conjugate or a Neutrokin- α complex may also be employed to inhibit T-cell proliferation by the inhibition of IL-2 biosynthesis for the treatment of T-cell mediated autoimmune diseases.

[0378] All of the above described applications also apply to veterinary medicine.

[0379] The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

Formulation and Administration

[0380] The timing of the administration can vary substantially. In one embodiment, the entire dose is provided in a single bolus. Alternatively, the

- 135 -

dose can be provided by multiple administrations, such as an extended infusion method or by repeated injections administered over a span of weeks, for example six to twelve weeks between radioimmunotherapeutic doses. In one embodiment, the Neutrokin- α complex is administered at two week intervals. In another embodiment, the Neutrokin- α complex is administered over a span of about 2 to about 4 days.

[0381] Alternatively, a slow intravenous infusion of the complex may be administered, having a period for the infusion of about one to about 24 hours.

[0382] In one embodiment of the treatment of B-cell lymphoma, intravenous administration is utilized to deliver the Neutrokin- α complex to the site of the tumor. In another embodiment, intralymphatic routes of administration, such as subcutaneous or intramuscular injection or by catheterization of lymphatic vessels, are utilized.

[0383] In one embodiment, a treatment regimen comprises two dosages of a Neutrokin- α conjugate or Neutrokin- α complex. The first dose is an imaging dose. The second dose is a therapeutic dose. The imaging dose can be used to confirm that the Neutrokin- α conjugate or Neutrokin- α complex localizes to specific organs, tissues, and/or cells. Upon confirmation, or sometime thereafter, that the Neutrokin- α conjugate or Neutrokin- α complex localizes to certain organs, tissues, and/or cells, the second dose comprising a therapeutically effective amount of a Neutrokin- α conjugate or Neutrokin- α complex is administered to the subject.

[0384] In one embodiment, an imaging dose of a Neutrokin- α complex comprising a ion suitable for imaging, e.g., ^{111}In , is administered to a subject. The distribution of the Neutrokin- α complex is then assessed. In one embodiment, a first image of the of the subject is obtained from about 0.5 to about 24 hours after administration of the Neutrokin-complex. In another embodiment, a second image is obtained of the subject from about 24 hours to about 72 hours after administration of the Neutrokin- α complex. In a further embodiment, a third image is obtained of the subject from about 72 hours to about 120 hours after administration of the Neutrokin- α

- 136 -

complex. After the first, second, and/or third images are obtained, the biodistribution is determined and analyzed, and, if the biodistribution is deemed acceptable, a therapeutic dose of a Neutrokin- α complex comprising a ion suitable for therapy, e.g., ^{90}Y , is administered to said subject.

[0385] Various delivery systems are known and can be used to administer a Neutrokin- α conjugate or a Neutrokin- α complex of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, receptor-mediated endocytosis (see, e.g., Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)), etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The Neutrokin- α conjugate, Neutrokin- α complex, or a composition thereof may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[0386] In a specific embodiment, it may be desirable to administer the Neutrokin- α conjugate or Neutrokin- α complex or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous

- 137 -

material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

[0387] In another embodiment, the Neutrokin- α conjugate or the Neutrokin- α complex or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat *et al.*, in *Liposomes in therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

[0388] In yet another embodiment, the Neutrokin- α conjugate or the Neutrokin- α complex or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald *et al.*, *Surgery* 88:507 (1980); Saudek *et al.*, *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Press, Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy *et al.*, *Science* 228:190 (1985); During *et al.*, *Ann. Neurol.* 25:351 (1989); Howard *et al.*, *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of therapeutic target, e.g., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)).

[0389] Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

[0390] When treating a subject with cancer, the Neutrokin- α conjugate or the Neutrokin- α complex may be used in conjunction with additional chemotherapeutic agents that are useful for the treatment of cancer. In

- 138 -

particular, in one embodiment of the present invention, the Neutrokin- α complex or Neutrokin- α conjugate is administered, to patient in need of such treatment, along with one or more chemotherapeutic agents. In a further embodiment, one or more chemotherapeutic agents are selected from the group consisting of chemotherapeutic agents used to treat or prevent one or more conditions selected from the group consisting of non-Hodgkin's lymphoma, chronic lymphocytic leukemia, multiple myeloma. Such agents are well-known in the art.

[0391] When treating a subject with autoimmune disease, the Neutrokin- α conjugate or the Neutrokin- α complex may be used in conjunction with additional agents that are useful for the treatment of autoimmune diseases. In particular, in one embodiment of the present invention, the Neutrokin- α complex or Neutrokin- α conjugate is administered, to patient in need of such treatment, along with one or more immunosuppressants. In a further embodiment, one or more chemotherapeutic agents are selected from the group consisting of chemotherapeutic agents used to treat or prevent one or more conditions selected from the group consisting of systemic lupus erythematosus, rheumatoid arthritis, multiple sclerosis, Crohn's disease, diabetes, Wegener's granulomatous, myasthenia gravis, and asthma. Such agents are well-known in the art.

[0392] The radiometric dosage to be applied can vary substantially. The Neutrokin- α complex or a composition comprising a Neutrokin- α conjugate and a radionuclide can be administered at a dose of about 0.1 to about 100 mCi per 70 kg body weight. In another embodiment, the Neutrokin- α complex or a composition comprising a Neutrokin- α conjugate and a radionuclide can be administered at a dose of about 0.1 to about 50 mCi per 70 kg body weight. In another embodiment, the Neutrokin- α complex or a composition comprising a Neutrokin- α conjugate and a radionuclide can be administered at a dose of about 0.1, 0.5, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90 or 100 mCi per 70 kg body weight.

- 139 -

[0393] For example, lymphomas are known to be radiosensitive tumors. For immunodiagnostic imaging, trace-labeling by the complex may be used, typically 1-20 mg of Neutrokin- α protein is labeled with about 1 to 60 mCi of radioisotope. The dose may be somewhat dependent upon the isotope used for imaging; amounts in the higher end of the range, preferably 40 to 60 mCi, may be used with ^{99m}Tc ; amounts in the lower end of the range, preferably 1-20 mCi, may be used with ^{111}In . For imaging purposes, about 1 to about 30 mg of Neutrokin- α complex can be given to the subject. For radioimmunotherapeutic purposes, the Neutrokin- α complex is administered to a subject in sufficient amount so that the whole body dose received is up to about 1100 cGy, but preferably less than or equal to 500 cGy. The total amount of Neutrokin- α protein administered to a subject, including Neutrokin- α protein, Neutrokin- α conjugate and Neutrokin- α complex, can range from 1.0 $\mu\text{g/kg}$ to 1.0 mg/kg of patient body weight. In another embodiment, total amount of Neutrokin- α protein administered to a subject, can range from 20 $\mu\text{g/kg}$ to 100 $\mu\text{g/kg}$ of patient body weight.

[0394] An amount of radioactivity which would provide approximately 500 cGy to the whole body of a human is estimated to be about 825 mCi of ^{131}I . The amounts of radioactivity to be administered depend, in part, upon the isotope chosen. For ^{90}Y therapy, from about 1 to about 200 mCi amounts of radioactivity are considered appropriate, with preferable amounts being 1 to 150 mCi, and 1 to 100 mCi (*e.g.*, 60 mCi) being most preferred. The preferred means of estimating tissue doses from the amount of administered radioactivity is to perform an imaging or other pharmacokinetic regimen with a tracer dose, so as to obtain estimates of predicted dosimetry. In determining the appropriate dosage of radiopharmaceutical to administer to an individual, it is necessary to consider the amount of radiation that individual organs will receive compared to the maximum tolerance for such organs. Such information is known to those skilled in the art, for example, see Emami *et al.*, *International Journal of Radiation Oncology, Biology, Physics* 21:109-22

- 140 -

(1991); and Meredith, Cancer Biotherapy & Radiopharmaceuticals 17:83-99 (2002), both of which are hereby incorporated by reference in their entireties.

[0395] A "high-dose" protocol, for example in the range of 200 to 600 cGy (or higher) to the whole body, may require the support of a bone-marrow replacement protocol, as the bone-marrow is the tissue which limits the radiation dosage due to toxicity.

[0396] The composition to be administered may be given in a single treatment or fractionated into several portions and administered at different times. Administering the composition in fractionated doses may make it possible to minimize certain damage to non-target tissue. Such multiple dose administration may be more effective.

[0397] The present method may further comprise administering a second composition comprising one or more basic amino acids, wherein said second composition is administered prior to the Neutrokin- α complex, Neutrokin- α conjugate, or Neutrokin- α composition. Said second composition reduces renal accumulation of radioactivity.

[0398] The Neutrokin- α complex according to the present invention may be used as an imaging agent. Imaging agents are useful in a number of applications, include planar imaging, magnetic resonance imaging (MRI) applications, ultrasound imaging applications, and X-ray applications. Other applications in which imaging agents are useful include scintigraphic, positron emission tomography (PET), single photon emission computed tomography (SPECT), gamma scintigraphy, electrical impedance, light, or magnetometric imaging applications.

[0399] When used as an imaging agent, the Neutrokin- α complex may contain any suitable metal ion in accordance with the invention. In one embodiment, the imaging agents of the invention contain radionuclides suitable for use in PET or SPECT imaging. In another embodiment, the radionuclide used in the imaging agent is a radionuclide selected from the group consisting of ^{99m}Tc , ^{68}Ga , ^{62}Cu , and ^{111}In . In another embodiment, the radionuclide is ^{111}In .

- 141 -

[0400] The Neutrokin- α complex of the present invention may be administered to patients for imaging in amounts sufficient to yield the desired contrast with the particular imaging technique. Generally, dosages of from about 0.001 to about 5.0 mmoles of chelated imaging metal ion per kilogram of patient bodyweight are effective to achieve adequate contrast enhancements. For most MRI applications, preferred dosages of imaging metal ion will be in the range of from about 0.02 to about 1.2 mmoles/kg bodyweight while, for X-ray applications, dosages of from about 0.05 to about 2.0 mmoles/kg are generally effective to achieve X-ray attenuation. Preferred dosages for most X-ray applications are from about 0.1 to about 1.2 mmoles of the lanthanide or heavy metal compound/kg bodyweight. Where the chelated species is a radionuclide, dosages of about 0.01 to about 100 mCi, preferably 0.1 to 50 mCi, will normally be sufficient per 70 kg bodyweight.

[0401] Another embodiment of the present invention is a method of imaging the site of infection or inflammation in a patient comprising administering a Neutrokin- α -complex or a composition comprising a Neutrokin- α conjugate and a metal ion to a patient, preferably by injection or infusion; and imaging the patient, preferably by using either planar or SPECT gamma scintigraphy.

[0402] Another embodiment of the present invention is a method of imaging the site of cancer in a patient comprising administering a Neutrokin- α conjugate or Neutrokin- α complex to a patient by injection or infusion; and imaging the patient using either planar or SPECT gamma scintigraphy.

[0403] The Neutrokin- α complex of the present invention may be used as a contrast agent.

[0404] Also in accordance with the present invention, a method for diagnostic examination or therapeutic treatment of a mammal is provided. This method is based on the mechanism of receptor-mediated endocytosis activity and involves i) the movement of a Neutrokin- α complex that can be detected by external imaging techniques into the interior of a cell through invagination of the cell membrane. The Neutrokin- α protein serves to deliver the

- 142 -

chelated metal or a chemotherapy agent into a cell that expresses Neutrokin- α binding protein or a Neutrokin α receptor, thereby enabling diagnostic examination, radiotherapy or chemotherapeutic treatment of an organ or tissue comprising the cell.

[0405] In one embodiment, the method of the present invention comprises the steps of (a) administering to a subject, preferably a mammal, a composition comprising a Neutrokin- α conjugate or a Neutrokin- α complex and a pharmaceutically acceptable carrier and (b) monitoring the biodistribution of a metal ion. Said metal ion can be administered as part of the complex or separately as a solution.

[0406] Paramagnetic metals are used in affecting the relaxation times of nuclei in mammalian tissue. This is useful for magnetic resonance imaging processes. On the basis of differences in proton density and relaxation times, images of biological tissues can be obtained which may be used for diagnostic purposes. The greater the differences in the relaxation times of the nuclei which are present in the tissues being examined, the greater will be the contrast in the image that is obtained.

[0407] It is known that the relaxation times of neighboring nuclei can be affected by the use of paramagnetic salts. In solution, the paramagnetic salts are toxic in mammals. Hence, to reduce the toxic effect of paramagnetic metal ions administered for diagnostic purposes, they are combined with complex compounds, *i.e.*, chelating agents. The Neutrokin- α complex increases the concentration of the metal at locations containing a site with affinity for Neutrokin- α , for example, a B-cell containing a Neutrokin- α receptor (*e.g.*, BAFF receptor, TACI receptor, or BCMA receptor), thus providing increased contrast of the tissue comprising said site.

[0408] Doses for administration of paramagnetic metals in the complex of the present invention can be from about 0.05 to about 0.3 mmol/kg of body weight.

[0409] The metal complexes of the present invention find utility as diagnostic and/or therapeutic agents. Thus, the present invention provides methods for

- 143 -

the diagnosis of the presence and/or status of a disease state, or for the treatment of a disease state, comprising the step of administering a metal complex of the present invention to a subject in need thereof. The metal complexes of the present invention may be administered by an appropriate route such as orally, parentally (for example, intravenously), intramuscularly or intraperitoneally or by any other suitable method. For example, the complexes of this invention may be administered to a subject by bolus or slow infusion intravenous injection.

[0410] In accordance with the present invention, the Neutrokin- α conjugate, Neutrokin- α complex, and pharmaceutical compositions of the present invention can be administered by any means that achieve their intended purpose. For example, administration can be by subcutaneous, intravenous, intramuscular, intraperitoneal, buccal, or ocular routes, rectally, parenterally, intrasystemically, intravaginally, topically (as by powders, ointments, drops or transdermal patch), or as an oral or nasal spray. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

[0411] The effective dose of radiation or metal content to be utilized for any application will also depend upon the particulars of that application. In treating tumors, for example, the dose will depend, *inter alia*, upon tumor burden, accessibility and the like. Somewhat similarly, the use of metal chelate conjugated antibodies for diagnostic purposes will depend, *inter alia*, upon the sensing apparatus employed, the location of the site to be examined, and other similar factors.

[0412] Certain embodiments of the invention can be practiced either with scintigraphic or magnetic resonance imaging agents. A combination of these imaging agents can also be used, although this requires more complex instrumentation and data processing. Scintigraphic imaging according to the method of the invention is effected by obtaining a scintigram of the tissue or organ of interest, using as an imaging agent a Neutrokin- α conjugate or

- 144 -

Neutrokin- α complex. The scintigram is normally taken by a gamma imaging camera having one or more windows for detection of energies in the 50-500 keV range. Use of radioisotopes with higher energy, beta, or positron emissions would entail use of imaging cameras with the appropriate detectors, all of which are conventional in the art. The scintigraphic data can be stored in a computer for later processing.

[0413] Neutrokin- α conjugates and Neutrokin- α complexes of the invention may be administered alone or in combination with each other or other agents. In preferred embodiments, Neutrokin- α conjugates and Neutrokin- α complexes of the invention may be administered alone or in combination with Neutrokin- α protein.

[0414] In other embodiments, Neutrokin- α conjugates and Neutrokin- α complexes of the invention may be in combination with other chemotherapeutic agents or regimens.

[0415] In other embodiments, Neutrokin- α conjugates and Neutrokin- α complexes of the invention may be in combination with toxins or cytotoxic prodrugs. By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, cytotoxins (cytotoxic agents), or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, α toxin, ricin, abrin, Pseudomonas exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, α -sarcin and cholera toxin. "Toxin" also includes a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, *e.g.*, α -emitters such as, for example, ^{213}Bi , or other radioisotopes such as, for example, ^{103}Pd , ^{133}Xe , ^{131}I , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{35}S , ^{90}Y , ^{153}Sm , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn ,

- 145 -

⁷⁵Se, ¹¹³Sn, ¹¹⁷Sn, ¹⁸⁶Rhenium, ¹⁶⁶Ho, and ¹⁸⁸Rhenium; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[0416] A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis- dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

[0417] By "cytotoxic prodrug" is meant a compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

Kits

[0418] For radiopharmaceutical or radiotherapy applications it is convenient to prepare the complexes of the present invention at, or near, the site where they are to be used. A single- or multi-vial kit comprising all or some of the components needed to prepare the complexes of this invention, other than the

- 146 -

radionuclide ion itself, is an additional part of this invention. In one embodiment, the kit comprises all of the components for preparing a Neutrokin- α complex, other than the radionuclide itself.

[0419] The present invention also provides a kit comprising one or more containers, wherein each of said containers is filled with one or more of the ingredients of the pharmaceutical composition of the invention. Optionally, associated with said one or more containers is a set of written materials in the form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals or biological products, said notice reflects approval by the agency of manufacture, use or sale for human administration. The pharmaceutical kit optionally further comprises instructions to prepare said pharmaceutical compositions.

[0420] In one embodiment, a kit comprises a first container, said first container containing a Neutrokin- α protein, and a second container, said second container containing a chelator. Optionally, associated with said kit is a set of instructions providing the appropriate guidance on how to prepare Neutrokin- α conjugate using the contents of the kit.

[0421] In another embodiment, a kit comprises a first container, said first container containing a Neutrokin- α conjugate, and optionally associated with said kit is a set of instructions providing the appropriate guidance on how to prepare Neutrokin- α complex using the contents of the kit.

[0422] In another embodiment of the present invention, the kit comprises a first container, the contents of said container comprising a buffer, preferably an acetate buffer with concentration from about 10 mM to about 100 mM, preferably about 50 mM; a second container, the contents of said second container comprising a radionuclide, preferably a radionuclide selected from the group consisting of ^{90}Y and ^{111}In , wherein said radionuclide preferably is in solution, with a concentration of about 20 mCi/mL to about 200 mCi/mL, preferably about 80 mCi/mL; and a third container, the contents of said third container comprising a Neutrokin- α conjugate, preferably a Neutrokin- α conjugate which comprises a Neutrokin- α protein that has the

- 147 -

sequence of SEQ ID NO:3 and a chelator that is α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, wherein said conjugate is preferably in solution having a concentration of about 1 mg/mL to about 10 mg/mL, preferably about 2 mg/mL. The kit further optionally comprises a fourth container, the contents of said fourth container comprising a buffer, preferably an acetate buffer having a concentration of about 1 to about 50 mM, preferably about 10 mM, having a NaCl concentration of about 1 to about 500 mM, preferably about 140 mM, having a human serum albumin (HSA) concentration of about 1% to about 20%, preferably about 7% to about 8%, more preferably about 7.5%, having a pH of about 3-8, preferably about 6, and having a MeO-DOTA-glycine concentration of about 0.01 mM to about 100 mM, preferably about 1 mM.

[0423] In another embodiment of the present invention, the kit according to the present invention optionally comprises written material describing the use of a Neutrokin- α complex, a Neutrokin- α conjugate, or a composition comprising said conjugate or complex in a radioimmunodiagnostic and/or radioimmunotherapeutic protocol. The written material can be applied directly to a container, such as by applying a label directly to a vial containing said conjugate or complex. Alternatively, said a first container holding said conjugate or complex can be placed in a second container, such as a box, and the written material, in the form of a packaging insert, can be placed in the second container together with the first container holding said conjugate or complex.

[0424] The written portion of the article of manufacture may describe indications for prescribing the conjugate or complex. Such indications could be, for example, presentation of lymphoma at any site in the body. The written material could further describe that the conjugate or complex is useful for the treatment of lymphoma or other neoplasm clonally derived from a cell of B cell lineage, indicated as set forth above. In a preferred embodiment of the invention, the written material describes a Neutrokin- α complex to be used in the treatment, wherein said complex comprises a DOTA chelator, a

- 148 -

Neutrokin- α protein, and a radionuclide selected from the group consisting of ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{111}In , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{97}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{47}Sc , ^{67}Ga , ^{51}Cr , $^{177\text{m}}\text{Sn}$, ^{67}Cu , ^{167}Tm , ^{95}Ru , ^{188}Re , ^{177}Lu , ^{199}Au , ^{203}Pb , and ^{141}Ce , more preferably a selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac . In a most preferred embodiment, the written material will describe that the Neutrokin- α conjugate or Neutrokin- α complex is used in the treatment of lymphoma. In other preferred embodiments, the written material describes a Neutrokin- α complex to be used in the treatment, wherein said complex comprises a MeO-DOTA-NCS chelator, a Neutrokin- α protein, and a radionuclide selected from the group consisting of a ^{90}Y and ^{111}In . Still further, the written material can describe that the appropriate radiometric dose to be administered for an immunodiagnostic scanning is provided by 1 to 35 mCi of radioisotope, while the appropriate dose for therapeutic administration should be below 150 cGy to the whole body if bone marrow replacement support cannot be provided, but can be as high as 600 cGy to the whole body if bone marrow replacement support is provided. The doses for particular isotopes, especially as set forth herein below, might also be described.

[0425] The written material is preferably provided in the form required by the Food and Drug Administration for a package insert for a prescription drug. The written material may indicate that the antibody would be prescribed for use in patients having a diagnosis of B cell lymphoma and can be administered to patients presenting lymphoma in any site in the body. The written material may indicate that the conjugate or complex as described herein is useful as an initial or secondary treatment or in combination with other treatments.

[0426] The written material may also describe any possible side effects. The written material may also describe any possible contraindications.

[0427] The written material also may indicate that general radiologic and nuclear medicine precautions appropriate to the isotope used for labeling the antibody should be observed.

- 149 -

Assays

[0428] The invention also provides a method of screening a Neutrokin- α conjugate or Neutrokin- α complex to identify a complex or conjugate which binds to a Neutrokin- α receptor.

[0429] In the assay of the invention for a complex or conjugate that binds a Neutrokin- α receptor (*e.g.*, BAFF-R (SEQ ID NO:5), TACI (SEQ ID NO:7), and BCMA (SEQ ID NO:9), a cellular compartment, such as a membrane or a preparation thereof, may be prepared from a cell that expresses a Neutrokin- α receptor. Exemplary cell lines that may be used for this purpose include, but are not limited to B lineage immortalized hematopoietic cell lines such as IM-9 (ATCC CCL-159); Reh (ATCC CRL-8286); ARH-77 (ATCC CRL-1621); Raji (ATCC-CCL-86); Namalwa (CRL-1432); and RPMI 8226 (ATCC CCL-155). The preparation is incubated with labeled Neutrokin- α (*e.g.*, fluorescently labeled Neutrokin- α or radiolabeled Neutrokin- α) in the absence or the presence of the candidate conjugate or complex. The ability of the candidate conjugate or complex to bind the Neutrokin- α receptor is reflected in decreased binding of the labeled Neutrokin- α . Alternatively, a similar assay may be performed on live intact cells, with the final detection step comprising flow cytometry analysis.

[0430] Additionally assays which measure the direct binding of Neutrokin- α conjugates or complexes to one or more Neutrokin- α receptors. One example of such an assay is one that would utilize surface plasmon resonance technology (Biacore, Piscataway, NJ) to analyze interactions between biomolecules. Biacore assays are well known and routine to those of skill in the art.

Definitions

[0431] The term "Neutrokin- α conjugate," when used herein, means a Neutrokin- α protein that is covalently bonded to a chelator molecule.

- 150 -

[0432] The term "Neutrokin- α complex," when used herein, means a Neutrokin- α conjugate that is associated with a radionuclide, metal ion, or other ion that is able to chelate with the chelator of said conjugate.

[0433] The term "chelator," as used herein, refers a molecule or molecular fragment, at least part of which is able to associate with, or chelate, a metal ion. As used herein, a chelator molecule may optionally contain a linker group which connects the metal chelating portion of the chelator to the Neutrokin- α protein. Chelators containing such linking moieties are well known in the art. See, for example, Saji, J. "Targeted Delivery of Radiolabeled Imaging and Therapeutic Agents: Bifunctional Radiopharmaceuticals," *Crit. Rev. Therap. Drug Carrier Systems* 16:209-244 (1999); Liu, S. et al., "Bifunctional Chelators for Therapeutic Lanthanide Radiopharmaceuticals," *Bioconjugate Chem.* 12:7-34 (2001); *Kirk-Othmer Concise Encyclopedia of Chemical Technology*, pages 242-244 (John Wiley & Sons, Inc. 1985); and *Van Nostrand's Scientific Encyclopedia*, pages 613-615, Editor: Douglas M. Considine, P.E. (Eighth Edition, Van Nostrand Reinhold, 1995), each of which is hereby incorporated by reference in its entirety. The term "chelator moiety" or "chelating moiety" refers to the portion of the chelator molecule which forms a noncovalent interaction with a metal ion.

[0434] The term "MeO-DOTA-NCS" as used herein refers to the compound α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid having the CAS registry number 130707-79-8.

[0435] The term "MeO-DOTA-glycine," as used herein, refers to the compound formed by reacting MeO-DOTA-NCS with glycine.

[0436] An "effective amount" of the formulation is used for therapy. As is known by one of ordinary skill in the art, the exact amount which constitutes an effective amount may vary from one situation to another. As used herein, the term "effective amount" refers to a dosage or amount of a conjugate, complex, or composition of the present invention, said dosage or amount being sufficient to effect a sufficient or significant, intended result. For example, the amount of a Neutrokin-complex used for imaging may, and

- 151 -

probably will, be different than the amount used to treat a B-cell mediated condition. Furthermore, the exact amount of a Neutrokin-alpha complex used to treat non-Hodgkin's lymphoma may be different than the amount used to treat multiple myeloma. The dose will vary depending on the disease being treated. Therapeutic doses will be administered in sufficient amounts to reduce pain, inhibit tumor growth, cause regression of tumors, and/or kill the tumor. The amount of radionuclide needed to provide the desired therapeutic dose may be determined experimentally and optimized for each particular composition. The amount of radioactivity required to deliver a therapeutic dose may vary with the individual composition employed. The dosage to be administered may be given in a single treatment or fractionated into several portions and administered at different times.

[0437] As used herein, "pharmaceutically acceptable salt" means any salt of a compound of formula (I) which is sufficiently non-toxic to be useful in therapy or diagnosis of mammals. Thus, the salts are useful in accordance with this invention. Representative of those salts, which are formed by standard reactions, from both organic and inorganic sources include, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, palmoic, mucic, glutamic, d-camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, steric, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic acids and other suitable acids. Also included are salts formed by standard reactions from both organic and inorganic sources such as ammonium, alkali metal ions, alkaline earth metal ions, and other similar ions. Particularly preferred are the salts of the compounds of formula (I) where the salt is potassium, sodium, ammonium, or mixtures thereof.

EXAMPLES

EXAMPLE 1

- 152 -

Rapid and Specific Targeting of Radiolabeled Neutrokin- α to Lymphoid Tissues

[0438] Here, biodistribution studies of radiolabeled Neutrokin- α are reported that demonstrate high *in vivo* targeting specificity of Neutrokin- α for lymphoid tissues. Neutrokin- α was radiolabeled with ^{125}I and injected intravenously into BALB/c mice. Three doses and 4 timepoints over a 24-hr period were studied. Biodistribution was measured by direct counting of the radioactivity in dissected whole organs or tissues and by whole body quantitative autoradiography (QAR).

[0439] Spleen and lymph nodes showed the highest concentration of radioactivity among the dissected organs and tissues. Three hours after injection of 0.01 mg/kg Neutrokin- α , 63% and 23% injected dose (ID)/g were measured in spleen and lymph node, respectively, compared to ~5% for both kidney and liver. As the dose was increased to 0.05 mg/kg or 0.3mg/kg, the %ID/g in spleen and lymph node decreased but was unchanged in liver and kidney, suggesting that targeting to spleen and lymph nodes is mediated by saturable binding. With increasing time, the ratio of the concentration in spleen and lymph node to the concentration in either kidney or liver increased. QAR confirmed the high uptake of radiolabeled Neutrokin- α in spleen and lymph nodes at 3 hr, and revealed high uptake in bone marrow, gut-associated lymphoid tissue (GALT) and intestinal contents as well. At 24 hr, spleen, lymph nodes and GALT were still strongly positive for radiolabeled Neutrokin- α by QAR whereas liver and kidney no longer had observable levels. A cytotoxic radionuclide coupled to Neutrokin- α could irradiate neoplastic B-cells trafficking through or residing in lymphoid tissues. Thus, the rapid and highly specific targeting of radiolabeled Neutrokin- α to lymphoid tissues provides a rationale for its application in the treatment of B-cell malignancies.

[0440] A similar biodistribution analysis was performed with Neutrokin- α attached chelator according to Formula I labeled with ^{111}In (NA-MeO-DOTA- ^{111}In). Female BALB/c mice weighing approximately 20g were used

- 153 -

to determine the biodistribution of NA-MeO-DOTA-¹¹¹In. Mice were injected intravenously via the tail vein with 0.1mL/g at 5µg/mL NA-MeO-DOTA-¹¹¹In to give a protein dose of 50 µg/kg. The specific activity of the NA-MeO-DOTA-¹¹¹In was 13.6 µCi/µg, resulting in approximately 13.6 µCi per injected animal. Prior to injection, 84% of the radioactivity in the dosing solution was determined to be TCA-precipitable (protein-associated). Biodistribution was determined by both tissue dissection and quantitative whole body autoradiography (QWBA). Timepoints of 3, 6, 24 72 and 96 hours post injection were studied. The following tissues were dissected and studied: spleen, mesenteric lymph node, kidney, part of the liver, stomach, part of the duodenum, heart, lung, thymus (72 and 96 hours only) skeletal muscle (biceps femoris) and part of the femur.

[0441] Similar to the biodistribution of ¹²⁵I labeled Neutrokin-α, the highest % injected dose (%ID) were found in the spleen and mesenteric lymph nodes (C max = 63.4 and 24.6 %ID/g, respectively). This localization agrees with the expression of the three known Neutrokin-α receptors (BCMA, TACI, and BAFF-R) found predominantly on mature, immunoglobulin-positive B cells. The activity observed in the femur Cmax=8.4%ID/g is believed to be due to the localization of NA-MeO-DOTA-¹¹¹In to mature B cells in the bone marrow and the activity in the duodenum (Cmax=7.2%ID/g) may represent localization to B cells in the lamina propria of the gut. Significant radioactivity was also found in the liver (Cmax=19%ID/g). The kidneys, stomach, heart, lung, muscle and thymus all had relatively low %ID/g. The high radioactivity seen in the liver is believed to be due to retention of a metabolic end product, ¹¹¹In-chelator-lysine. Because of its ionic character at intralysosomal pH, ¹¹¹In-chelator-lysine becomes trapped intracellularly at the site of metabolism. In contrast, ¹²⁵I labeled Neutrokin-α studies did not reveal high levels of radioactivity in liver because the free ¹²⁵I released by the intracellular metabolism of halogenated proteins can rapidly diffuse across cell membranes and is not retained at the site of metabolism.

- 154 -

EXAMPLE 2:

Pharmacological Effects of ^{131}I -labeled Neutrokin- α in BCL1 Tumor-Bearing Mice and J558 Tumor Bearing Mice

^{131}I -Neutrokin- α administration to BCL1 Tumor-Bearing Mice

[0442] The BCL1 tumor cell line was derived from a spontaneous murine B cell tumor. Intraperitoneal inoculation of the BCL1 cell line in BALB/c mice results in splenomegaly, and subsequent death. The BCL1 tumor cell phenotype is IgM positive, complement receptor negative, Fc receptor positive and has marginal IgD expression (Knapp *et al.*, *J. Immunol.* 123:992-999 (1979) and Vitetta *et al. Blood* 89:4425-36. (1997)). Based on FACS analysis using biotinylated Neutrokin- α , BCL1 cells freshly isolated from the spleens of BALB/c mice express Neutrokin- α receptors on their cell surface. The BCL1 tumor model is a relevant mouse model for human B cell lymphoma, providing a means to test the ability of ^{131}I -labeled Neutrokin- α to kill leukemic B cells and consequently prolong survival of tumor-bearing mice. Three lots of ^{131}I -labeled Neutrokin- α (Lots TX1, TX2 and TX3) were prepared by MDS Nordion (Ontario, Canada) and used in 3 different experiments to evaluate the effects of ^{131}I -labeled Neutrokin- α in this murine model.

[0443] Female BALB/c mice were injected intraperitoneally (ip) on Day 0 with 1×10^5 viable BCL1 cells that had been propagated in vivo. Treatment groups for the 3 experiments are described in Table VI. Ten days after injection of tumor cells, the animals were administered ^{131}I -labeled Neutrokin- α iv in 110 μL . The doses administered were 11.9 or 15.3 mCi/kg (TX1), 17.5 mCi/kg (TX2), or 37.7 mCi/kg (TX3) for the 3 experiments. To identify potentially toxic effects of the administered ^{131}I -labeled Neutrokin- α , age-matched control BALB/c mice without BCL1 tumors were injected with identical doses of the ^{131}I -labeled protein. An additional group of BALB/c mice, bearing BCL1 tumors and receiving an iv

- 155 -

injection of the vehicle, served as the normal tumor control group. Survival was then monitored for 48, 44, or 40 days for the TX1, TX2, and TX3 experiments, respectively.

Table VI Treatment groups for TX1, TX2 and TX3 experiments

<u>Exp.</u>	<u>Group</u>	¹³¹ I-Neutrokin- α Dose (mCi/kg)	<u>n</u>	BCL1 Tumor Inoculated ip (No. of cells)
1 (TX1)	1 Vehicle	0	15	1 x 10 ⁵
	2 ¹³¹ I-Neutrokin- α	11.9	10	1 x 10 ⁵
	3 ¹³¹ I-Neutrokin- α	15.3	10	1 x 10 ⁵
	4 ¹³¹ I-Neutrokin- α	11.9	10	0
	5 ¹³¹ I-Neutrokin- α	15.3	10	0
2 (TX2)	1 Vehicle	0	12	1 x 10 ⁵
	2 ¹³¹ I-Neutrokin- α	17.5	12	1 x 10 ⁵
	3 ¹³¹ I-Neutrokin- α	17.5	8	0
3 (TX3)	1 Vehicle	0	14	1 x 10 ⁵
	2 ¹³¹ I-Neutrokin- α	37.7	14	1 x 10 ⁵
	3 ¹³¹ I-Neutrokin- α	37.7	8	0

[0444] The endpoint monitored in the 3 experiments was survival (days) following ip inoculation of BCL1 tumor cells. All animals were examined daily. The day post-inoculation that mice were either found dead or in moribund condition (the latter being immediately euthanized for humane reasons) was recorded.

[0445] A single iv administration of either 11.9 or 15.3 mCi/kg (TX1), 17.5 mCi/kg (TX2), or 37.7 mCi/kg (TX3) of ¹³¹I-labeled Neutrokin- α injected 10 days after intraperitoneal inoculation of BCL1 cells in BALB/c mice significantly improved survival compared with mice inoculated with tumor and treated with the ¹³¹I-labeled Neutrokin- α vehicle (Figures 2-4; in Figures 2-4, ¹³¹I-labeled Neutrokin- α is indicated as LR131). The median survival time for the vehicle-treated, tumor-bearing mice was 18, 21, and 19 days post-tumor cell injection for the TX1, TX2, and TX3 experiments, respectively. In the TX1 experiment, ¹³¹I-labeled Neutrokin- α administration at dose levels of 11.9 and 15.3 mCi/kg doubled the median survival time of tumor-bearing mice to 35.5 (11.9 mCi/kg) and 34 (15.3

- 156 -

mCi/kg) days post-treatment, respectively. In the TX2 and TX3 experiments, ^{131}I -labeled Neutrokin- α administration at a dose of 17.5 or 37.7 mCi/kg increased the median survival time of tumor-bearing mice to 30 and 22 days post-treatment, respectively. Tumor-bearing mice treated with all doses of ^{131}I -labeled Neutrokin- α in the 3 experiments had a significantly lower risk of dying than tumor-bearing mice treated with vehicle (Table VII).

Table VII Incidence of mortality for TX1 – TX3 experiments

<u>Experiment</u>	<u>Treatment Group</u>	<u>Median Survival Time (Days)</u>
TX1	1, BCL1 + ^{131}I -labeled Neutrokin- α (11.9 mCi/kg)	35.5
	2, BCL1 + ^{131}I -labeled Neutrokin- α (15.3 mCi/kg)	34
	3, BCL1 Tumor Only	18
	4, No Tumor + ^{131}I -labeled Neutrokin- α (11.9 mCi/kg)	> 48
	5, No Tumor + ^{131}I -labeled Neutrokin- α (15.3 mCi/kg)	> 48
TX2	1, BCL1 + ^{131}I -labeled Neutrokin- α (17.5 mCi/kg)	30
	2, BCL1 Tumor Only + vehicle	21
	3, No Tumor + ^{131}I -labeled Neutrokin- α (17.5 mCi/kg)	> 44
TX3	1, BCL1 + vehicle	19
	2, BCL1 + ^{131}I -labeled Neutrokin- α (37.7 mCi/kg)	22
	3, No tumor + ^{131}I -labeled Neutrokin- α (37.7 mCi/kg)	> 40

[0446] In the TX1 – TX3 series of experiments, the effect that increasing the dose of ^{131}I -labeled Neutrokin- α had on the survival of the BCL1 tumor-bearing animals was investigated. A maximal survival benefit was achieved with the low doses of ^{131}I -labeled Neutrokin- α (11.9 and 15.3 mCi/kg). The much reduced effectiveness of ^{131}I -labeled Neutrokin- α in TX3 may be due to toxicity associated with the high dose of the material used.

[0447] In conclusion, a single iv administration of ^{131}I -labeled Neutrokin- α administered to mice bearing BCL1 leukemia cell splenic tumors significantly improved survival compared with tumor-bearing mice treated with vehicle.

^{131}I -Neutrokin- α administration to J558 Tumor-Bearing Mice

- 157 -

[0448] In a similar experiment as that described above, BALB/c mice were injected subcutaneously with J558 plasmacytoma cells (ATCC # TIB-6) and treated with a single intravenous treatment of 25mCi/kg of ^{131}I -labeled Neutrokin- α . 24 BALB/c mice (NCI, 4 weeks old, average weight 18 g) were divided into 2 groups (12 mice per group) and injected sc with 2.5×10^5 J558 cells in 100 mL of PBS. At Day 9 after injection, mice in Group 1 were injected intravenously with 100 mL of formulation buffer, and mice in Group 2 were injected iv with a dose of 25 mCi/kg of ^{131}I -Neutrokin- α in 100 mL of formulation buffer. The average body weight at the time of ^{131}I -Neutrokin- α injection was 19.5 g.

[0449] Two parameters were evaluated during this study the tumor size and the time to tumor response. To evaluate tumor size the short and long axes of the tumor were measured using an electronic digital caliper. Tumor size was calculated by multiplication of the lengths of the short and long axes and expressed in mm^2 . The time to tumor response was characterized by the day after cell inoculation when a visible tumor ($> 2\text{mm}$) was detected on a mouse. In addition, mice were monitored for survival and signs of radiation induced toxicity (general appearance, activity, breathing frequency, stool consistence).

[0450] One mouse in the ^{131}I -Neutrokin- α -treated group died on Day 25 (16 days after ^{131}I -Neutrokin- α treatment) with no obvious signs of radiation related toxicity. A second mouse died in the same group on Day 30, when all animals in the control group were terminated because of large tumor size.

[0451] The first tumors of measurable size were detected at Day 14 in the buffer control group, where 4 out of 12 animals developed tumors. In the ^{131}I -Neutrokin- α treated animals, tumor formation was delayed by 6 days. Only one mouse out of 12 developed a tumor at Day 20. At Day 22, there was only one tumor-bearing mouse in the ^{131}I -Neutrokin- α treated group out of 12 animals, whereas in the buffer control group, 11 out of 12 mice developed tumors of different sizes. At Day 27, the mean tumor size in the buffer control group was 489 mm^2 (all tumor positive mice in this group were

- 158 -

terminated at this time point). In the ^{131}I -Neutrokin- α treated group, the mean tumor size was 32.7 mm², 15 times smaller than in the buffer control group. Taken together, these data suggest a strong inhibition of J558 tumor development in mice treated with ^{131}I -Neutrokin- α at a dose of 25 mCi/kg and tumor load of 2.5×10^5 cells/mouse.

[0452] In conclusion, a single intravenous administration of ^{131}I -Neutrokin- α into BALB/c mice at a dose of 25 mCi/kg significantly inhibits subcutaneous growth of J558 plasma cell tumors. At the initial tumor load of 2.5×10^5 cells/mouse, a 6 day delay in tumor formation and a 15-fold reduction in tumor size was observed in ^{131}I -Neutrokin- α treated animals.

[0453] The anti-neoplastic effects of ^{131}I -Neutrokin- α were accompanied by the expected B lymphocyte hypoplasia and a transient (<20 days) depletion of cKit⁺ bone marrow precursors and peripheral platelets. Peripheral neutrophil, red blood cell, and monocyte counts were unaffected by ^{131}I -Neutrokin- α treatment. Taken together, the results demonstrate that ^{131}I -Neutrokin- α inhibits in vivo tumor growth in two models of B cell neoplasia. Moreover, ^{131}I -Neutrokin- α efficacy was not accompanied by significant bone marrow toxicities or peripheral myelosuppression.

EXAMPLE 3:

Method For Producing Neutrokin-Alpha Using A Stringent Promoter And Low Expression Level

[0454] Neutrokin- α has been produced in *Escherichia coli* K-12 from the periplasmic fraction of the cell lysate. Using this system, soluble, properly folded, active Neutrokin- α is not obtainable from simple shake flask experiments. Yields of soluble Neutrokin- α from complex media fermentations in small and large-scale bioreactors are on the order of 1-5 mg/L. Greater yields (25-38 mg/L) of soluble, properly folded, active Neutrokin- α can be accomplished in bioreactors at low to medium cell

- 159 -

density under defined medium conditions. Moreover, this low quantity of protein is difficult to purify via conventional methods.

[0455] This example describes a method for the production of high yields of soluble, properly folded, active Neutrokin- α in the periplasm of *Escherichia coli*, which permits the use of conventional methods for Neutrokin- α purification, such as those described below. Additionally, Neutrokin- α protein may be purified using affinity columns comprising Neutrokin- α binding peptides such as those described in WO 02/02641, which is herein incorporated by reference in its entirety. Purified Neutrokin- α may be quantified using RP-HPLC.

[0456] This method relies on the expression of Neutrokin- α protein from the bacterial *phoA* promoter. The *phoA* promoter is a very tightly regulated system that exhibits a very low level of transcription in the presence of excess phosphate. As the phosphate level in the medium decreases below a threshold of ~ 4 micromolar (Wanner, B.L., *J Cell Biochem* 51:47 (1993)), transcription is induced about 1000-fold. The *phoA* promoter yields a gradual build-up of recombinant protein, instead of a sharp increase of induction that occurs with other systems. This gradual or steady increase in recombinant protein minimizes the chance of overwhelming the components of the bacterial expression system and may also minimize the formation of inclusion bodies. Furthermore, this gradual build up permits the expression of proteins that might have been toxic to the cell if they were induced to high levels over a short period of time.

Expression Vector pML124

[0457] The expression vector, pML124, was created using pBR322 as the starting backbone. First, the endogenous NdeI site of pBR322 was eliminated by digesting it with NdeI, filling in the overhanging ends with the Klenow enzyme, then re-ligating the two blunt-ends back together (this created pML123). Next, pML123 was digested with EcoRI and BamHI restriction

- 160 -

enzymes and the linear plasmid (loss of ~375 bp of DNA) was agarose gel purified (Qiagen).

[0458] The *phoA* promoter region was PCR-amplified from the *E. coli* K-12 chromosome (W3110; ATCC Catalogue No. 27325) with EcoRI (5') and BamHI (3') engineered sites. NdeI and KpnI sites were also engineered downstream of the *phoA* promoter to facilitate cloning of recombinant genes. Finally, the Shine-Dalgarno (SD) box was optimized for protein expression. The wild-type SD box and its adjacent sequence is as follows (the putative SD boxes are underlined and in bold):

5'-TTTGTACATGGAGAAAATAAA (SEQ ID NO:12):-[ATG, start of coding sequence]-3'

Optimized SD box and adjacent sequence is as follows:

5'-CACGTAAAGGAAGTATCTCAT (SEQ ID NO:13):-[ATG, start of coding sequence]-3'

[0459] The digested (EcoRI and BamHI) and purified *phoA* promoter PCR product was ligated into the agarose gel purified pML123 (described above). The ligation mixture was transformed into highly competent *E. coli* cells using standard techniques. Positive clones were identified via restriction analysis and DNA sequencing.

[0460] pML124 contains a gene for ampicillin resistance, a ColE1 replicon (pBR322-based), Rop, *phoA* promoter, the optimized Shine-Dalgarno (SD) box (above) and a multiple cloning site. Figure 5 is a plasmid map of pML124 and SEQ ID NO:14 is the nucleotide sequence of pML124. Additionally, plasmid pML124 was deposited at the American Type Culture Collection (ATCC) on October 8, 2001 and given ATCC Deposit No. PTA-3778. ATCC Deposit Nos. PTA-3778 was made pursuant to the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. The ATCC (American

- 161 -

Type Culture Collection) is located at 10801 University Boulevard, Manassas, Virginia 20110-2209.

Neutrokin- α Expression Vector pML124-MBPssBLyS

[0461] A fusion construct of the maltose binding protein signal sequence (MBPss) and Neutrokin- α was placed behind the *phoA* promoter in pML124 as follows. A 549 bp NdeI/KpnI MBPss-Neutrokin- α containing DNA insert was ligated into NdeI/KpnI digested and gel purified pML124 to form pML124-MBPss-BLyS. (Figure 6, SEQ ID NO:15 ATCC Deposit No. PTA-3867, deposited November 16, 2001). ATCC Deposit No. PTA-3867 was made pursuant to the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. The ATCC (American Type Culture Collection) is located at 10801 University Boulevard, Manassas, Virginia 20110-2209.

[0462] The pML124 plasmid (Figure 5, SEQ ID NO:14) is described in above and in U.S. Provisional Applications 60/329,508 filed October 17, 2001, 60/329,747 filed October 18, 2001 and 60/331,478 filed November 16, 2001 which are herein incorporated by reference in their entireties. The *phoA* promoter region is located at nucleotides 111-410 SEQ ID NOs:14 and 15. The MBP signal sequence is encoded by nucleotides 423-500 of SEQ ID NO:15 and nucleotides 501-959 of SEQ ID NO:15 encode amino acids 134-285 of Neutrokin- α (SEQ ID NO:2). The amino acid sequence of the MBP signal sequence is shown in SEQ ID NO:16 and the amino acid sequence of the full length MBP signal sequence-Neutrokin- α protein encoded by the pML124-MBPss-BLyS vector is shown in SEQ ID NO:17.

Neutrokin- α Expression in *E. coli*

[0463] Plasmid pML124-MBPss-BLyS was transformed into *E. coli* cells, e.g. K-12 based strains, by standard methods. Ampicillin resistant transformants were screened for the proper DNA insert by restriction enzyme analysis and

- 162 -

DNA sequence. For example, digestion of pML124-MBPss-BLyS™ with NdeI and KpnI results in two nucleotide fragments: 549 and 4,431 base pairs in length. Positive clones were subsequently grown in City Broth-Low Phosphate media (see recipe below). Neutrokin- α expression levels were examined via SDS-PAGE and subsequent Coomassie staining. Using simple shake flask experiments, more than 260 mg/L of Neutrokin- α was obtained.

[0464] Next, positive clones were grown to high cell density in complex media in small scale bioreactors, similar to the method described by Joly *et al.*, *PNAS* 95:2773-2777 (1998), which is hereby incorporated by reference in its entirety. Specifically, the initial fermentation medium for the 5L bioreactor was composed of 55.7 mM ammonium sulfate, 13.9 mM sodium monobasic phosphate, 21.9 mM potassium dibasic phosphate, 5 mM sodium citrate, 29.6 mM potassium chloride, 14.7 mM magnesium sulfate, 1.11% NZ-amine AS, 1.11% yeast extract, 5 g/L glucose, 0.002% ferric chloride, 25 μ g/ml kanamycin. A trace element solution (2.5 ml/3.4 L) was added containing 100 mM ferric chloride plus 30 mM of the following components: zinc sulfate, cobalt chloride, sodium molybdate, copper sulfate, boric acid and manganese sulfate. The fermenter was operated at 30°C, 650 rpm agitation, 10 standard liter/minute aeration. When the initial glucose was depleted, a concentrated glucose solution (50%) was added until the dissolved oxygen (DO) concentration reached 20% of air saturation as measured by an on-line oxygen electrode. When the optical density (600 nm) reached 40 OD600, a solution of 20% NZ amine AS, 20% yeast extract was fed at 0.2 ml/min for the rest of the fermentation. Neutrokin- α production was on the order of 260-570mg/L.

- 163 -

Low Phosphate Containing Media:

City Broth-Low Phosphate:

[0465] 30mM $(\text{NH}_4)_2\text{SO}_4$; 2.25 mM NaCitrate- $2\text{H}_2\text{O}$; 12mM MgSO_4 ; 15 mM KCl; 5% Yeast extract; 2% Casamino acids; 110 mM MOPS; 33 mM Glucose; pH 7.3

Vegan City Broth-Low Phosphate:

[0466] 30mM $(\text{NH}_4)_2\text{SO}_4$; 2.25 mM NaCitrate- $2\text{H}_2\text{O}$; 12mM MgSO_4 ; 15 mM KCl; 5% Phytone; 2% Casamino acids; 110 mM MOPS; 33 mM Glucose; pH 7.3

[0467] The only difference between the two media is that Phytone is substituted for Yeast extract in the Vegan recipe.

Purification of Neutrokin- α

[0468] 10 grams of *E. coli* cell paste are suspended in 50 milliliters of 5mM sodium citrate, pH 6.0 and placed at 4°C for 1 hour with gentle shaking. Cells are then disrupted by passing them through an M-Y110 Microfluidizer® Processor (Microfluidics, Inc., Newton, MA) set at 7500 psi four times. The suspension is then centrifuged at 22,000 x g for twenty minutes at 4°C using a Sorvall SLA-1500 rotor. The supernatant is then collected and filtered through a 0.45 micron bottle top filter (Nalgene).

[0469] Filtered supernatant is then loaded at 9 centimeters/hour on a Fast Flow Sepharose DEAE column (Amersham Biosciences, Piscataway, NJ) previously equilibrated with 5mM sodium citrate, pH6.0 (equilibration buffer). After loading, the column is washed with 5 to 10 column volumes of equilibration buffer. The Neutrokin- α protein is eluted with a 200mM NaCl step in equilibration buffer. Buffers used with the Fast Flow Sepharose DEAE chromatography column are pre-filtered using a 0.22 micron CA bottle

- 164 -

top filter (Nalgene) and pre-chilled to 4°C. The Fast Flow Sepharose DEAE column is used at 4°C. Prior to use, columns are cleaned with 0.5 M NaOH.

[0470] Relevant fractions, as determined by the ratio of contaminating proteins to Neutrokin- α protein seen in Coomassie stained SDS-PAGE gels, are pooled and diluted 1:1 with 10mM sodium citrate, pH 6.0, 2M (NH₄)SO₄. Pooled fractions are loaded at 17 centimeters/hour onto a Polypropylene Glycol Hydrophobic Interaction chromatography column (Tosoh Biosep, Montgomeryville, PA) previously equilibrated with 10mM sodium citrate, pH 6.0, 1M (NH₄)SO₄ (loading buffer). After loading, the column is washed with 5-10 volumes of loading buffer. The Neutrokin- α protein is eluted with a 5 column volume gradient from loading buffer to elution buffer (10mM sodium citrate, pH 6.0). Neutrokin- α elutes in the second peak toward the end of the gradient absorbance at 280nm. Buffers used with the Polypropylene Glycol Hydrophobic Interaction chromatography column are pre-filtered using a 0.22 micron CA bottle top filter (Nalgene) and used at room temperature. The Polypropylene Glycol Hydrophobic Interaction chromatography column is also used at room temperature. Prior to use, columns are cleaned with 0.5 M NaOH.

[0471] Relevant fractions, determined by the ratio of contaminating proteins to Neutrokin- α protein as monitored by Coomassie stained SDS-PAGE gels, are pooled and are dialyzed overnight (12 hours) into 50mM Tris, pH 7.4, 50mM NaCl at 4°C. The dialyzed pool is then loaded onto a POROS PI-50 anion exchange chromatography column (Applied Biosystems, Foster City, CA), previously equilibrated with 50mM Tris, pH 7.4, 50mM NaCl, at 17 centimeters/hour. After loading, column is washed with 5-10 volumes of loading buffer. Neutrokin- α is eluted using a pH step from 50mM Tris, pH 7.4, 50mM NaCl buffer to 50mM sodium citrate, pH 6.0. Relevant fractions, as determined by the ratio of contaminating proteins to Neutrokin- α protein seen in Coomassie stained SDS-PAGE gels, are pooled and stored at 4°C. Buffers used with the POROS PI-50 anion exchange chromatography column are pre-filtered using a 0.22 micron CA bottle top

- 165 -

filter (Nalgene) and pre-chilled to 4°C. The POROS PI-50 anion exchange chromatography column is used at 4°C. Prior to use, columns are cleaned with 0.5 M NaOH.

[0472] This purification protocol yields 0.5-1 milligram per gram of starting cell paste based on BCA protein assay (Pierce Biotechnology, Rockford, IL) and absorbance at 280 nanometers. The protein is 96% pure as determined by reverse phase-high performance liquid chromatography (RP-HPLC). Native-PAGE and size exclusion chromatography-HPLC (SEC-HPLC) analysis indicates the protein is predominantly in trimeric form.

[0473] The production of MBPss-Neutrokin- α under control of the *phoA* promoter allowed more stringent, slower expression, and resulted in increased yields. In summary, the production of Neutrokin- α from the *phoA* system is scaleable and achieves 10 to 20-fold more soluble, properly folded, active material than the current system.

[0474] Alternatively, Neutrokin- α can be purified by any method known in the art. In a non limiting example, Neutrokin- α proteins may be purified by affinity chromatography using columns to which Neutrokin- α binding peptides are attached. Neutrokin- α binding peptides that may be used in affinity purification of Neutrokin α are described for example, in WO2002/16411 and WO2002/16412 which are herein incorporated by reference in their entireties).

EXAMPLE 4

Preparation of Neutrokin- α -DOTA Conjugate

[0475] The following materials are used in the preparation of the NA-chelator conjugate:

- Neutrokin- α protein (in citrate buffer (see below); concentration of about 2.4 mg/mL);
- MeO-DOTA-NCS;
- HEPES buffer (about 500 mM and about pH 8.5);

- 166 -

- citrate buffer (10 mM sodium citrate, 140 mM NaCl, pH 6.0);
- 1 M Glycine-HCl (Glycine buffer);
- 1 N NaOH solution;
- water for injection (WFI);
- diafiltration buffer (10 mM sodium acetate, 140 mM NaCl, pH 6.0);
- Millipore Pellicon XL regenerated cellulose membrane - 10 kD molecular weight cut off (MWCO).
- Reaction vessels for preparing Neutrokin- α -DOTA conjugates and Neutrokin- α -DOTA complexes of the invention are preferably disposable plastic containers to minimize contamination with heavy metal ions

[0476] Into a sterile container (reaction vessel) was placed 708.3 mL of Neutrokin- α protein solution (concentration about 2.4 mg/mL in citrate buffer). The Neutrokin- α protein is a homotrimer of three molecules of Neutrokin- α each having a sequence of amino acids 134-285 of SEQ ID NO:2. Citrate buffer, 311.2 mL, was added into the sterile container and gently mixed. 60 μ L of the diluted Neutrokin- α protein was mixed with 40 μ L of citrate buffer and set aside in a 1.5 mL Eppendorf tube labeled "Control". Freeze down sample or keep it on ice for a short period of time and then freeze.

[0477] About 153 mL of HEPES buffer were added to the reaction vessel and gently swirled to mix.

[0478] The MeO-DOTA-NCS solution is prepared in a separate container. MeO-DOTA-NCS, 11.9 g, was placed into a separate container to which 170 mL of water for Injection and 34 mL 1N NaOH solution were added. The container was mixed by inversion until all MeO-DOTA-NCS was dissolved.

[0479] The quantity MeO-DOTA-NCS used is established by the approximate 1:11 molar ratio of chelator bonding sites in Neutrokin- α to MeO-DOTA-NCS molecules optimal for giving one MeO-DOTA moiety, on average, per Neutrokin- α monomer (or 3 MeO-DOTA moieties per Neutrokin- α

- 167 -

trimer). In this case, the Neutrokin- α protein contains 11 chelator bonding sites (Ala-134, Lys-160, Lys-173, Lys-181, Lys-184, Lys-188, Lys-204, Lys-215, Lys-216, Lys-252, and Lys-283).

[0480] The MeO-DOTA-NCS solution was then added to the reaction vessel containing the Neutrokin- α protein. The reaction vessel was then gently mixed. The container that contained the MeO-DOTA-NCS solution was then rinsed with about 153 mL of water for injection, which was then added to the reaction vessel.

[0481] If pH of the solution in the reaction vessel was not at pH 8.5 ± 0.1 , the pH was adjusted with 1 M NaOH or 1 M citric acid as necessary. The volume of titration necessary to adjust pH was recorded.

[0482] The reaction vessel was placed into a water bath and was gently agitated for 4.5 to 5.5 hours at about 25 °C. At the end of incubation time, 68 mL of glycine buffer was added to the reaction vessel to stop the reaction. The reaction was incubated for approximately an additional 15 minutes at about 25 °C. The volume of the reaction solution was adjusted to about 1700 mL by addition of citrate buffer (10mM sodium citrate, 140mM sodium chloride, pH 6.0). One sample (0.5 mL) of the reaction solution was obtained and labeled as "prediafiltration sample." The prediafiltration sample was lyophilized or kept on ice.

EXAMPLE 5

Diafiltration Procedure

[0483] One 0.1 m² Pellicon XL™ regenerate cellulose membrane cartridge (Millipore) was prepared according to manufacturer's instruction manual included with the cartridge. The flow rate of the peristaltic pump is set to about 400 mL/min. The membrane was sanitized with 0.1 N NaOH for at least about 30 minutes. At least about 1.0 L of diafiltration buffer was flushed through the retentate port, and at least about 1.0 L diafiltration buffer was

- 168 -

flushed through the permeate port to equilibrate system. Neutrokin- α conjugate was concentrated approximately 2.5-fold.

[0484] Maintaining the volume within the reaction volume, ten diafiltration volumes (DV) of buffer were exchanged.

[0485] A sample of the solution, about 0.5 mL, was obtained and labeled as "post diafiltration (DF)" sample. The post DF sample was lyophilized or kept on ice. The membrane was cleaned according to manufacturer's instructions.

[0486] Samples were analyzed by reverse phase HPLC for purity and by mass spectrometry to estimate the amount of chelator molecules covalently linked to each Neutrokin- α monomer. A Bradford assay was used to determine the Neutrokin- α conjugate concentration.

EXAMPLE 6

Incorporation Of ^{90}Y Into Neutrokin- α Conjugate

[0487] The Neutrokin- α conjugate, as prepared in Example 5, was used to prepare a Neutrokin- α complex comprising ^{90}Y . The ratio of chelator to protein in the conjugate was about 1.13. 50 μL of Neutrokin- α conjugate (1 mg/mL in an acetate buffer (0.1 M, pH 6.0)) was placed in a vial. About 2 μL of the following yttrium solution was added: yttrium solution consisted essentially of 19 μL of 15 mM Y^{3+} in 1 mM HCl and 1 μL (10 μCi) of $^{90}\text{Y}^{3+}$ in 1 mM HCl. The reaction solution was mixed and allowed to stand for about 30 minutes at room temperature (about 23 $^{\circ}\text{C}$). 2 μL of 23.6 mM DPTA was added. Reaction mixture was allowed to stand for about 5 minutes.

[0488] Chelation efficiency may be determined using an instantaneous thin layer chromatography (ITLC) kit which is commercially available from BioDex Medical, Shirley, New York (Tec-control Kit #151-770). In ITLC, radionuclide not associated with protein will migrate with the solvent front whereas radionuclide associated with a Neutrokin- α complex will migrate more slowly. Comparison of the amount of the radioactivity in the

- 169 -

dye front compared to the amount of radioactivity in the portions of the ITLC strip closer to the sample will give the chelation efficiency. Briefly, the final drug product is diluted 1 to 10 in saline (0.9%). 2 microliters of the diluted product is deposited on the ITLC strip. The strip is developed in saline (1 milliliter). The strip is dried and cut in two halves; each half is deposited in a glass tube. The free metal is measured by the Gamma Well counter (Packard Instruments Model 5003) and reported as a percentage of the total activity.

[0489] The above procedure was repeated varying the time of the chelating reaction (0, 5, 10, 20, and 30 minutes before adding EDTA or DPTA) and varying the concentration of the yttrium (2x concentration).

EXAMPLE 7

Preparation Of Neutrokin-Alpha Complex For Therapeutic Use

[0490] A Neutrokin- α complex is prepared by reacting a Neutrokin- α conjugate with $^{90}\text{YCl}_3$. 1.0 (2mg/ml) mL of a first solution comprising the Neutrokin- α conjugate as prepared in Example 5 (2 mg/mL) is added to a reaction vial containing 1.0 mL of acetate buffer (275 mM, pH8.4). 1.0 mL of a solution of $^{90}\text{YCl}_3$ (60 mCi/mL) is added to the reaction vial. The reaction vial is allowed to stand or gently mixed or agitated for up to 45, preferably 30-45 minutes, more preferably 30 minutes. After the given amount of time, 7.0 mL of a solution comprising an acetate buffer (10 mM), NaCl (140 mM) ascorbate (4.0%) and DTPA (1 mM) at pH 6.0 is added to the reaction vial. The labeling reaction can be optimized using routine techniques that are known in the art. For example, by varying the pH, reactant concentrations (neutrokin- α conjugate, metal ion), reaction vessel material, and temperature, percent incorporation of the metal ion into a complex can be maximized and reaction time minimized.

^{90}Y labeling Protocol

[0491] In an 10cc vial from West (unwashed) combine:

- 170 -

- 1.8 milliliters of sodium acetate (NaOAc, 275 mM) with 0.4 milliliters of sodium bicarbonate (NaHCO₃, 377mM).
- 70 mCi of sterile ⁹⁰Y (MDS Nordion, Ontario, Canada)
- 1 milliliter of Neutrokine-alpha conjugate as prepared in Examples 4-5 (2 mg/mL).

[0492] Mix the vial by gentle inversion and react for 45 minutes at room temperature. Stop the reaction by adding diluent (140 mM NaCl, 2 mM DPTA, and 10% sodium ascorbate) to a final volume of 9.95 milliliters. Let the mixture stand for five minutes prior to proceeding further, i.e. prior to ITLC analysis, HPLC analysis (see Example 12) confirmation that pH is in the range of 6.0 to 7.5, testing in a receptor binding assay, for example as described in Example 11. Neutrokine-alpha complex prepared as described above was shown to be stable for 8 hours at room temperature. Storage of the Neutrokine-alpha complex at 2-8 °C would be expected to extend its shelf life.

[0493] Optimally, therapeutic drug product will meet or exceed the following specifications: 10% (or less) free ⁹⁰Y as determined by ITLC, 10% (or less) protein aggregates relative to trimer as determined by HPLC (see Example 12), pH 6.5-7.5, with at least 75% specific binding in a receptor binding assay (see polystyrene bead assay in Example 11).

¹¹¹In labeling Protocol

[0494] In a 10 cc vial from West (pressure washed with water for injection) combine:

- 1.8 milliliters of sodium acetate (NaOAc, 275 mM) with 0.4 milliliters of sodium bicarbonate (NaHCO₃, 377mM).
- 10 mCi of sterile ¹¹¹In (MDS Nordion, Ontario, Canada) in 0.08 mM InCl₃
- 1 milliliter of Neutrokine-alpha conjugate as prepared in Examples 4-5 (2 mg/mL).

[0495] Mix the vial by gentle inversion and react for 45 minutes at room temperature. Stop the reaction by adding diluent (140mM NaCl, 2mM DPTA,

- 171 -

and 10% sodium ascorbate) to a final volume of 9.95 milliliters. Let the mixture stand for five minutes prior to proceeding further, i.e. prior to ITLC analysis, HPLC analysis (see Example 12) confirmation that pH is in the range of 6.0 to 7.5, testing in a receptor binding assay, for example as described in Example 11. Neutrokin-alpha complex prepared as described above was shown to be stable for 8 hours at room temperature. Storage of the Neutrokin-alpha complex at 2-8 °C would be expected to extend its shelf life.

[0496] In a preferred embodiment, therapeutic drug product will meet or exceed the following specifications: 10% (or less) free ¹¹¹In as determined by ITLC, 10% (or less) protein aggregates relative to trimer as determined by HPLC (see Example 12), pH 6.5-7.5, with at least 75% specific binding in a receptor binding assay (e.g., see polystyrene bead assay in Example 11).

[0497] Reagents for the labeling reaction may be provided in kit form. The kit would contain one or more of the following: (a) a vial pre-filled with reaction buffer (e.g., 275 mM sodium acetate, 377 mM sodium bicarbonate in a ratio of 4:1, respectively), (b) a vial pre-filled with diluent (e.g., 140mM NaCl, 2mM DPTA, and 10% sodium ascorbate), (c) a vial containing Neutrokin-alpha conjugate (e.g., 2 milligrams per milliliter), (d) an empty reaction vial and (e) a vial containing the radionuclide.

[0498] In a specific embodiment, a kit of the invention contains vials (a), (b) and (c). In this case, where no empty reaction vial is included in the kit, one may be provided separately; alternatively, the vial containing the Neutrokin-alpha complex or the reaction buffer may be used as the reaction vial. The reaction vial should be large enough to contain the full volume of the final drug product, (e.g., at least 10cc in this example).

EXAMPLE 8

Spect Imaging

[0499] The test dosage of radioactivity for these studies is 10 mCi of a Neutrokin-alpha complex prepared according to Example 6 or 7. The

- 172 -

preparation of the injection material (*i.e.*, sterilization, final calibration of dosage and loading the syringe) is done in a radiopharmacy. A Rhesus monkey is anesthetized by inhalation Metophane, and an intravenous catheter implanted for injection of the Neutrokin- α complex and transported to the SPECT suite. The animal is maintained under anesthesia for the duration of the procedure. The monkey's head is placed in a custom-designed animal-sized collimator for the scan and data is obtained continuously for 120 minutes following an intravenous bolus of the complex. The binding sites of Neutrokin- α are visualized using the SPECT data. This test is performed at approximately the peak of receptor occupancy. Following the completion of the scan the monkey is kept in a containment facility until the radioactive material is clear from the system.

[0500] Alternatively, a full body scan is performed to determine total distribution of the Neutrokin- α complex.

EXAMPLE 9

Treatment Of Lung Cancer Patient

[0501] A human patient having small-cell carcinoma of the right lung is infused intravenously with a sterile, pyrogen-free solution containing 4 mg of the Neutrokin- α complex in sterile, buffered saline, prepared as described herein, for example according to Example 6 or 7 hereof. After 5 days, the conjugate is well localized in the lung and has substantially cleared from the circulation of the patient, as seen by imaging scanning at daily intervals.

EXAMPLE 10

Therapy Of Lymphoma

[0502] A human patient suffering from lymphoma is infused intravenously with a sterile, pyrogen-free solution containing 4 mg of the Neutrokin- α complex in sterile, buffered saline, prepared as described herein, for example

- 173 -

according to Example 6 or 7 hereof. After 6 days, the conjugate is well localized at the target and has substantially cleared from the circulation of the patient, as seen by imaging scanning at daily intervals.

EXAMPLE 11

Neutrokin- α Receptor or Neutrokin- α Antibody Binding Assay

[0503] An in vitro direct binding assay is used to assess the ability of the radiolabeled Neutrokin- α (*e.g.*, Neutrokin- α complex) protein to bind to one of its known cellular receptors, *e.g.*, TACI, BCMA or BAFF-R, or to an anti-Neutrokin- α antibody (*e.g.*, a Neutrokin- α antibody that neutralizes the ability of Neutrokin- α to bind to one of its cellular receptors). Exemplary antibodies that could be used in this assay are described in International Patent Application Publication Number WO03/55979, WO03/02641 (in particular, antibodies produced by the cell lines deposited in association with WO02/02641 and described on page 145 of WO02/02641) and WO03/33658 which are hereby incorporated by reference in their entirety. Purified Neutrokin- α receptor or anti-Neutrokin- α antibody is coated onto polystyrene beads, which are suspended in solution with an aliquot of radiolabeled protein. The coated beads and ligand are allowed to incubate for 30 minutes, at room temperature, with shaking on an Eppendorf thermomixer at 1400 rpm. The beads are then pelleted by centrifugation and frozen in liquid nitrogen. The tube is sectioned to separate the liquid phase from the pellets and both are counted on a gamma-well counter. Non-specific binding is assessed in parallel with beads that do not have Neutrokin- α receptor or anti-Neutrokin- α antibody bound to them. Percent specific binding is calculated as: $[(\text{total radioactivity bound to coated beads} - \text{radioactivity bound to uncoated beads}) / (\text{total radioactivity bound to coated beads})] \times 100\%$.

- 174 -

[0504] Alternatively, live cells that express Neutrokin- α receptors such as IM-9 cells may be used to assess the ability of the radiolabeled Neutrokin- α protein (*e.g.*, Neutrokin- α complex) to bind to one of its cellular receptors. In such an assay, the Neutrokin- α complex is incubated with the cells for a period sufficient to allow binding of Neutrokin- α complex with Neutrokin- α receptors. The cells are washed to remove unbound or non-specifically bound Neutrokin- α complex. Cell-associated radioactivity is then measured.

EXAMPLE 12

HPLC Analysis of final drug product

[0505] The Neutrokin- α complex for therapeutic use may be analyzed for radiochemical purity aggregates and identity using an HPLC method with a size exclusion column. A sample of undiluted Neutrokin- α complex (as prepared in Example 6 or 7, for example) is loaded onto the size exclusion column. 10mM sodium acetate, with 140 mM NaCl is used as the mobile phase. The concentration of Neutrokin- α trimer in the final drug product may be determined by comparison of the results to a calibration curve for 0.15, 0.20, 0.25, and 0.30 milligram/milliliter of Neutrokin- α trimer standards. Radiochemical purity and identity can be established by comparing the test sample HPLC profile with the standard HPLC profile for trimeric Neutrokin- α protein. Identity can be established by HPLC retention time. Radiochemical purity can be established by the percentage labeled protein trimer versus the total labeled protein trimer and aggregates.

[0506] Each of the preceding examples can be repeated by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

[0507] Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without

- 175 -

affecting the scope of the invention or any embodiment thereof. All patents and publications cited herein are fully incorporated by reference herein in their entirety.

}

- 176 -

WHAT IS CLAIMED IS:

1. A Neutrokin- α conjugate having the formula $NA-(Chel)_n$,
wherein

NA is a Neutrokin- α protein;

Chel is said chelator; and

n is an integer from 1 to about 30;

wherein said Neutrokin- α protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of amino acid residues 134 to 285 of SEQ ID NO:2;

(b) the amino acid sequence of amino acid residues n to 285 of SEQ ID NO:2, where n is an integer in the range of 2-190;

(c) the amino acid sequence of amino acid residues 1 to m of SEQ ID NO:2, where m is an integer in the range of 274-284;

(d) the amino acid sequence of amino acid residues n to m of SEQ ID NO:2, where n is an integer in the range of 2-190 and m is an integer in the range of 274-284; and

(e) an amino acid sequence which has at least 80% identity to any of the proteins described in (a), (b), (c), and (d);

wherein said Neutrokin- α protein binds a Neutrokin- α receptor.

2. The conjugate according to claim 1, wherein said Neutrokin- α protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of amino acid residues n to 285 of SEQ ID NO:2, where n is an integer in the range of 2-190;

(b) the amino acid sequence of amino acid residues 1 to m of SEQ ID NO:2, where m is an integer in the range of 274-284; and

- 177 -

(c) the amino acid sequence of amino acid residues n to m of SEQ ID NO:2, where n is an integer in the range of 2-190 and m is an integer in the range of 274-284;


wherein said Neutrokin- α protein binds a Neutrokin- α receptor.

3. The conjugate of claim 1, wherein n is 1, 2, 3, 4, 5, or 6.
4. The conjugate according to claim 3, wherein n is 3.
5. The conjugate according to claim 3, wherein n is 1.
6. The conjugate according to claim 4, wherein said Neutrokin- α protein is a mature, soluble Neutrokin- α protein.
7. The conjugate according to claim 6, wherein said protein comprises a sequence that is at least 85% identical to amino acids 134-285 of SEQ ID NO:2.
8. The conjugate according to claim 7, wherein said Neutrokin- α consists of a trimer of Neutrokin- α monomeric subunits and wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2.
9. The conjugate according to claim 8, wherein at least one or more lysine residues or N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.
10. The conjugate according to claim 9, wherein at least one or more N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

- 178 -

11. The conjugate according to claim 1, wherein said chelator is DOTA, a DOTA derivative, or a DOTA analogue, optionally containing a linker moiety.

12. The conjugate according to claim 11, having the formula



wherein

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen;

each R^5 independently is hydrogen, $\text{C}_1\text{-C}_4$ alkyl or $-(\text{C}_1\text{-C}_2\text{ alkyl})\text{phenyl}$;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

n' is 1 to 10;

m is an integer from 0 to 10 inclusive;

p is 1 or 2;

r is 0 or 1;

w is 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

- 179 -


R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

R^3 is selected from the group consisting of C_1 - C_4 alkoxy, $-OCH_2CO_2H$, hydroxy and hydrogen;

R^6 is a group formed from a chemical group selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido, and carboxyl;

with the proviso that R^2 and R^6 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

13. The conjugate according to claim 12, having the formula



or a pharmaceutically acceptable salt thereof, wherein n^1 is 1 to 10.

14. The conjugate according to claim 11, wherein NA is a human, mature, soluble Neutrokin- α protein.

15. The conjugate according to claim 14, wherein said NA has a sequence that is at least 85% identical to amino acids 134-285 of SEQ ID NO:2.

- 180 -


16. The conjugate according to claim 15, wherein said Neutrokin- α consists of a trimer of Neutrokin- α monomeric subunits and wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2.

17. The conjugate according to claim 16, wherein at least one or more lysine residues or N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with $(\text{Chel})_n$.

18. The conjugate according to claim 17, wherein at least one or more N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with $(\text{Chel})_n$.

19. The conjugate according to claim 18, wherein n is 1.

20. A Neutrokin- α conjugate having the formula:



or a

pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokin- α protein that consists of a trimer of Neutrokin- α monomeric subunits wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1, 2, 3, 4, 5 or 6;

N' is a nitrogen from the amino terminus or from a lysine residue of said Neutrokin- α protein.

- 181 -

21. A Neutrokin- α complex comprising a Neutrokin- α conjugate and a metal ion wherein said metal ion is associated with the chelator moiety of said Neutrokin- α conjugate.

22. The complex of claim 21, wherein said Neutrokin- α conjugate has the formula $NA-(Chel)_n$, wherein

NA is a Neutrokin- α protein;

Chel is said chelator; and

n is an integer from 1 to about 10;

wherein said Neutrokin- α protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of amino acid residues n to 285 of SEQ ID NO:2, where n is an integer in the range of 2-190;

(b) the amino acid sequence of amino acid residues 1 to m of SEQ ID NO:2, where m is an integer in the range of 274-284;

(c) the amino acid sequence of amino acid residues n to m of SEQ ID NO:2, where n is an integer in the range of 2-190 and m is an integer in the range of 274-284; and

(d) an amino acid sequence which has at least 80% identity to any of the proteins described in (a), (b), and (c);

wherein said Neutrokin- α protein binds a Neutrokin- α receptor.

23. The complex according to claim 22 wherein said Neutrokin- α protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of amino acid residues n to 285 of SEQ ID NO:2, where n is an integer in the range of 2-190;

(b) the amino acid sequence of amino acid residues 1 to m of SEQ ID NO:2, where m is an integer in the range of 274-284; and

- 182 -

(c) the amino acid sequence of amino acid residues n to m of SEQ ID NO:2, where n is an integer in the range of 2-190 and m is an integer in the range of 274-284;

wherein said Neutrokin- α protein binds a Neutrokin- α receptor.

24. The complex of claim 22, wherein n is 1, 2, 3, 4, 5, or 6.

25. The complex according to claim 24, wherein n is 3.

26. The complex according to claim 25, wherein n is 1.

27. The complex according to claim 25, wherein said Neutrokin- α protein is a mature, soluble Neutrokin- α protein.

28. The complex according to claim 27, wherein said Neutrokin- α protein has a sequence that is at least 85% identical to amino acids 134-285 of SEQ ID NO:2.

29. The complex according to claim 28, wherein said Neutrokin- α consists of amino acids 134-285 of SEQ ID NO:2.


30. The complex according to claim 29, wherein at least one or more lysine residues or N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

31. The complex according to claim 30, wherein at least one or more N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

32. The complex according to claim 22, wherein the chelator is DOTA, a DOTA derivative, or a DOTA analog, optionally containing a linker moiety.

- 183 -

33. The complex according to claim 32, wherein the conjugate has the formula



wherein

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;

with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen;

each R^5 independently is hydrogen, $\text{C}_1\text{-C}_4$ alkyl or $-(\text{C}_1\text{-C}_2\text{ alkyl})\text{phenyl}$;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1;

n' is 1 to 10;

m is an integer from 0 to 10 inclusive;

p is 1 or 2;

r is 0 or 1;

w is 0 or 1;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;


R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl;

- 184 -

R^3 is selected from the group consisting of C_1 - C_4 alkoxy, $-OCH_2CO_2H$, hydroxy and hydrogen;

R^6 is a functional group formed from a group selected from the group consisting of nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl; with the proviso that R^2 and R^6 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

34. The complex according to claim 33, wherein the conjugate has the formula



or a pharmaceutically acceptable salt thereof, wherein n^1 is 1 to 10.

35. The complex according to claim 34, wherein NA is a human, mature, soluble Neutrokin- α protein.

36. The complex according to claim 35, wherein said NA has a sequence that is at least 85% identical to amino acids 134-285 of SEQ ID NO:2.


37. The complex according to claim 36, wherein said Neutrokin- α consists of a trimer of Neutrokin- α monomeric subunits and wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2.

- 185 -

38. The complex according to claim 37, wherein at least one or more lysine residues or N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

39. The complex according to claim 38, wherein at least one or more N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

40. A Neutrokin- α complex comprising a Neutrokin- α conjugate and a metal ion wherein said metal ion is associated with the chelator moiety of said Neutrokin- α conjugate, wherein said conjugate has the formula



or a pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokin- α protein that consists of a trimer of Neutrokin- α monomeric subunits wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1, 2, 3, 4, 5 or 6;

N' is a nitrogen from the amino terminus or from a lysine residue of said Neutrokin- α protein.

41. The complex according to claim 40, wherein said metal ion is selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac .

- 186 -

42. The complex according to claim 41, wherein said metal ion is ^{90}Y .
43. The complex according to claim 41, wherein said metal ion is ^{111}In .
44. The complex according to claim 41, wherein said metal ion is ^{177}Lu .
45. The complex according to claim 41, wherein said metal ion is ^{166}Ho .
46. The complex according to claim 41, wherein said metal ion is ^{215}Bi .
47. The complex according to claim 41, wherein said metal ion is ^{225}Ac .
48. A composition comprising a Neutrokin- α conjugate according to any of claims 1-21 and a suitable carrier.
49. A composition comprising a Neutrokin- α conjugate according to claim 20 and a pharmaceutically acceptable carrier.
50. The composition according to claim 49, wherein said carrier is sterile water.
51. The composition according to claim 50 further comprising a buffer.
52. The composition according to claim 51 wherein said buffer is an acetate buffer having a concentration of from about 10 mM to about 200 mM.

- 187 -

53. The composition according to claim 49 further comprising a metal ion selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac .

54. A composition comprising a Neutrokin- α complex according to any of claims 22-47 and a suitable carrier.

55. A composition comprising a Neutrokin- α complex according to claim 40 and a pharmaceutically acceptable carrier.

56. The composition according to claim 55 wherein said carrier is sterile water.

57. The composition according to claim 56 further comprising a buffer.

58. The composition according to claim 57 wherein said buffer is an acetate buffer having a concentration of from about 10 mM to about 200 mM.


59. A method of preparing the conjugate of claim 1 comprising reacting a Neutrokin- α protein with a chelator.

60. The method according to claim 59, wherein said chelator is an activated chelator.

61. The method according to claim 60, wherein said activated chelator is a DOTA derivative or a DOTA analogue.

62. The method according to claim 61, wherein said chelator has the formula activated chelator having the formula:

- 188 -



wherein

each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$, preferably $-\text{CH}_2\text{CO}_2\text{R}$, more preferably $-\text{CH}_2\text{CO}_2\text{H}$;

Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$, preferably CO_2R , more preferably COOH ;

each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl; preferably hydrogen or $\text{C}_1\text{-C}_4$ alkyl, more preferably hydrogen;

with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen;

each R^5 independently is hydrogen, $\text{C}_1\text{-C}_4$ alkyl or $-(\text{C}_1\text{-C}_2\text{ alkyl})\text{phenyl}$;

X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;

n is 0 or 1; preferably 0

m is an integer from 0 to 10 inclusive, preferably 0 to 1, more preferably 0;

p is 1 or 2, preferably 1;

r is 0 or 1, preferably 0;

w is 0 or 1, preferably 0;

with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato;

R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{CO}_2\text{H}$, hydroxy and hydrogen, preferably $\text{C}_1\text{-C}_4$ alkoxy, more preferably methoxy;

- 189 -

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, maleimido, bromoacetamido and carboxyl, preferably hydrogen or isothiocyanato; with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

63. The method according to claim 62, wherein said activated chelator is α -(5-isothiocyanato-2-methoxyphenyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.

64. The method according to claim 63, wherein the Neutrokin- α protein is a human, mature, soluble Neutrokin- α protein.

65. The method according to claim 64, wherein the Neutrokin- α protein has a sequence that is at least 85% identical to amino acids 134-285 of SEQ ID NO:2.

66. The method according to claim 65, wherein said Neutrokin- α consists of a trimer of Neutrokin- α monomeric subunits and wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2.

67. The method according to claim 66, wherein at least one or more lysine residues or N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with the chelator.

68. The conjugate according to claim 67, wherein at least one or more N-terminal alanine residues of said Neutrokin- α protein forms the covalent bond with said chelator.

69. The method according to claim 68, comprising mixing, agitating, or preparing a solution comprising said Neutrokin- α protein and said chelator at a temperature of about 0 °C to about 50 °C for about 0.5

- 190 -

hours to about 10 hours, wherein said solution has a pH of about 8.0 to about 9.0.

70. The method according to claim 69, wherein said mixing, agitating, or preparing occurs at a temperature of about 20 °C to about 30 °C for about 3 hours to about 5 hours.

71. The method according to claim 70, further comprising adding a quenching agent after from about 3 hours to about 5 hours.

72. The method according to claim 71, wherein said pH is about 8.5.

73. The method according to claim 72, wherein the molar ratio of chelator to chelator bonding sites in the Neutrokin- α protein is from about 10:1 to about 12:1.

74. The method according to claim 73, wherein said quenching agent is glycine or glycine hydrochloride.

75. The method according to claim 74, wherein said solution further comprises citrate buffer and HEPES.

76. The method according to claim 75, wherein the concentration of said Neutrokin- α protein is from about 1.5 to about 3.0 mg/mL, and wherein the molar ratio of said chelator to chelator bonding sites in the Neutrokin- α protein is from about 10:1 to about 12:1.

77. The method according to claim 76, further comprising purifying said Neutrokin- α conjugate.

78. The method according to claim 77, wherein said purifying comprises a diafiltration method.

- 191 -

79. A method of preparing a Neutrokin complex according to claim 22, comprising mixing, agitating, or preparing a solution comprising a Neutrokin- α conjugate and a metal ion capable of complexing with a Neutrokin- α conjugate;

wherein said Neutrokin- α conjugate having the formula $NA-(Chel)_n$, wherein

(a) NA is a Neutrokin- α protein;

(b) Chel is said chelator; and

(c) n is an integer from 1 to about 10;

wherein said Neutrokin- α protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of amino acid residues n to 285 of SEQ ID NO:2, where n is an integer in the range of 2-190;

(b) the amino acid sequence of amino acid residues 1 to m of SEQ ID NO:2, where m is an integer in the range of 274-284;

(c) the amino acid sequence of amino acid residues n to m of SEQ ID NO:2, where n is an integer in the range of 2-190 and m is an integer in the range of 274-284; and


(d) an amino acid sequence which has at least 80% identity to any of the proteins described in (a), (b), and (c);

wherein said Neutrokin- α protein binds a Neutrokin- α receptor

80. A method of preparing the complex according to claim 40, comprising mixing, agitating, or preparing a solution comprising a Neutrokin- α conjugate and a metal ion capable of associating with a Neutrokin- α conjugate;

wherein said conjugate has the formula

- 192 -



or a pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokin- α protein that consists of a trimer of Neutrokin- α monomeric subunits and wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1;

N' is a nitrogen from the amino terminus or from a lysine residue of said Neutrokin- α protein.

81. The method according to claim 80, further comprising removing excess metal ion.

82. The method according to claim 81, wherein said removing comprises adding a chelating agent selected from the group consisting of DPTA, EDTA, and MeO-DOTA-glycine.

83. The method according to claim 80, wherein said solution further comprises an acetate buffer having a concentration of from about 1 mM to about 20 mM and NaCl having a concentration of about 100 mM to about 200 mM.

84. The method according to claim 80, wherein the metal ion is selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac .

- 193 -

85. The method according to claim 81, wherein the solution further comprises an acetate buffer having a concentration of from about 1 mM to about 20 mM; and

further comprising allowing the solution to mix, agitate, or stand for from about 5 minutes to about 60 minutes at a temperature from about 20 °C to about 30 °C; and

adding a second solution, said second solution comprising

an acetate buffer having a concentration of about 10 mM,

NaCl having a concentration of about 140 mM,

HSA having a concentration of about 7.5 %, and

DPTA having a concentration of about 1 mM, wherein said second solution has a pH of about 6.

86. A method of administering radiotherapy to a subject in need thereof, comprising administering to said subject an effective amount of a Neutrokin- α complex according to claim 22.

87. A method of administering radiotherapy to a subject in need thereof, comprising administering to said subject an effective amount of a Neutrokin- α complex according to claim 40.

88. The method of claim 87, wherein said Neutrokin- α complex is administered as an injectable solution.

89. The method according to claim 88, wherein said solution is administered intravenously.

90. The method according to claim 87, wherein a dosage of radioactivity from about 5 mCi to about 200 mCi is administered.

91. The method of claim 90, wherein said subject is a human.

- 194 -

92. The method of claim 90, wherein said subject has a B-cell mediated disease.

93. The method of claim 90, wherein said subject has a condition selected from the group consisting of non-Hodgkin's lymphoma, chronic lymphocytic leukemia, multiple myeloma, systemic lupus erythematosus, rheumatoid arthritis, multiple sclerosis, Crohn's disease, diabetes, Wegener's granulomatous, myasthenia gravis, and asthma.

94. The method of claim 93, wherein said subject has non-Hodgkin's lymphoma.

95. A method of treating cancer comprising administering to a subject with cancer, an effective amount of a Neutrokin-alpha complex according to claim 40.

96. The method of claim 95 wherein a cell of said cancer expresses a Neutrokin-alpha receptor on its surface.

97. The method of claim 95 wherein said cancer is a B cell cancer.

98. The method of claim 97 wherein said B cell cancer is selected from the group consisting of:

- (a) Non-Hodgkin's Lymphoma;
- (b) Multiple Myeloma; and
- (c) Chronic Lymphocytic Leukemia.

99. A method of treating an autoimmune disease or disorder comprising administering to a subject with an autoimmune disease or disorder, an effective amount of a Neutrokin-alpha complex according to claim 40.

- 195 -

100. The method of claim 99 wherein said autoimmune disease or disorder is selected from the group consisting of:

- (a) Systemic Lupus Erythematosus;
- (b) Rheumatoid Arthritis; and
- (c) Sjögren's Syndrome.

101. A method of killing a cell selected from the group consisting of:

- (a) a cell bearing a Neutrokin- α receptor; and
- (b) a cell in close proximity to a cell bearing Neutrokin- α receptors;

wherein said method comprises contacting said cell with a composition according to claim 22 in an amount effective to kill a said cell.

102. The method of claim 101 wherein said cell is (a).

103. The method of claim 101 wherein said cell is (b).

104. The method of claim 101 wherein said cell is a lymphocyte.

105. The method of claim 104 wherein said cell is B cell.

106. The method of claim 101 wherein said cell is cancerous cell that has metastasized into the lymphatic system.


107. A method of diagnostic imaging, comprising administering a Neutrokin- α complex according to any of claims 22-47.

108. A kit comprising a first vial containing a Neutrokin- α conjugate.

109. The kit according to claim 108, wherein

- 196 -

said first vial contains a Neutrokin- α conjugate in a solution of acetate buffer (10 mM sodium acetate, 140 mM sodium chloride, pH 6.0) said conjugate having the formula



or a pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokin- α protein that consists of a trimer of Neutrokin- α monomeric subunits wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1, 2, 3, 4, 5 or 6; and

N^1 is a nitrogen from the amino terminus or from a lysine residue of said Neutrokin- α protein.

110. The kit of claim 108 comprising


- (a) a second vial containing a radionuclide; and
- (b) a third vial containing a buffer solution.

111. The kit according to claim 110 which further comprises a fourth vial wherein said vial is empty.

112. The kit according to claim 110, wherein

said first vial contains a Neutrokin- α conjugate in a solution of acetate buffer (10 mM sodium acetate, 140 mM sodium chloride, pH 6.0) said conjugate having the formula

- 197 -



or a pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokine-alpha protein that consists of a trimer of Neutrokine-alpha monomeric subunits wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1, 2, 3, 4, 5 or 6; and

N' is a nitrogen from the amino terminus or from a lysine residue of said Neutrokine-alpha protein;

said second vial contains a radionuclide selected from the group consisting of ^{90}Y , ^{111}In , ^{177}Lu , ^{166}Ho , ^{215}Bi , and ^{225}Ac ; and

said third vial contains a buffer solution, wherein said buffer solution is an acetate buffer solution having a concentration in the range of about 50 mM to about 300mM.

113. The kit of claim 108 comprising

- (c) a second vial containing buffer solution; and
- (d) a third vial containing a diluent.


114. The kit according to claim 113 which further comprises a fourth vial wherein said vial is empty.

115. The kit according to claim 113, wherein

said first vial contains a Neutrokine-alpha conjugate in a solution comprising acetate buffer having a concentration of from about 1 mM

- 198 -

to about 20 mM and NaCl having a concentration of about 100 mM to about 200 mM and having a pH of about 6, said conjugate having the formula



or a pharmaceutically acceptable salt thereof,

wherein NA is a Neutrokinine- α protein that consists of a trimer of Neutrokinine- α monomeric subunits wherein each subunit consists of amino acids 134-285 of SEQ ID NO:2;

n^1 is 1, 2, 3, 4, 5 or 6; and

N' is a nitrogen from the amino terminus or from a lysine residue of said Neutrokinine- α protein;

wherein said buffer solution in said second vial comprises sodium acetate in a concentration range of about 50 mM to about 300mM and sodium bicarbonate in a concentration range of about 50 mM to about 300mM; and

said diluent in said third vial comprises about 100mM to about 200 mM NaCl, about 1mM to about 10mM DPTA, and about from 7% to about 10% sodium ascorbate

116. The kit according to claim 115, wherein said solution in said first vial comprises 10 mM sodium acetate, 140 mM NaCl.

- 199 -

117. The kit according to claim 115, wherein said buffer solution in said second vial comprises 220 mM sodium acetate and 75 mM sodium bicarbonate.

118. The kit according to claim 115, wherein said diluent in said third vial comprises 140 mM NaCl, 2 mM DPTA and 10% sodium ascorbate.


119. The method according to claim 80, further comprising adding to the solution a second solution to form a mixture, said second solution comprising about 220 mM sodium acetate and about 75 mM sodium bicarbonate and allowing the mixture of the solution containing the Neutrokin- α conjugate, the metal ion, and the second solution to mix, agitate, or stand for about 5 minutes to about 60 minutes at a temperature from about 20°C to about 30°C.

120. The method according to claim 113, further comprising adding, after the mixture has mixed, agitated, or stood, a third solution comprising about 140 mM NaCl, about 2 mM DPTA, and about 10% sodium ascorbate.

121. A composition comprising a Neutrokin- α conjugate according to claim 20 and a suitable carrier.

122. A composition comprising a Neutrokin- α complex according to claim 40 and a suitable carrier.

123. A method of diagnostic imaging, comprising administering a Neutrokin- α complex according to claim 40.

Figure 1: Proposed labeling of BLYS-DOTA conjugate




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

-1-

SEQUENCE LISTING

<110> Human Genome Sciences, Inc.
 Parmelee, David
 Yeh, Ren-Hwa
 Galperina, Olga
 Hilbert, David
 Rosen, Craig A.

<120> Neutrokin- α Conjugate, Neutrokin- α Complex, and Uses Thereof

<130> 1488.181PC03

<150> US 60/435,262
 <151> 2002-12-23

<150> US 60/467,198
 <151> 2003-05-02

<160> 17

<170> PatentIn version 3.1

<210> 1
 <211> 1100
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (147) .. (1001)
 <223>

<400> 1
 aaattcagga taactctcct gaggggtgag ccaagccctg ccatgtagtg cacgcaggac 60
 atcaacaaac acagataaca ggaaatgata cattccctgt ggtaacttat tctaaaggcc 120
 ccaaccttca aagttcaagt agtgat atg gat gac tcc aca gaa agg gag cag 173
 Met Asp Asp Ser Thr Glu Arg Glu Gln
 1 5
 tca cgc ctt act tct tgc ctt aag aaa aga gaa gaa atg aaa ctg aag 221
 Ser Arg Leu Thr Ser Cys Leu Lys Lys Arg Glu Glu Met Lys Leu Lys
 10 15 20 25
 gag tgt gtt tcc atc ctc cca cgg aag gaa agc ccc tct gtc cga tcc 269
 Glu Cys Val Ser Ile Leu Pro Arg Lys Glu Ser Pro Ser Val Arg Ser
 30 35 40
 tcc aaa gac gga aag ctg ctg gct gca acc ttg ctg ctg gca ctg ctg 317
 Ser Lys Asp Gly Lys Leu Leu Ala Ala Thr Leu Leu Leu Ala Leu Leu
 45 50 55
 tct tgc tgc ctc acg gtg gtg tct ttc tac cag gtg gcc gcc ctg caa 365
 Ser Cys Cys Leu Thr Val Val Ser Phe Tyr Gln Val Ala Ala Leu Gln
 60 65 70
 ggg gac ctg gcc agc ctc cgg gca gag ctg cag ggc cac cac gcg gag 413
 Gly Asp Leu Ala Ser Leu Arg Ala Glu Leu Gln Gly His His Ala Glu
 75 80 85

-2-

aag ctg cca gca gga gca gga gcc ccc aag gcc ggc ctg gag gaa gct 461
Lys Leu Pro Ala Gly Ala Gly Ala Pro Lys Ala Gly Leu Glu Glu Ala
90 95 100 105

cca gct gtc acc gcg gga ctg aaa atc ttt gaa cca cca gct cca gga 509
Pro Ala Val Thr Ala Gly Leu Lys Ile Phe Glu Pro Pro Ala Pro Gly
110 115 120

gaa ggc aac tcc agt cag aac agc aga aat aag cgt gcc gtt cag ggt 557
Glu Gly Asn Ser Ser Gln Asn Ser Arg Asn Lys Arg Ala Val Gln Gly
125 130 135

cca gaa gaa aca gtc act caa gac tgc ttg caa ctg att gca gac agt 605
Pro Glu Glu Thr Val Thr Gln Asp Cys Leu Gln Leu Ile Ala Asp Ser
140 145 150

gaa aca cca act ata caa aaa gga tct tac aca ttt gtt cca tgg ctt 653
Glu Thr Pro Thr Ile Gln Lys Gly Ser Tyr Thr Phe Val Pro Trp Leu
155 160 165

ctc agc ttt aaa agg gga agt gcc cta gaa gaa aaa gag aat aaa ata 701
Leu Ser Phe Lys Arg Gly Ser Ala Leu Glu Glu Lys Glu Asn Lys Ile
170 175 180 185

ttg gtc aaa gaa act ggt tac ttt ttt ata tat ggt cag gtt tta tat 749
Leu Val Lys Glu Thr Gly Tyr Phe Phe Ile Tyr Gly Gln Val Leu Tyr
190 195 200

act gat aag acc tac gcc atg gga cat cta att cag agg aag aag gtc 797
Thr Asp Lys Thr Tyr Ala Met Gly His Leu Ile Gln Arg Lys Lys Val
205 210 215

cat gtc ttt ggg gat gaa ttg agt ctg gtg act ttg ttt cga tgt att 845
His Val Phe Gly Asp Glu Leu Ser Leu Val Thr Leu Phe Arg Cys Ile
220 225 230

caa aat atg cct gaa aca cta ccc aat aat tcc tgc tat tca gct ggc 893
Gln Asn Met Pro Glu Thr Leu Pro Asn Asn Ser Cys Tyr Ser Ala Gly
235 240 245

att gca aaa ctg gaa gaa gga gat gaa ctc caa ctt gca ata cca aga 941
Ile Ala Lys Leu Glu Glu Gly Asp Glu Leu Gln Leu Ala Ile Pro Arg
250 255 260 265

gaa aat gca caa ata tca ctg gat gga gat gtc aca ttt ttt ggt gca 989
Glu Asn Ala Gln Ile Ser Leu Asp Gly Asp Val Thr Phe Phe Gly Ala
270 275 280

ttg aaa ctg ctg tgacctactt acaccatgtc tgtagctatt ttcctccctt 1041
Leu Lys Leu Leu
285

tctctgtacc tctaagaaga aagaatctaa ctgaaaatac caaaaaaaaa aaaaaaaaa 1100

<210> 2
<211> 285
<212> PRT
<213> Homo sapiens

<400> 2

-3-

Met Asp Asp Ser Thr Glu Arg Glu Gln Ser Arg Leu Thr Ser Cys Leu
 1 5 10 15

Lys Lys Arg Glu Glu Met Lys Leu Lys Glu Cys Val Ser Ile Leu Pro
 20 25 30

Arg Lys Glu Ser Pro Ser Val Arg Ser Ser Lys Asp Gly Lys Leu Leu
 35 40 45

Ala Ala Thr Leu Leu Leu Ala Leu Leu Ser Cys Cys Leu Thr Val Val
 50 55 60

Ser Phe Tyr Gln Val Ala Ala Leu Gln Gly Asp Leu Ala Ser Leu Arg
 65 70 75 80

Ala Glu Leu Gln Gly His His Ala Glu Lys Leu Pro Ala Gly Ala Gly
 85 90 95

Ala Pro Lys Ala Gly Leu Glu Glu Ala Pro Ala Val Thr Ala Gly Leu
 100 105 110

Lys Ile Phe Glu Pro Pro Ala Pro Gly Glu Gly Asn Ser Ser Gln Asn
 115 120 125

Ser Arg Asn Lys Arg Ala Val Gln Gly Pro Glu Glu Thr Val Thr Gln
 130 135 140

Asp Cys Leu Gln Leu Ile Ala Asp Ser Glu Thr Pro Thr Ile Gln Lys
 145 150 155 160

Gly Ser Tyr Thr Phe Val Pro Trp Leu Leu Ser Phe Lys Arg Gly Ser
 165 170 175

Ala Leu Glu Glu Lys Glu Asn Lys Ile Leu Val Lys Glu Thr Gly Tyr
 180 185 190

Phe Phe Ile Tyr Gly Gln Val Leu Tyr Thr Asp Lys Thr Tyr Ala Met
 195 200 205

Gly His Leu Ile Gln Arg Lys Lys Val His Val Phe Gly Asp Glu Leu
 210 215 220

Ser Leu Val Thr Leu Phe Arg Cys Ile Gln Asn Met Pro Glu Thr Leu
 225 230 235 240

Pro Asn Asn Ser Cys Tyr Ser Ala Gly Ile Ala Lys Leu Glu Glu Gly
 245 250 255

-4-

Asp Glu Leu Gln Leu Ala Ile Pro Arg Glu Asn Ala Gln Ile Ser Leu
 260 265 270

Asp Gly Asp Val Thr Phe Phe Gly Ala Leu Lys Leu Leu
 275 280 285

<210> 3
 <211> 152
 <212> PRT
 <213> Homo sapiens

<400> 3

Ala Val Gln Gly Pro Glu Glu Thr Val Thr Gln Asp Cys Leu Gln Leu
 1 5 10 15

Ile Ala Asp Ser Glu Thr Pro Thr Ile Gln Lys Gly Ser Tyr Thr Phe
 20 25 30

Val Pro Trp Leu Leu Ser Phe Lys Arg Gly Ser Ala Leu Glu Glu Lys
 35 40 45

Glu Asn Lys Ile Leu Val Lys Glu Thr Gly Tyr Phe Phe Ile Tyr Gly
 50 55 60

Gln Val Leu Tyr Thr Asp Lys Thr Tyr Ala Met Gly His Leu Ile Gln
 65 70 75 80

Arg Lys Lys Val His Val Phe Gly Asp Glu Leu Ser Leu Val Thr Leu
 85 90 95

Phe Arg Cys Ile Gln Asn Met Pro Glu Thr Leu Pro Asn Asn Ser Cys
 100 105 110

Tyr Ser Ala Gly Ile Ala Lys Leu Glu Glu Gly Asp Glu Leu Gln Leu
 115 120 125

Ala Ile Pro Arg Glu Asn Ala Gln Ile Ser Leu Asp Gly Asp Val Thr
 130 135 140

Phe Phe Gly Ala Leu Lys Leu Leu
 145 150

<210> 4
 <211> 898
 <212> DNA
 <213> Homo sapiens

<220>

-5-

<221> CDS

<222> (6)..(560)

<223>

<400> 4

```

gcacc atg agg cga ggg ccc cgg agc ctg cgg ggc agg gac gcg cca gcc      50
      Met Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala
        1              5              10              15

ccc acg ccc tgc gtc ccg gcc gag tgc ttc gac ctg ctg gtc cgc cac      98
Pro Thr Pro Cys Val Pro Ala Glu Cys Phe Asp Leu Leu Val Arg His
          20              25              30

tgc gtg gcc tgc ggg ctc ctg cgc acg ccg cgg ccg aaa ccg gcc ggg      146
Cys Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala Gly
          35              40              45

gcc agc agc cct gcg ccc agg acg gcg ctg cag ccg cag gag tcg gtg      194
Ala Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu Ser Val
          50              55              60

ggc gcg ggg gcc ggc gag gcg gcg ctg ccc ctg ccc ggg ctg ctc ttt      242
Gly Ala Gly Ala Gly Glu Ala Ala Leu Pro Leu Pro Gly Leu Leu Phe
        65              70              75

ggc gcc ccc gcg ctg ctg ggc ctg gca ctg gtc ctg gcg ctg gtc ctg      290
Gly Ala Pro Ala Leu Leu Gly Leu Ala Leu Val Leu Ala Leu Val Leu
      80              85              90              95

gtg ggt ctg gtg agc tgg agg cgg cga cag cgg cgg ctt cgc ggc gcg      338
Val Gly Leu Val Ser Trp Arg Arg Arg Gln Arg Arg Leu Arg Gly Ala
          100              105              110

tcc tcc gca gag gcc ccc gac gga gac aag gac gcc cca gag ccc ctg      386
Ser Ser Ala Glu Ala Pro Asp Gly Asp Lys Asp Ala Pro Glu Pro Leu
          115              120              125

gac aag gtc atc att ctg tct ccg gga atc tct gat gcc aca gct cct      434
Asp Lys Val Ile Ile Leu Ser Pro Gly Ile Ser Asp Ala Thr Ala Pro
          130              135              140

gcc tgg cct cct cct ggg gaa gac cca gga acc acc cca cct ggc cac      482
Ala Trp Pro Pro Pro Gly Glu Asp Pro Gly Thr Thr Pro Pro Gly His
          145              150              155

agt gtc cct gtg cca gcc aca gag ctg ggc tcc act gaa ctg gtg acc      530
Ser Val Pro Val Pro Ala Thr Glu Leu Gly Ser Thr Glu Leu Val Thr
          160              165              170              175

acc aag acg gcc ggc cct gag caa caa tag cagggagccg gcaggagggtg      580
Thr Lys Thr Ala Gly Pro Glu Gln Gln
          180

gcccctgccc tccctctgga ccccagcca ggggcttgga aatcaaattc agctcttcac      640

tocagcatgc acatgccctc tttctgggac caggctaacc tgcagaagca cagacactac      700

agaccacagc attcagcccc catggagttt ggtgtgcttg cctttggctt cagacctcac      760

catctttgac agcccttgaa ggtggtagcc cagctcctgt tcctgtgcct tcaaaaggct      820

ggggcactat gagtaaaaga ccgcttttaa aatggggaag gcaccattaa gccaaaatga      880

```

-6-

atctgaaaaa agacaaaa

898

<210> 5
 <211> 184
 <212> PRT
 <213> Homo sapiens

<400> 5

Met Arg Arg Gly Pro Arg Ser Leu Arg Gly Arg Asp Ala Pro Ala Pro
 1 5 10 15

Thr Pro Cys Val Pro Ala Glu Cys Phe Asp Leu Leu Val Arg His Cys
 20 25 30

Val Ala Cys Gly Leu Leu Arg Thr Pro Arg Pro Lys Pro Ala Gly Ala
 35 40 45

Ser Ser Pro Ala Pro Arg Thr Ala Leu Gln Pro Gln Glu Ser Val Gly
 50 55 60

Ala Gly Ala Gly Glu Ala Ala Leu Pro Leu Pro Gly Leu Leu Phe Gly
 65 70 75 80

Ala Pro Ala Leu Leu Gly Leu Ala Leu Val Leu Ala Leu Val Leu Val
 85 90 95

Gly Leu Val Ser Trp Arg Arg Arg Gln Arg Arg Leu Arg Gly Ala Ser
 100 105 110

Ser Ala Glu Ala Pro Asp Gly Asp Lys Asp Ala Pro Glu Pro Leu Asp
 115 120 125

Lys Val Ile Ile Leu Ser Pro Gly Ile Ser Asp Ala Thr Ala Pro Ala
 130 135 140

Trp Pro Pro Pro Gly Glu Asp Pro Gly Thr Thr Pro Pro Gly His Ser
 145 150 155 160

Val Pro Val Pro Ala Thr Glu Leu Gly Ser Thr Glu Leu Val Thr Thr
 165 170 175

Lys Thr Ala Gly Pro Glu Gln Gln
 180

<210> 6
 <211> 1377
 <212> DNA
 <213> Homo sapiens

-7-

<220>

<221> CDS

<222> (14) .. (895)

<223>

<400> 6

```

agcatcctga gta atg agt ggc ctg ggc cgg agc agg cga ggt ggc cgg      49
      Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg
      1              5              10

agc cgt gtg gac cag gag gag cgc ttt cca cag ggc ctg tgg acg ggg      97
Ser Arg Val Asp Gln Glu Glu Arg Phe Pro Gln Gly Leu Trp Thr Gly
      15              20              25

gtg gct atg aga tcc tgc ccc gaa gag cag tac tgg gat cct ctg ctg      145
Val Ala Met Arg Ser Cys Pro Glu Glu Gln Tyr Trp Asp Pro Leu Leu
      30              35              40

ggg acc tgc atg tcc tgc aaa acc att tgc aac cat cag agc cag cgc      193
Gly Thr Cys Met Ser Cys Lys Thr Ile Cys Asn His Gln Ser Gln Arg
      45              50              55              60

acc tgt gca gcc ttc tgc agg tca ctc agc tgc cgc aag gag caa ggc      241
Thr Cys Ala Ala Phe Cys Arg Ser Leu Ser Cys Arg Lys Glu Gln Gly
      65              70              75

aag ttc tat gac cat ctc ctg agg gac tgc atc agc tgt gcc tcc atc      289
Lys Phe Tyr Asp His Leu Leu Arg Asp Cys Ile Ser Cys Ala Ser Ile
      80              85              90

tgt gga cag cac cct aag caa tgt gca tac ttc tgt gag aac aag ctc      337
Cys Gly Gln His Pro Lys Gln Cys Ala Tyr Phe Cys Glu Asn Lys Leu
      95              100              105

agg agc cca gtg aac ctt cca cca gag ctc agg aga cag cgg agt gga      385
Arg Ser Pro Val Asn Leu Pro Pro Glu Leu Arg Arg Gln Arg Ser Gly
      110              115              120

gaa gtt gaa aac aat tca gac aac tcg gga agg tac caa gga ttg gag      433
Glu Val Glu Asn Asn Ser Asp Asn Ser Gly Arg Tyr Gln Gly Leu Glu
      125              130              135              140

cac aga ggc tca gaa gca agt cca gct ctc ccg ggg ctg aag ctg agt      481
His Arg Gly Ser Glu Ala Ser Pro Ala Leu Pro Gly Leu Lys Leu Ser
      145              150              155

gca gat cag gtg gcc ctg gtc tac agc acg ctg ggg ctc tgc ctg tgt      529
Ala Asp Gln Val Ala Leu Val Tyr Ser Thr Leu Gly Leu Cys Leu Cys
      160              165              170

gcc gtc ctc tgc tgc ttc ctg gtg gcg gtg gcc tgc ttc ctc aag aag      577
Ala Val Leu Cys Cys Phe Leu Val Ala Val Ala Cys Phe Leu Lys Lys
      175              180              185

agg ggg gat ccc tgc tcc tgc cag ccc cgc tca agg ccc cgt caa agt      625
Arg Gly Asp Pro Cys Ser Cys Gln Pro Arg Ser Arg Pro Arg Gln Ser
      190              195              200

ccg gcc aag tct tcc cag gat cac gcg atg gaa gcc ggc agc cct gtg      673
Pro Ala Lys Ser Ser Gln Asp His Ala Met Glu Ala Gly Ser Pro Val
      205              210              215              220

```

-8-

```

agc aca tcc ccc gag cca gtg gag acc tgc agc ttc tgc ttc cct gag      721
Ser Thr Ser Pro  Glu Pro Val Glu Thr Cys Ser Phe Cys Phe Pro Glu
                225                230                235

tgc agg gcg ccc acg cag gag agc gca gtc acg cct ggg acc ccc gac      769
Cys Arg Ala Pro  Thr Gln Glu Ser Ala Val Thr Pro Gly Thr Pro Asp
                240                245                250

ccc act tgt gct gga agg tgg ggg tgc cac acc agg acc aca gtc ctg      817
Pro Thr Cys Ala Gly Arg Trp Gly Cys His Thr Arg Thr Thr Val Leu
                255                260                265

cag cct tgc cca cac atc cca gac agt ggc ctt ggc att gtg tgt gtg      865
Gln Pro Cys Pro His Ile Pro Asp Ser Gly Leu Gly Ile Val Cys Val
                270                275                280

cct gcc cag gag ggg ggc cca ggt gca taa atgggggtca gggaggga      915
Pro Ala Gln Glu Gly Gly Pro Gly Ala
285                290

ggaggaggga gagagatgga gaggagggga gagagaaaga gaggtgggga gaggggagag      975

agatatgagg agagagagac agaggaggca gaaagggaga gaaacagagg agacagagag    1035

ggagagagag acagagggag agagagacag aggggaagag aggcagagag ggaaagaggg    1095

agagaaggaa agagacaggc agagaaggag agaggcagag agggagagag gcagagaggg    1155

agagaggcag agagacagag agggagagag ggacagagag agatagagca ggaggtcggg    1215

gcactctgag tcccagttcc cagtgcagct gtaggtcgtc atcacctaac cacacgtgca    1275

ataaagtcct cgtgcctgct gtcacagcc cccgagagcc cctcctcctg gagaataaaa    1335

cctttggcag ctgcccttcc tcaaaaaaaaa aaaaaaaaaa aa                      1377

```

<210> 7

<211> 293

<212> PRT

<213> Homo sapiens

<400> 7

```

Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg Ser Arg Val Asp
1          5          10          15

```

```

Gln Glu Glu Arg Phe Pro Gln Gly Leu Trp Thr Gly Val Ala Met Arg
20          25          30

```

```

Ser Cys Pro Glu Glu Gln Tyr Trp Asp Pro Leu Leu Gly Thr Cys Met
35          40          45

```

```

Ser Cys Lys Thr Ile Cys Asn His Gln Ser Gln Arg Thr Cys Ala Ala
50          55          60

```

```

Phe Cys Arg Ser Leu Ser Cys Arg Lys Glu Gln Gly Lys Phe Tyr Asp
65          70          75          80

```

-9-

His Leu Leu Arg Asp Cys Ile Ser Cys Ala Ser Ile Cys Gly Gln His
 85 90 95

Pro Lys Gln Cys Ala Tyr Phe Cys Glu Asn Lys Leu Arg Ser Pro Val
 100 105 110

Asn Leu Pro Pro Glu Leu Arg Arg Gln Arg Ser Gly Glu Val Glu Asn
 115 120 125

Asn Ser Asp Asn Ser Gly Arg Tyr Gln Gly Leu Glu His Arg Gly Ser
 130 135 140

Glu Ala Ser Pro Ala Leu Pro Gly Leu Lys Leu Ser Ala Asp Gln Val
 145 150 155 160

Ala Leu Val Tyr Ser Thr Leu Gly Leu Cys Leu Cys Ala Val Leu Cys
 165 170 175

Cys Phe Leu Val Ala Val Ala Cys Phe Leu Lys Lys Arg Gly Asp Pro
 180 185 190

Cys Ser Cys Gln Pro Arg Ser Arg Pro Arg Gln Ser Pro Ala Lys Ser
 195 200 205

Ser Gln Asp His Ala Met Glu Ala Gly Ser Pro Val Ser Thr Ser Pro
 210 215 220

Glu Pro Val Glu Thr Cys Ser Phe Cys Phe Pro Glu Cys Arg Ala Pro
 225 230 235 240

Thr Gln Glu Ser Ala Val Thr Pro Gly Thr Pro Asp Pro Thr Cys Ala
 245 250 255

Gly Arg Trp Gly Cys His Thr Arg Thr Thr Val Leu Gln Pro Cys Pro
 260 265 270

His Ile Pro Asp Ser Gly Leu Gly Ile Val Cys Val Pro Ala Gln Glu
 275 280 285

Gly Gly Pro Gly Ala
 290

<210> 8
 <211> 995
 <212> DNA
 <213> Homo sapiens

<220>

<222> (219) . . (773)

<223>

aagactcaaa cttagaaact tgaattagat gtggtattca aatccttacg tgccgcgaag 60

agctgctctt gctgcatttg ctctggaatt cttgtagaga tattacttgt ccttccaggc 180

tggtctttct gtagctccct tgttttcttt ttgtgatc atg ttg cag atg gct ggg 236

cag tgc tcc caa aat gaa tat ttt gac agt ttg ttg cat gct tgc ata 284

cct tgt caa ctt cga tgt tct tct aat act cct cct cta aca tgt caa 332

cgt tat tgt aat gca agt gtg acc aat tca gtg aaa gga acg aat gcg 380

att ctc tgg acc tgt ttg gga ctg agc tta ata att tct ttg qca qtt 428

ttc gtg cta atg ttt ttg cta agg aag ata agc tct gaa cca tta aag 476

gac gag ttt aaa aac aca gga tca ggt ctc ctg ggc atg gct aac att 524

gac ctg gaa aag agc agg act ggt gat gaa att att ctt ccg aga ggc 572

ctc gag tac acg gtg gaa gaa tgc acc tgt gaa gac tgc atc aag agc 620

aaa ccg aag gtc gac tct gac cat tgc ttt cca ctc cca gct atg gaq 668

Lys Pro Lys Val Asp Ser Asp His Cys Phe Pro Leu Pro Ala Met Glu
135 140 145 150

gaa ggc gca acc att ctt gtc acc acg aaa acg aat gac tat tgc aag 716

Glu Gly Ala Thr Ile Leu Val Thr Thr Lys Thr Asn Asp Tyr Cys Lys
155 160 165

agc ctg cca gct gct ttg agt gct acg gag ata gag aaa tca att tct 764

Ser Leu Pro Ala Ala Leu Ser Ala Thr Glu Ile Glu Lys Ser Ile Ser
170 175 180

gct agg taa ttaaccattt cgactcgagc agtgccactt taaaaatctt 813

Ala Arg

-11-

ttgtcagaat agatgatgtg tcagatctct ttaggatgac tgtatttttc agttgccgat 873
 acagctttttt gtcctctaac tgtggaaact ctttatgtta gatataatttc tctagggttac 933
 tgttggggagc ttaatggtag aaacttcctt ggtttcatga ttaaagtctt tttttttcct 993
 ga 995

<210> 9
 <211> 184
 <212> PRT
 <213> Homo sapiens
 <400> 9

Met Leu Gln Met Ala Gly Gln Cys Ser Gln Asn Glu Tyr Phe Asp Ser
 1 5 10 15

Leu Leu His Ala Cys Ile Pro Cys Gln Leu Arg Cys Ser Ser Asn Thr
 20 25 30

Pro Pro Leu Thr Cys Gln Arg Tyr Cys Asn Ala Ser Val Thr Asn Ser
 35 40 45

Val Lys Gly Thr Asn Ala Ile Leu Trp Thr Cys Leu Gly Leu Ser Leu
 50 55 60

Ile Ile Ser Leu Ala Val Phe Val Leu Met Phe Leu Leu Arg Lys Ile
 65 70 75 80

Ser Ser Glu Pro Leu Lys Asp Glu Phe Lys Asn Thr Gly Ser Gly Leu
 85 90 95

Leu Gly Met Ala Asn Ile Asp Leu Glu Lys Ser Arg Thr Gly Asp Glu
 100 105 110

Ile Ile Leu Pro Arg Gly Leu Glu Tyr Thr Val Glu Glu Cys Thr Cys
 115 120 125

Glu Asp Cys Ile Lys Ser Lys Pro Lys Val Asp Ser Asp His Cys Phe
 130 135 140

Pro Leu Pro Ala Met Glu Glu Gly Ala Thr Ile Leu Val Thr Thr Lys
 145 150 155 160

Thr Asn Asp Tyr Cys Lys Ser Leu Pro Ala Ala Leu Ser Ala Thr Glu
 165 170 175

-12-

Ile Glu Lys Ser Ile Ser Ala Arg
180

<210> 10

<211> 1348

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (282)..(1034)

<223>

<400> 10

```

ggtacgaggc ttcctagagg gactggaacc taattctoct gaggctgagg gaggggtggag      60
ggtctcaagg caacgctggc cccacgacgg agtgccagga gcactaacag tacccttagc      120
ttgcttttct cctccctcct ttttattttc aagttccttt ttatttctcc ttgcgtaaca      180
acctttcttcc cttctgcacc actgcccgtc cccttaccgc ccccgccacc tccttgctac      240
cccactcttg aaaccacagc tggtggcagg gtccccagct c atg cca gcc tca tct      296
                                         Met Pro Ala Ser Ser
                                         1           5

cct ttc ttg cta gcc ccc aaa ggg cct cca ggc aac atg ggg ggc cca      344
Pro Phe Leu Leu Ala Pro Lys Gly Pro Pro Gly Asn Met Gly Gly Pro
              10              15              20

gtc aga gag ccg gca ctc tca gtt gcc ctc tgg ttg agt tgg ggg gca      392
Val Arg Glu Pro Ala Leu Ser Val Ala Leu Trp Leu Ser Trp Gly Ala
              25              30              35

gct ctg ggg gcc gtg gct tgt gcc atg gct ctg ctg acc caa caa aca      440
Ala Leu Gly Ala Val Ala Cys Ala Met Ala Leu Leu Thr Gln Gln Thr
              40              45              50

gag ctg cag agc ctc agg aga gag gtg agc cgg ctg cag ggg aca gga      488
Glu Leu Gln Ser Leu Arg Arg Glu Val Ser Arg Leu Gln Gly Thr Gly
              55              60              65

ggc ccc tcc cag aat ggg gaa ggg tat ccc tgg cag agt ctc ccg gag      536
Gly Pro Ser Gln Asn Gly Glu Gly Tyr Pro Trp Gln Ser Leu Pro Glu
              70              75              80              85

cag agt tcc gat gcc ctg gaa gcc tgg gag aat ggg gag aga tcc cgg      584
Gln Ser Ser Asp Ala Leu Glu Ala Trp Glu Asn Gly Glu Arg Ser Arg
              90              95              100

aaa agg aga gca gtg ctc acc caa aaa cag aag aag cag cac tct gtc      632
Lys Arg Arg Ala Val Leu Thr Gln Lys Gln Lys Lys Gln His Ser Val
              105              110              115

ctg cac ctg gtt ccc att aac gcc acc tcc aag gat gac tcc gat gtg      680
Leu His Leu Val Pro Ile Asn Ala Thr Ser Lys Asp Asp Ser Asp Val
              120              125              130

aca gag gtg atg tgg caa cca gct ctt agg cgt ggg aga ggc cta cag      728
Thr Glu Val Met Trp Gln Pro Ala Leu Arg Arg Gly Arg Gly Leu Gln
              135              140              145

```

-13-

gcc caa gga tat ggt gtc cga atc cag gat gct gga gtt tat ctg ctg 776
 Ala Gln Gly Tyr Gly Val Arg Ile Gln Asp Ala Gly Val Tyr Leu Leu
 150 155 160 165

tat agc cag gtc ctg ttt caa gac gtg act ttc acc atg ggt cag gtg 824
 Tyr Ser Gln Val Leu Phe Gln Asp Val Thr Phe Thr Met Gly Gln Val
 170 175 180

gtg tct cga gaa ggc caa gga agg cag gag act cta ttc cga tgt ata 872
 Val Ser Arg Glu Gly Gln Gly Arg Gln Glu Thr Leu Phe Arg Cys Ile
 185 190 195

aga agt atg ccc tcc cac ccg gac cgg gcc tac aac agc tgc tat agc 920
 Arg Ser Met Pro Ser His Pro Asp Arg Ala Tyr Asn Ser Cys Tyr Ser
 200 205 210

gca ggt gtc ttc cat tta cac caa ggg gat att ctg agt gtc ata att 968
 Ala Gly Val Phe His Leu His Gln Gly Asp Ile Leu Ser Val Ile Ile
 215 220 225

ccc cgg gca agg gcg aaa ctt aac ctc tct cca cat gga acc ttc ctg 1016
 Pro Arg Ala Arg Ala Lys Leu Asn Leu Ser Pro His Gly Thr Phe Leu
 230 235 240 245

ggg ttt gtg aaa ctg tga ttgtgttata aaaagtggct ccagcttgg 1064
 Gly Phe Val Lys Leu
 250

aagaccaggg tgggtacata ctggagacag ccaagagctg agtatataaa ggagaggggaa 1124

tgtgcaggaa cagaggcatc ttcttggtt tggtccccc ttctcactt ttcccttttc 1184

attcccaccc cctagacttt gattttacgg atatcttgct tctgttcccc atggagctcc 1244

gaattcttgc gtgtgtgtag atgaggggag ggggacgggc gccaggcatt gttcagacct 1304

ggtcggggcc cactggaagc atccagaaca gcaccaccat ctta 1348

<210> 11
 <211> 250
 <212> PRT
 <213> Homo sapiens

<400> 11

Met Pro Ala Ser Ser Pro Phe Leu Leu Ala Pro Lys Gly Pro Pro Gly
 1 5 10 15

Asn Met Gly Gly Pro Val Arg Glu Pro Ala Leu Ser Val Ala Leu Trp
 20 25 30

Leu Ser Trp Gly Ala Ala Leu Gly Ala Val Ala Cys Ala Met Ala Leu
 35 40 45

Leu Thr Gln Gln Thr Glu Leu Gln Ser Leu Arg Arg Glu Val Ser Arg
 50 55 60

-14-

Leu Gln Gly Thr Gly Gly Pro Ser Gln Asn Gly Glu Gly Tyr Pro Trp
 65 70 75 80

Gln Ser Leu Pro Glu Gln Ser Ser Asp Ala Leu Glu Ala Trp Glu Asn
 85 90 95

Gly Glu Arg Ser Arg Lys Arg Arg Ala Val Leu Thr Gln Lys Gln Lys
 100 105 110

Lys Gln His Ser Val Leu His Leu Val Pro Ile Asn Ala Thr Ser Lys
 115 120 125

Asp Asp Ser Asp Val Thr Glu Val Met Trp Gln Pro Ala Leu Arg Arg
 130 135 140

Gly Arg Gly Leu Gln Ala Gln Gly Tyr Gly Val Arg Ile Gln Asp Ala
 145 150 155 160

Gly Val Tyr Leu Leu Tyr Ser Gln Val Leu Phe Gln Asp Val Thr Phe
 165 170 175

Thr Met Gly Gln Val Val Ser Arg Glu Gly Gln Gly Arg Gln Glu Thr
 180 185 190

Leu Phe Arg Cys Ile Arg Ser Met Pro Ser His Pro Asp Arg Ala Tyr
 195 200 205

Asn Ser Cys Tyr Ser Ala Gly Val Phe His Leu His Gln Gly Asp Ile
 210 215 220

Leu Ser Val Ile Ile Pro Arg Ala Arg Ala Lys Leu Asn Leu Ser Pro
 225 230 235 240

His Gly Thr Phe Leu Gly Phe Val Lys Leu
 245 250

<210> 12
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> wild-type Shine-Dalgarno box and its adjacent sequence

<400> 12
 tttgtacatg gagaaaataa a

21

<210> 13
 <211> 21

-15-

<212> DNA

<213> Artificial

<220>

<223> Optimized Shine-dalgarno box and adjacent sequence

<400> 13

cacgtaaagg aagtatctca t

21

<210> 14

<211> 4458

<212> DNA

<213> Artificial

<220>

<223> pML124

<400> 14

gaattcaact tctccatact ttggataagg aaatacagac atgaaaaatc tcattgctga	60
gttggttattt aagcttgccc aaaaagaaga agagtcgaat gaactgtgtg cgcaggtaga	120
agctttggag attatcgtca ctgcaatgct tcgcaatatg gcgcaaatg accaacagcg	180
gttgattgat caggtagagg gggcgctgta cgaggtaaag cccgatgccg gcattcctga	240
cgacgatacg gagctgctgc gcgattacgt aaagaagtta ttgaagcatc ctcgtcagta	300
aaaagttaat cttttcaaca gctgtcataa agttgtcacg gccgagactt atagtcgctt	360
tggtttttatt ttttaatgta tttgtaacta gtacgcaagt tcacgtaaag gaagtatctc	420
atatgcagat cttcgtgtaa accggtaccg gcaaaaccat cactctagaa gtggatcctc	480
tacgccggac gcacgtggc cggcatcacc ggcgccacag gtgcggttgc tggcgccat	540
atgcccgaca tcaccgatgg ggaagatcgg gctcgccact tcgggctcat gagcgcttgt	600
ttcggcgctgg gtatggtggc agggcccgctg gccggggggac tggttgggcgc catctccttg	660
catgcacccat tccttgccggc ggcggtgctc aacggcctca acctactact gggctgcttc	720
ctaatgcagg agtcgcataa gggagagcgt cgaccgatgc ccttgagagc cttcaaccca	780
gtcagctcct tcgggtgggc gcggggcatg actatcgtcg ccgcacttat gactgtcttc	840
tttatcatgc aactcgtagg acaggtgccg gcagcgctct gggtcatttt cggcgaggac	900
cgttttcgct ggagcgcgac gatgatcggc ctgtcgcttg cggtatcgg aatcttgca	960
gccctcgctc aagccttcgt cactgggtccc gccaccaaag gtttcggcga gaagcaggcc	1020
attatcgccg gcatggcggc cgacgcgctg ggctacgtct tgctggcggt cgcgacgcga	1080
ggctggatgg ccttccccat tatgattctt ctgccttcgg gcggcatcgg gatgcccgcg	1140
ttgcaggcca tgctgtccag gcaggtagat gacgaccatc agggacagct tcaaggatcg	1200
ctcgcggctc ttaccagcct aacttcgac actggaccgc tgatcgtcac ggcgatttat	1260
gccgcctcgg cgagcacatg gaacggggtg gcatggattg taggcgccgc cctatacctt	1320

-16-

gtctgcctcc	cgcggttgcg	tcgcggtgca	tggagccggg	ccacctcgac	ctgaatggaa	1380
gccggcgcca	cctcgctaac	ggattcacca	ctccaagaat	tggagccaat	caattcttgc	1440
ggagaactgt	gaatgcgcaa	accaaccctt	ggcagaacat	atccatcgcg	tccgccatct	1500
ccagcagccg	cacgcggcgc	atctcgggca	gcgttgggtc	ctggccacgg	gtgcgcattga	1560
tcgtgctcct	gtcgttgagg	acccggctag	gctggcgggg	ttgccttact	ggtttagcaga	1620
atgaatcacc	gatacgcgag	cgaacgtgaa	gcgactgctg	ctgcaaaacg	tctgcgacct	1680
gagcaacaac	atgaatggtc	ttcggtttcc	gtgttttcgta	aagtctggaa	acgcggaagt	1740
cagcgccctg	caccattatg	ttccggatct	gcctcgcagg	atgctgctgg	ctaccctgtg	1800
gaacacctac	atctgtatta	acgaagcgct	ggcattgacc	ctgagtgatt	tttctctggt	1860
cccgcgcgat	ccataccgcc	agttgtttac	cctcacaacg	ttccagtaac	cgggcattgtt	1920
catcatcagt	aacccgatc	gtgagcatcc	tctctcgttt	catcggtatc	attaccccca	1980
tgaacagaaa	tcccccttac	acggaggcat	cagtgcacaa	acaggaaaaa	accgccctta	2040
acatggcccc	ctttatcaga	agccagacat	taacgcttct	ggagaaactc	aacgagctgg	2100
acgcggatga	acaggcagac	atctgtgaat	cgcttcacga	ccacgctgat	gagctttacc	2160
gcagctgcct	cgcgcgtttc	ggtgatgacg	gtgaaaacct	ctgacacatg	cagctcccgg	2220
agacggtcac	agcttgctcg	taagcggatg	ccgggagcag	acaagcccg	cagggcgcgt	2280
cagcgggtgt	tggcgggtgt	cggggcgcag	ccatgaccca	gtcacgtagc	gatagcggag	2340
tgtatactgg	cttaactatg	cggcatcaga	gcagattgta	ctgagagtgc	accatatatg	2400
cgggtgtgaa	taccgcacag	atgcgttaagg	agaaaatacc	gcctcaggcg	ctcttccgct	2460
tcctcgctca	ctgaactcgt	gcgctcggtc	gttcggctgc	ggcgagcgg	atcagctcac	2520
tcaaaggcgg	taatacgggt	atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	2580
gcaaaaggcc	agcaaaaggc	caggaaccgt	aaaaaggccg	cgttgctggc	gtttttccat	2640
aggctccgcc	cccctgacga	gcctcacaaa	aatcgacgct	caagtcagag	gtggcgaaac	2700
ccgacaggac	tataaagata	ccaggcgttt	ccccctggaa	gctccctcgt	gcgctctcct	2760
gttccgaccc	tgccgcttac	cggatacctg	tccgcctttc	tcccttcggg	aagcgtggcg	2820
ctttctcata	gctcacgctg	taggtatctc	agttcgggtg	aggctcgttcg	ctccaagctg	2880
ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctgcg	ccttatccgg	taactatcgt	2940
cttgagtcca	acccggtaag	acacgactta	tcgccactgg	cagcagccac	tggtaacagg	3000
attagcagag	cgaggatatgt	aggcgggtgt	acagagttct	tgaagtgggtg	gcctaactac	3060
ggctacacta	gaaggacagt	atttgggtatc	tcgctctctg	tgaagccagt	taccttcgga	3120
aaaagagttg	gtagctcttg	atccggcaaa	caaaccaccg	ctggtagcgg	tggttttttt	3180

-17-

```

gtttgcaagc agcagattac ggcagaaaa aaaggatctc aagaagatcc tttgatcttt 3240
tctacggggt ctgacgctca gtggaacgaa aactcacgtt aagggatttt ggtcatgaga 3300
ttatcaaaaa ggatcttcac ctagatcctt ttaaattaaa aatgaagttt taaatcaatc 3360
taaagtatat atgagtaaac ttggtctgac agttaccaat gcttaatcag tgaggcacct 3420
atctcagcga tctgtctatt tcgttcaccc atagttgcct gactccccgt cgtgtagata 3480
actacgatac gggaggggctt accatctggc ccagtgctg caatgatacc gcgagaccca 3540
cgctcaccgg ctccagattt atcagcaata aaccagccag ccggaagggc cgagcgcaga 3600
agtggtcctg caactttatc cgctccatc cagtctatta attgttgccg ggaagctaga 3660
gtaagtagtt cgccagttaa tagtttgccg aacgttggtt ccattgctgc aggcacgtg 3720
gtgtcacgct cgtcgtttgg tatggcttca ttcagctccg gttccaacg atcaaggcga 3780
gttacatgat ccccatgtt gtgcaaaaaa gcggttagct ccttcggtcc tccgatcgtt 3840
gtcagaagta agttggccgc agtggttatca ctcatgggta tggcagcact gcataattct 3900
cttactgtca tgccatccgt aagatgcttt tctgtgactg gtgagtactc aaccaagtca 3960
ttctgagaat agtgtatgcy gcgaccgagt tgctcttgcc cggcgtcaac acgggataat 4020
accgcgccac atagcagaac tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga 4080
aaactctcaa ggatcttacc gctgttgaga tccagttcga tgtaaccac tcgtgcaccc 4140
aactgatctt cagcatcttt tactttcacc agcgtttctg ggtgagcaaa aacaggaagg 4200
caaatgccg caaaaaaggg aataagggcg acacggaaat gttgaatact catactcttc 4260
ctttttcaat attattgaag catttatcag ggttattgtc tcatgagcgg atacatattt 4320
gaatgtattt agaaaaataa acaaataggg gttccgcgca catttccccg aaaagtgcc 4380
cctgacgtct aagaaccat tattatcatg acattaacct ataaaaatag gcgtatcacg 4440
aggccctttc gtcttcaa 4458

```

```

<210> 15
<211> 4980
<212> DNA
<213> Artificial

```

```

<220>
<223> pML124:MBPss-BLyS

```

```

<400> 15
gaattcaact tctccatact ttggataagg aaatacagac atgaaaaatc tcattgctga 60
gttgttattt aagcttgccc aaaaagaaga agagtccaat gaactgtgtg cgcaggtaga 120
agctttggag attatcgtca ctgcaatgct tcgcaatatg gcgcaaaatg accaacagcg 180
gttgattgat caggtagagg gggcgtgta cgaggtaaag cccgatgccg gcattcctga 240

```

-18-

cgacgatacg gagctgctgc gcgattacgt aaagaagtta ttgaagcatc ctcgtcagta	300
aaaagttaat cttttcaaca gctgtcataa agttgtcacg gccgagactt atagtcgctt	360
tgtttttatt ttttaatgta tttgtaacta gtacgcaagt tcacgtaaag gaagtatctc	420
atatgaaaat aaaaacaggt gcacgcatcc tcgcattatc cgcattaacg acgatgatgt	480
tttccgcctc ggctctcgcc gctgttcagg gtccggaaga aaccgttact caggactgcc	540
ttcagctgat cgcagactct gaaactccga ccatccagaa aggttcttac acctttgttc	600
cttggtctgct ttctttcaaa cgtgggtctg ccctggaaga gaaagaaaac aaaatcctgg	660
ttaaagaaac tggttacttc tttatctacg gtcagggttct ttacactgat aagacctacg	720
ccatgggtca cctgattcag cgtaagaaag ttcacgtttt cggtgacgag ctgtctctgg	780
ttactctgtt tcgctgcatt cagaacatgc cggaaactct tctaacaac tcctgctact	840
ctgctggcat cgcaaaactg gaagaggggtg atgaactgca gctggcaatt cctcgtgaaa	900
acgcacaaat ttctctggac ggtgatgtaa ccttctttgg tgcactgaaa cttctgtaat	960
aataaggtac cggcaaaacc atcactctag aagtggatcc tctacgccgg acgcatcgtg	1020
gccggcatca ccggcgccac aggtgcgggt gctggcgctt atatcgccga catcaccgat	1080
ggggaagatc gggctcgcca ctccgggctc atgagcgctt gtttcggcgt gggatatgggtg	1140
gcaggccccg tggccggggg actgttgggc gccatctcct tgcattgcacc attccttgcg	1200
gcggcggtgc tcaacggcct caacctacta ctgggctgct tctaatagca ggagtcgcat	1260
aaggagagac gtcgaccgat gcccttgaga gccttcaacc cagtcagctc cttccgggtg	1320
gcgcggggca tgactatcgt cgcgcactt atgactgtct tctttatcat gcaactcgta	1380
ggacaggtgc cggcagcgt ctgggtcatt ttcggcgagg accgctttcg ctggagcgcg	1440
acgatgatcg gcctgtcgct tgcggtatct ggaatcttgc acgccctcgc tcaagccttc	1500
gtcactggtc ccgccaccaa acgtttcggc gagaagcagg ccattatcgc cggcatggcg	1560
gccgacgcgc tgggctacgt cttgctggcg ttcgcgacgc gaggtggat ggccttcccc	1620
attatgattc ttctcgcttc cggcggcatc gggatgcccc cgttgcaggc catgctgtcc	1680
aggcaggtag atgacgacca tcagggacag cttcaaggat cgctcgccgc tcttaccagc	1740
ctaacttcga tcaactggacc gctgatcgtc acggcgatct atgccgcctc ggcgagcaca	1800
tggaaacgggt tggcatggat tgtaggcgc gccctatacc ttgtctgcct ccccgcggtg	1860
cgtcgcgggtg catggagccg ggccacctcg acctgaatgg aagccggcgg cacctcgcta	1920
acggattcac cactccaaga attggagcca atcaattctt gcggagaact gtgaatgcgc	1980
aaaccaaccc ttggcagaac atatccatcg cgtccgccat ctccagcagc cgcacgcggc	2040
gcatctcggg cagcgttggg tcctggccac ggggtgcgcat gatcgtgctc ctgctggtga	2100

-19-

ggacccggct	aggctggcgg	ggttgcccta	ctgggttagca	gaatgaatca	ccgatacgcg	2160
agcgaacgtg	aagcgactgc	tgctgcaaaa	cgtctgcgac	ctgagcaaca	acatgaatgg	2220
tcttcggttt	ccgtgtttcg	taaagtctgg	aaacgcggaa	gtcagcgccc	tgaccatta	2280
tgttccggat	ctgcatcgca	ggatgctgct	ggctaccctg	tggaacacct	acatctgtat	2340
taacgaagcg	ctggcattga	ccctgagtga	tttttctctg	gtcccgcgcg	atccataccg	2400
ccagttgttt	accctcacia	cgttccagta	accgggcatg	ttcatcatca	gtaacccgta	2460
tcgtgagcat	cctctctcgt	ttcatcggtg	tcattacccc	catgaacaga	aatccccctt	2520
acacggaggc	atcagtgacc	aaacaggaaa	aaaccgccct	taacatggcc	cgctttatca	2580
gaagccagac	attaacgctt	ctggagaaa	tcaacgagct	ggacgcggat	gaacaggcag	2640
acatctgtga	atcgcttcac	gaccacgctg	atgagcttta	ccgcagctgc	ctcgcgcgtt	2700
tcggtgatga	cgggtgaaa	ctctgacaca	tgagctccc	ggagacggtc	acagcttgct	2760
tgtaagcggg	tgccgggagc	agacaagccc	gtcagggcgc	gtcagcgggt	gttggcgggt	2820
gtcggggcgc	agccatgacc	cagtcacgta	gcgatagcgg	agtgtatact	ggcttaacta	2880
tgccgcatca	gagcagattg	tactgagagt	gcaccatata	tgccgtgtga	aataccgcac	2940
agatgcgtaa	ggagaaaata	ccgcatcagg	cgctcttccg	cttcctcgtc	cactgactcg	3000
ctgcgctcgg	tcgttcggct	gcggcgagcg	gtatcagctc	actcaaaggc	ggtaatacgg	3060
ttatccacag	aatcagggga	taacgcagga	aagaacatgt	gagcaaaagg	ccagcaaaag	3120
gccaggaacc	gtaaaaaggc	cgcgttgctg	gcgtttttcc	ataggctccg	ccccctgac	3180
gagcatcaca	aaaatcgacg	ctcaagtcag	aggtggcgaa	acccgacagg	actataaaga	3240
taccaggcgt	ttccccctgg	aagctccctc	gtgcgctctc	ctgttccgac	cctgccgctt	3300
accggatacc	tgtccgcctt	tctcccttcg	ggaagcgtgg	cgctttctca	tagctcacgc	3360
tgtaggtatc	tcagttcggt	gtaggtcgtt	cgctccaagc	tgggctgtgt	gcacgaaccc	3420
cccgttcagc	ccgaccgctg	cgccttatcc	ggtaactatc	gtcttgagtc	caacccggta	3480
agacacgact	tatcgccact	ggcagcagcc	actggtaaca	ggattagcag	agcgaggat	3540
gtaggcgggtg	ctacagagtt	cttgaagtgg	tggcctaact	acggctacac	tagaaggaca	3600
gtatttggtg	tctgcgctct	gctgaagcca	gttaacctcg	gaaaaagagt	tggtagctct	3660
tgatccggca	aacaaaccac	cgctggtagc	ggtggttttt	ttgtttgcaa	gcagcagatt	3720
acgcgcagaa	aaaaaggatc	tcaagaagat	cctttgatct	ttctacggg	gtctgacgct	3780
cagtggaaacg	aaaactcacg	ttaagggatt	ttggatcatg	gattatcaaa	aaggatcttc	3840
acctagatcc	ttttaaatta	aaaatgaagt	tttaaatcaa	tctaaagtat	atatgagtaa	3900
acttggctctg	acagttacca	atgcttaatc	agtgaggcac	ctatctcagc	gatctgtcta	3960

-20-

```

tttcgttcat ccatagttgc ctgactcccc gtcgtgtaga taactacgat acgggagggc 4020
ttaccatctg gcccagtgcc tgcaatgata ccgcgagacc cacgctcacc ggctccagat 4080
ttatcagcaa taaaccagcc agccggaagg gccgagcgca gaagtgggtcc tgcaacttta 4140
tccgcctcca tccagtctat taattgttgc cgggaagcta gagtaagtag ttcgccagtt 4200
aatagtttgc gcaacgttgt tgccattgct gcaggcatcg tgggtgtcacg ctcgtcgttt 4260
ggtatggctt cattcagctc cggttcccaa cgatcaaggc gagttacatg atcccccatg 4320
ttgtgcaaaa aagcggtttag ctcccttcggt cctccgatcg ttgtcagaag taagttggcc 4380
gcagtgttat cactcatggg tatggcagca ctgcataatt ctcttactgt catgccatcc 4440
gtaagatgct tttctgtgac tgggtgagtac tcaaccaagt cattctgaga atagtgtatg 4500
cggcgaccga gttgctcttg cccggcgcta acacgggata ataccgcgcc acatagcaga 4560
actttaaaag tgctcatcat tggaaaacgt tcttcggggc gaaaactctc aaggatctta 4620
ccgctgttga gatccagttc gatgtaacct actcgtgcac ccaactgac ttcagcatct 4680
tttactttca ccagcgtttc tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag 4740
ggaataaggg cgacacggaa atgttgaata ctcatctct tcctttttca atattattga 4800
agcattttatc agggttattg tctcatgagc ggatacatat ttgaatgtat ttagaaaaat 4860
aaacaaatag gggttccgcg cacatttccc cgaaaagtgc cacctgacgt ctaagaaacc 4920
attattatca tgacattaac ctataaaaat aggcgtatca cgaggccctt tcgtcttcaa 4980

```

<210> 16
 <211> 26
 <212> PRT
 <213> Artificial

<220>
 <223> MBP signal sequence

<400> 16

```

Met Lys Ile Lys Thr Gly Ala Arg Ile Leu Ala Leu Ser Ala Leu Thr
1           5           10           15

```

```

Thr Met Met Phe Ser Ala Ser Ala Leu Ala
          20           25

```

<210> 17
 <211> 178
 <212> PRT
 <213> Artificial

<220>
 <223> MBPss-Neutrokin-α

<400> 17

-21-

Met Lys Ile Lys Thr Gly Ala Arg Ile Leu Ala Leu Ser Ala Leu Thr
 1 5 10 15

Thr Met Met Phe Ser Ala Ser Ala Leu Ala Ala Val Gln Gly Pro Glu
 20 25 30

Glu Thr Val Thr Gln Asp Cys Leu Gln Leu Ile Ala Asp Ser Glu Thr
 35 40 45

Pro Thr Ile Gln Lys Gly Ser Tyr Thr Phe Val Pro Trp Leu Leu Ser
 50 55 60

Phe Lys Arg Gly Ser Ala Leu Glu Glu Lys Glu Asn Lys Ile Leu Val
 65 70 75 80

Lys Glu Thr Gly Tyr Phe Phe Ile Tyr Gly Gln Val Leu Tyr Thr Asp
 85 90 95

Lys Thr Tyr Ala Met Gly His Leu Ile Gln Arg Lys Lys Val His Val
 100 105 110

Phe Gly Asp Glu Leu Ser Leu Val Thr Leu Phe Arg Cys Ile Gln Asn
 115 120 125

Met Pro Glu Thr Leu Pro Asn Asn Ser Cys Tyr Ser Ala Gly Ile Ala
 130 135 140

Lys Leu Glu Glu Gly Asp Glu Leu Gln Leu Ala Ile Pro Arg Glu Asn
 145 150 155 160

Ala Gln Ile Ser Leu Asp Gly Asp Val Thr Phe Phe Gly Ala Leu Lys
 165 170 175

Leu Leu

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/40979

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 49/00; C07K 14/525

US CL : 424/1.41; 530/351

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/1.41; 530/351; 534/10,11; 540/474

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98/18921 A1 (HUMAN GENOME SCIENCES, INC.) 07 May 1998 (07.05.1998) See Page 45, lines 1-40, and sequence 2.	1-123
Y	US 6,541,224 B2 (YU et al) 01 April 2003 (01.04.2003) , See Column 64, lines 1-40.	1-123
Y	US 5,605,671 A (LYLE et al) 25 February 1997 (25.02.1997), See Column 5, lines 1-30.	1-123
Y	US 5,281,704 A (LOVE et al) 25 January 1994 (25.01.1994) See example 23.	1-123

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"B"	earlier application or patent published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

04 April 2004 (04.04.2004)

Date of mailing of the international search report

07 MAY 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Valerie Bell-Hausfn
Edward C. Ward

Telephone No. 571/272-1600